

# Two Classes of Integral Regular Graphs \*

Ligong Wang<sup>1</sup>, Xueliang Li<sup>2</sup> and Cornelis Hoede<sup>3</sup>

<sup>1</sup>Department of Applied Mathematics,  
Northwestern Polytechnical University, Xi'an, Shaanxi, 710072, P. R.China.  
E-mail: ligongwangnpu@yahoo.com.cn

<sup>2</sup>Center for Combinatorics, Nankai University, Tianjin, 300071, P. R. China.  
E-mail: x.li@eyou.com

<sup>3</sup>Faculty of Mathematical Science, University of Twente,  
P.O.Box 217,7500 AE Enschede, The Netherlands.  
E-mail: hoede@math.utwente.nl

## Abstract

A graph  $G$  is called integral or Laplacian integral if all the eigenvalues of the adjacency matrix  $A(G)$  or the Laplacian matrix  $Lap(G) = D(G) - A(G)$  of  $G$  are integers, where  $D(G)$  denote the diagonal matrix of the vertex degrees of  $G$ . Let  $K_{n,n+1} \equiv K_{n+1,n}$  and  $K_{1,p}[(p-1)K_p]$  denote the  $(n+1)$ -regular graph with  $4n+2$  vertices and the  $p$ -regular graph with  $p^2+1$  vertices, respectively. In this paper, we shall give the spectra and characteristic polynomials of  $K_{n,n+1} \equiv K_{n+1,n}$  and  $K_{1,p}[(p-1)K_p]$  from the theory on matrices. We derive the characteristic polynomials for their complement graphs, their line graphs, the complement graphs of their line graphs and the line graphs of their complement graphs. We also obtain the numbers of spanning trees for such graphs. When  $p = n^2 + n + 1$ , these graphs are not only integral but also Laplacian integral. The discovery of these integral graphs is a new contribution to the search of integral graphs.

**Key Words:** Integral graph, Regular graph, Block circulant matrix, Graph spectrum, Laplacian integral.

**AMS Subject Classification (2000):** 05C05, 11D09, 11D41.

---

\*Supported by National Science Foundation of China (No. 2000CG0103), the Fund of "The Developing Program for Outstanding Persons" in NPU and S & T Innovative Foundation for Young Teachers of Northwestern Polytechnical University.

The research of this paper was partially done while the first author was visiting the University of Twente during the period from November 25, 2001 to December 22, 2001.

# I. Introduction

We use  $G$  to denote a simple graph with vertex set  $V(G) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E(G)$ . The adjacency matrix  $A = A(G) = [a_{ij}]$  of  $G$  is an  $n \times n$  symmetric matrix of 0's and 1's with  $a_{ij} = 1$  if and only if  $v_i$  and  $v_j$  are joined by an edge. The characteristic polynomial of  $G$  is the polynomial  $P(G) = P(G, x) = \det(xI_n - A)$ , where and in the sequel  $I_n$  always denotes the  $n \times n$  identity matrix. The spectrum of  $A(G)$  is also called the spectrum of  $G$ . If the eigenvalues are ordered by  $\lambda_1 > \lambda_2 > \dots > \lambda_r$ , and their multiplicities are  $m_1, m_2, \dots, m_r$ , respectively, then we shall write

$$\text{Spec}(G) = \begin{pmatrix} \lambda_1 & \lambda_2 & \dots & \lambda_r \\ m_1 & m_2 & \dots & m_r \end{pmatrix}.$$

Let  $D(G) = \text{diag}(d(v_1), d(v_2), \dots, d(v_n))$  be the diagonal matrix of the vertex degrees of  $G$ . Then  $\text{Lap}(G) = D(G) - A(G)$  is called the *Laplacian matrix* of  $G$ . Clearly,  $\text{Lap}(G)$  is a real symmetric matrix. If all the eigenvalues of the Laplacian matrix  $\text{Lap}(G)$  of  $G$  are integers, we say that  $G$  is *Laplacian integral*.

The notion of integral graphs was first introduced by F. Harary and A.J. Schwenk in 1974 (see [1]). A graph  $G$  is called *integral* if all the zeros of the characteristic polynomial  $P(G, x)$  of  $G$  are integers. In general, the problem of characterizing integral graphs seems to be very difficult. Thus, it makes sense to restrict our investigations to some interesting families of graphs, for instance, cubic graphs [2, 3], complete  $r$ -partite graphs [4, 24], graphs with three eigenvalues [5], graphs with maximum degree 4 [6], etc. Trees present another important family of graphs for which the problem has been considered in [7-20]. Some graph operations, which when applied on integral graphs produce again integral graphs, are described in [1] or [22]. Other results on integral graphs can be found in [21-23, 28, 34]. For all other facts or terminology on graph spectra, see [22, 23].

For a graph  $G$ , let  $\overline{G}$  be the complement graph of  $G$  and  $L(G)$  denote the *line graph* of  $G$ , in which  $V(L(G)) = E(G)$ , and where two vertices are adjacent if and only if they are adjacent as edges of  $G$ . The  $m$ -iterated line graph of  $G$  is defined recursively by  $L^0(G) = G$  and  $L^m(G) = L(L^{m-1}(G))$ . A graph is said to be *regular of degree  $k$  (or  $k$ -regular)* if each of its vertices has degree  $k$ . Denote  $\kappa(G)$  the number of spanning trees in a graph  $G$ . We denote by  $G_1 \cup G_2$  the union of two disjoint graphs  $G_1$  and  $G_2$ , and by  $nG$  the disjoint union of  $n$  copies of  $G$ . A complete bipartite graph  $K_{p_1, p_2}$  is a graph with vertex classes  $V_1$  and  $V_2$  if  $V = V_1 \cup V_2$ ,  $V_1 \cap V_2 = \emptyset$ , where  $V_i$  are nonempty disjoint sets,  $|V_i| = p_i$  for  $i = 1, 2$ , such that two vertices in  $V$  are adjacent if and only if they belong to different classes. The  $(n+1)$ -regular graph  $K_{n, n+1} \equiv K_{n+1, n}$  on  $4n+2$  vertices is obtained by adding the edges  $\{v_i w_i | i = 1, 2, \dots, n+1\}$  from two disjoint copies of  $K_{n, n+1}$  with vertex classes  $V_1 = \{u_i | i = 1, 2, \dots, n\}$ ,  $V_2 = \{v_i | i = 1, 2, \dots, n+1\}$  and  $U_1 = \{z_i | i = 1, 2, \dots, n\}$ ,  $U_2 = \{w_i | i = 1, 2, \dots, n+1\}$ , respectively. Let  $K_{1, p}$  be a graph with vertex classes  $V_1 = \{u_1\}$  and  $V_2 = \{v_i | i = 1, 2, \dots, p\}$ . The  $i$ -th graph

$K_p$  of  $(p-1)K_p$  has the vertex set  $\{w_{ij} | j = 1, 2, \dots, p\}$ , where  $i = 1, 2, \dots, p-1$ . Then the  $p$ -regular graph  $K_{1,p}[(p-1)K_p]$  on  $p^2 + 1$  vertices is obtained by adding the edges  $\{v_i w_{ji} | j = 1, 2, \dots, p-1\}$  for  $i = 1, 2, \dots, p$  between the graph  $K_{1,p}$  and the graph  $(p-1)K_p$ . In this paper, we shall give the spectra and characteristic polynomials of  $K_{n,n+1} \equiv K_{n+1,n}$  and  $K_{1,p}[(p-1)K_p]$  from the theory on matrices. We derive the characteristic polynomials for their complement graphs, their line graphs, the complement graphs of their line graphs and the line graphs of their complement graphs. We also obtain the numbers of spanning trees for such graphs. When  $p = n^2 + n + 1$ , these graphs are not only integral but also Laplacian integral. The discovery of these integral graphs is a new contribution to the search of integral graphs.

## II. Preliminaries

In this section, we shall give some useful properties of circulant matrices.

First of all, we give the following notations.

- (1)  $C$  and  $R$  denote the set of complex and real numbers, respectively.
- (2)  $C^{m \times n}$  and  $R^{m \times n}$  denote the set of  $m \times n$  matrices whose entries are in  $C$  and  $R$ , respectively.
- (3)  $A^T$  denotes the transpose of the matrix  $A$ .
- (4)  $A^*$  denotes the conjugate transpose of the matrix  $A$ .
- (5)  $J_{m \times n}$  and  $0_{m \times n}$  denotes the  $m \times n$  all 1 and all 0 matrix, respectively.

All other notations and terminology on matrices can be found in [25,32].

Let  $A \in \mathcal{BC}(m, r)$  be a block circulant matrix given as follows

$$A = \begin{bmatrix} A_0 & A_1 & \cdots & A_{m-1} \\ A_{m-1} & A_0 & \cdots & A_{m-2} \\ \vdots & \vdots & \ddots & \vdots \\ A_1 & A_2 & \cdots & A_0 \end{bmatrix},$$

where  $A_k \in R^{r \times r}$ ,  $k = 0, 1, \dots, m-1$ .

Obviously,  $A$  can be expressed as

$$A = \sum_{k=0}^{m-1} \Pi_m^k \otimes A_k, \quad (1)$$

where  $\Pi_m \in R^{m \times m}$  is the permutation matrix

$$\Pi_m = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \end{bmatrix}, \quad (2)$$

and  $\otimes$  denotes the Kronecker product.

We can easily obtain that the characteristic polynomial of  $\Pi_m$  is  $|xI_m - \Pi_m| = x^m - 1$ . Let  $\omega_m = \exp(\frac{2\pi i}{m}) = \cos\frac{2\pi}{m} + i\sin\frac{2\pi}{m}$ , where  $i = \sqrt{-1}$ . Then the eigenvalues of  $\Pi_m$  are  $1, \omega_m, \omega_m^2, \dots, \omega_m^{m-1}$ . Note that the sequence  $\omega_m^k$  ( $k = 0, 1, \dots$ ) is periodic.

Let  $F_l \in C^{l \times l}$  be a matrix as follow

$$F_l = \frac{1}{\sqrt{l}}(\omega_l^{(t-1)(s-1)}) = \frac{1}{\sqrt{l}} \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & \omega_l & \omega_l^2 & \cdots & \omega_l^{l-1} \\ 1 & \omega_l^2 & \omega_l^4 & \cdots & \omega_l^{2(l-1)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \omega_l^{l-1} & \omega_l^{2(l-1)} & \cdots & \omega_l^{(l-1)(l-1)} \end{bmatrix}, \quad (3)$$

where  $\omega_l = \exp(\frac{2\pi i}{l})$ . Obviously,  $F_l$  is a unitary matrix.

The following Lemmas 1 and 2 can be found in [25, 26].

**Lemma 1.** Let  $A \in \mathcal{BC}(m, r)$  be symmetric, then  $A$  is unitarily similar to an Hermitian block diagonal matrix, i.e.,  $A$  is of the form

$$A = (F_m \otimes I_r) \text{diag}(\widetilde{M}_0, \widetilde{M}_1, \dots, \widetilde{M}_{m-1}) (F_m \otimes I_r)^*,$$

where  $\widetilde{M}_j \in C^{r \times r}$ ,  $j = 0, 1, \dots, m-1$ , are given as follows

(1) For  $m \geq 2$  even ,

$$\widetilde{M}_j = A_0 + \sum_{k=1}^{m/2-1} (\omega_m^{kj} A_k + \varpi_m^{kj} A_k^T) + (-1)^j A_{m/2}.$$

(2) For  $m \geq 3$  odd ,

$$\widetilde{M}_j = A_0 + \sum_{k=1}^{(m-1)/2} (\omega_m^{kj} A_k + \varpi_m^{kj} A_k^T).$$

**Lemma 2.** Let  $A \in \mathcal{BC}(m, r)$  be symmetric, then we have that

(1) For  $m \geq 2$  even , all the eigenvalues  $x$  of  $A$  are given by

$$\begin{aligned} & \{x \mid AX = xX, X \in C^{mr}\} \\ & = \{x \mid [A_0 + \sum_{k=1}^{m/2-1} (A_k + A_k^T) + A_{m/2}]Y = xY, Y \in C^r\} \\ & \cup \{x \mid [A_0 + \sum_{k=1}^{m/2-1} (\omega_m^k A_k + \varpi_m^k A_k^T) - A_{m/2}]Z = xZ, Z \in C^r\} \cup \dots \\ & \cup \{x \mid [A_0 + \sum_{k=1}^{m/2-1} (\omega_m^{k(m-1)} A_k + \varpi_m^{k(m-1)} A_k^T) + (-1)^{m-1} A_{m/2}]W = xW, \\ & W \in C^r\}. \end{aligned}$$

(2) For  $m \geq 3$  odd, all the eigenvalues  $x$  of  $A$  are given by

$$\begin{aligned} & \{x \mid AX = xX, X \in C^{mr}\} \\ & = \{x \mid [A_0 + \sum_{k=1}^{(m-1)/2} (A_k + A_k^T)]Y = xY, Y \in C^r\} \\ & \cup \{x \mid [A_0 + \sum_{k=1}^{(m-1)/2} (\omega_m^k A_k + \varpi_m^k A_k^T)]Z = xZ, Z \in C^r\} \cup \dots \\ & \cup \{x \mid [A_0 + \sum_{k=1}^{(m-1)/2} (\omega_m^{k(m-1)} A_k + \varpi_m^{k(m-1)} A_k^T)]W = xW, W \in C^r\}. \end{aligned}$$

**Lemma 3.** Let  $A \in \mathcal{BC}(2, r)$  be symmetric, then the eigenvalues of  $A$  are those of  $A_0 + A_1$  together with those of  $A_0 - A_1$ .

**Proof.** It is easy to check the correctness by Lemma 2.  $\square$

### III. The Characteristic Polynomials of Two classes of Regular Graphs

In this section, we shall determine the characteristic polynomials of  $K_{n,n+1} \equiv K_{n+1,n}$  and  $K_{1,p}[(p-1)K_p]$  from the theory on matrices.

**Theorem 1.** For the regular graph  $K_{n,n+1} \equiv K_{n+1,n}$  of degree  $(n+1)$  with  $4n+2$  vertices, its characteristic polynomial is

$$P(K_{n,n+1} \equiv K_{n+1,n}, x) = (x+n+1)(x+n)(x+1)^n x^{2n-2} (x-1)^n (x-n)(x-n-1).$$

**Proof.** By properly ordering the vertices of the graph  $K_{n,n+1} \equiv K_{n+1,n}$ , the adjacency matrix  $A = A(K_{n,n+1} \equiv K_{n+1,n})$  of  $K_{n,n+1} \equiv K_{n+1,n}$  can be written as the  $(4n+2) \times (4n+2)$  symmetric block circulant matrix such that  $A = A(K_{n,n+1} \equiv K_{n+1,n}) \in \mathcal{BC}(2, 2n+1)$  and

$$A = A(K_{n,n+1} \equiv K_{n+1,n}) = \begin{bmatrix} A_0 & A_1 \\ A_1 & A_0 \end{bmatrix},$$

$$\text{where } A_0 = \begin{bmatrix} 0_{n \times n} & J_{n \times (n+1)} \\ J_{(n+1) \times n} & 0_{(n+1) \times (n+1)} \end{bmatrix} \text{ and } A_1 = \begin{bmatrix} 0_{n \times n} & 0_{n \times (n+1)} \\ 0_{(n+1) \times n} & I_{n+1} \end{bmatrix}.$$

From Lemma 3, we distinguish the following two cases.

**Case 1.** Let  $b_0 = |xI_{2n+1} - (A_0 + A_1)|$ . Then we have

$$b_0 = \begin{vmatrix} xI_n & -J_{n \times (n+1)} \\ -J_{(n+1) \times n} & (x-1)I_{n+1} \end{vmatrix}$$

By careful calculation, we can obtain that  $b_0 = x^{n-1}(x-1)^n(x+n)(x-n-1)$ .

**Case 2.** Let  $b_1 = |xI_{2n+1} - (A_0 - A_1)|$ . Then we have

$$b_1 = \begin{vmatrix} xI_n & -J_{n \times (n+1)} \\ -J_{(n+1) \times n} & (x+1)I_{n+1} \end{vmatrix}$$

By careful calculation, we can obtain that  $b_1 = x^{n-1}(x+1)^n(x-n)(x+n+1)$ .

Hence, the characteristic polynomial of  $K_{n,n+1} \equiv K_{n+1,n}$  is

$$P(K_{n,n+1} \equiv K_{n+1,n}, x) = (x+n+1)(x+n)(x+1)^n x^{2n-2} (x-1)^n (x-n)(x-n-1).$$

The proof is complete.  $\square$

We note that the graph  $K_{1,2} \equiv K_{2,1}$  is the cycle  $C_6$  and the graph  $K_{2,3} \equiv K_{3,2}$  is the graph 3.20 of [22, P.293] or  $G_{10}$  of [3].

**Theorem 2.** For the regular graph  $K_{1,p}[(p-1)K_p]$  of degree  $p$  with  $p^2 + 1$  vertices, its characteristic polynomial is

$$P(K_{1,p}[(p-1)K_p], x) = (x+1)^{(p-1)(p-2)}(x-p+1)^{p-2}(x-p)(x^2+x-p+1)^p.$$

**Proof.** By properly ordering the vertices of the graph  $K_{1,p}[(p-1)K_p]$ , the adjacency matrix  $A = A(K_{1,p}[(p-1)K_p])$  of  $K_{1,p}[(p-1)K_p]$  can be written as the  $(p^2 + 1) \times (p^2 + 1)$  matrix such that

$$A = A(K_{1,p}[(p-1)K_p]) = \begin{bmatrix} A_1 & A_2 & A_3 & \cdots & A_{p-2} & A_{p-1} & I_p & 0_{p \times 1} \\ A_{p-1} & A_1 & A_2 & \cdots & A_{p-3} & A_{p-2} & I_p & 0_{p \times 1} \\ A_{p-2} & A_{p-1} & A_1 & \cdots & A_{p-4} & A_{p-3} & I_p & 0_{p \times 1} \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots & \vdots & \vdots \\ A_3 & A_4 & A_5 & \cdots & A_1 & A_2 & I_p & 0_{p \times 1} \\ A_2 & A_3 & A_4 & \cdots & A_{p-1} & A_1 & I_p & 0_{p \times 1} \\ I_p & I_p & I_p & \cdots & I_p & I_p & 0_{p \times p} & J_{p \times 1} \\ 0_{1 \times p} & 0_{1 \times p} & 0_{1 \times p} & \cdots & 0_{1 \times p} & 0_{1 \times p} & J_{1 \times p} & 0 \end{bmatrix},$$

where  $A_1 = J_{p \times p} - I_p$  and  $A_2 = A_3 = \cdots = A_{p-1} = 0_{p \times p}$ .

Then we have

$$P(K_{1,p}[(p-1)K_p], x) = |xI_{p^2+1} - A(K_{1,p}[(p-1)K_p])| = \begin{vmatrix} (x+1)I_p - J_{p \times p} & 0_{p \times p} & \cdots & 0_{p \times p} & -I_p & 0_{p \times 1} \\ 0_{p \times p} & (x+1)I_p - J_{p \times p} & \cdots & 0_{p \times p} & -I_p & 0_{p \times 1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0_{p \times p} & 0_{p \times p} & \cdots & (x+1)I_p - J_{p \times p} & I_p & 0_{p \times 1} \\ -I_p & -I_p & \cdots & -I_p & xI_p & -J_{p \times 1} \\ 0_{1 \times p} & 0_{1 \times p} & \cdots & 0_{1 \times p} & -J_{1 \times p} & x \end{vmatrix}$$

By careful calculation, we can obtain that the characteristic polynomial of  $K_{1,p}[(p-1)K_p]$  is

$$P(K_{1,p}[(p-1)K_p], x) = (x+1)^{(p-1)(p-2)}(x-p+1)^{p-2}(x-p)(x^2+x-p+1)^p.$$

The proof is complete.  $\square$

We note that the graph  $K_{1,3}[2K_3]$  is the graph 3.16 of [22, P.293] or  $G_{11}$  of [3].

## IV. Other Results

In this section, we shall give the characteristic polynomials for  $\overline{K_{n,n+1} \equiv K_{n+1,n}}$ ,  $L(K_{n,n+1} \equiv K_{n+1,n})$ ,  $\overline{L(K_{n,n+1} \equiv K_{n+1,n})}$ ,  $L(\overline{K_{n,n+1} \equiv K_{n+1,n}})$ ,  $\overline{K_{1,p}[(p-1)K_p]}$ ,  $L(K_{1,p}[(p-1)K_p])$ ,  $\overline{L(K_{1,p}[(p-1)K_p])}$  and  $L(\overline{K_{1,p}[(p-1)K_p]})$ . We also obtain the numbers of spanning trees for these graphs, the graph  $K_{n,n+1} \equiv K_{n+1,n}$  and the graph  $K_{1,p}[(p-1)K_p]$ . For integers  $n \geq 0$  and  $m \geq 0$ , if a regular graph  $G$  is integral, then the graphs  $L^m(G)$  and  $L^n(\overline{L^m(G)})$  are not only integral but also Laplacian integral. We note some interesting characteristic polynomials of integral graphs, see [1-24, 28] and [34].

The following Lemmas 4, 5, 6, 7 and 8 can be found in [27, 28].

**Lemma 4.** If  $G$  is a regular graph of degree  $k$ , then its line graph  $L(G)$  is regular of degree  $2k - 2$ .

**Lemma 5.** If  $G$  is a regular graph of degree  $k$  with  $n$  vertices and  $m = \frac{1}{2}nk$  edges, then

$$P(\overline{G}, x) = (-1)^n \frac{x - n + k + 1}{x + k + 1} P(G, -x - 1).$$

**Lemma 6.** If  $G$  is a regular graph of degree  $k$  with  $n$  vertices and  $m = \frac{1}{2}nk$  edges, then

$$P(L(G), x) = (x + 2)^{\frac{1}{2}n(k-2)} P(G, x + 2 - k).$$

**Lemma 7.** If  $G$  is a regular graph of degree  $k$  with  $n$  vertices and  $m = \frac{1}{2}nk$  edges, then the number of spanning trees  $\kappa(G)$  of  $G$  is given by

$$\kappa(G) = \frac{1}{n} P'(G, x)|_{x=\lambda_1} = \frac{1}{n} \prod_{i=2}^n (\lambda_1 - \lambda_i),$$

where  $\lambda_i$  ( $1 \leq i \leq n$ ) are the roots of  $P(G, x)$  and  $\lambda_1 = k$ .

**Lemma 8.** If  $G$  is a regular graph of degree  $k$  with  $n$  vertices and  $m = \frac{1}{2}nk$  edges, then the number of spanning trees of  $L(G)$  is given by

$$\kappa(L(G)) = 2^{m-n+1} k^{m-n-1} \kappa(G).$$

**Theorem 3.** For the complement of the regular graph  $K_{n,n+1} \equiv K_{n+1,n}$ , the characteristic polynomial of  $\overline{K_{n,n+1} \equiv K_{n+1,n}}$  is

$$P(\overline{K_{n,n+1} \equiv K_{n+1,n}}, x) = (x + n + 1)(x + 2)^n (x + 1)^{2n-2} x^n (x - n + 1)(x - n)(x - 3n).$$

**Proof.** It is easy to check the correctness by Theorem 1 and Lemmas 4 and 5.  $\square$

**Theorem 4.** For the line graph, the complement of the line graph and the line graph of the complement of the regular graph  $K_{n,n+1} \equiv K_{n+1,n}$ , we have the following results.

(1) The characteristic polynomial of  $L(K_{n,n+1} \equiv K_{n+1,n})$  is

$$P[L(K_{n,n+1} \equiv K_{n+1,n}), x] = (x+2)^{n(2n-1)}(x+1)(x-n+2)^n(x-n+1)^{2n-2} \cdot (x-n)^n(x-2n+1)(x-2n).$$

(2) The characteristic polynomial of  $\overline{L(K_{n,n+1} \equiv K_{n+1,n})}$  is

$$P[\overline{L(K_{n,n+1} \equiv K_{n+1,n})}, x] = (x+2n)(x+n+1)^n(x+n)^{2n-2}(x+n-1)^n \cdot x(x-1)^{n(2n-1)}(x-2n^2-n).$$

(3) The characteristic polynomial of  $L(\overline{K_{n,n+1} \equiv K_{n+1,n}})$  is

$$P[L(\overline{K_{n,n+1} \equiv K_{n+1,n}}), x] = (x+2)^{(2n+1)(3n-2)}(x-2n+3)(x-3n+4)^n \cdot (x-3n+3)^{2n-2}(x-3n+2)^n(x-4n+3)(x-4n+2)(x-6n+2).$$

**Proof.** It is easy to check the correctness by Theorems 1, 3 and Lemmas 4, 5, 6.  $\square$

**Corollary 1.** The graphs  $K_{n,n+1} \equiv K_{n+1,n}$ ,  $\overline{K_{n,n+1} \equiv K_{n+1,n}}$ ,  $L(K_{n,n+1} \equiv K_{n+1,n})$ ,  $\overline{L(K_{n,n+1} \equiv K_{n+1,n})}$  and  $L(\overline{K_{n,n+1} \equiv K_{n+1,n}})$  are integral.

**Proof.** It is easy to check the correctness by Theorems 1, 3 and 4.  $\square$

**Theorem 5.** For the numbers of spanning trees in the graphs  $\overline{K_{n,n+1} \equiv K_{n+1,n}}$ ,  $K_{n,n+1} \equiv K_{n+1,n}$ ,  $L(K_{n,n+1} \equiv K_{n+1,n})$ ,  $\overline{L(K_{n,n+1} \equiv K_{n+1,n})}$  and  $L(\overline{K_{n,n+1} \equiv K_{n+1,n}})$ , we have the the following results.

- (1)  $\kappa(K_{n,n+1} \equiv K_{n+1,n}) = n^n(n+1)^{2n-1}(n+2)^n.$
- (2)  $\kappa(\overline{K_{n,n+1} \equiv K_{n+1,n}}) = 3^n n^{n+1}(3n+1)^{2n-2}(3n+2)^n(4n+1).$
- (3)  $\kappa(L(K_{n,n+1} \equiv K_{n+1,n})) = 2^{n(2n-1)}n^n(n+1)^{(2n+3)(n-1)}(n+2)^n$
- (4)  $\kappa(\overline{L(K_{n,n+1} \equiv K_{n+1,n})}) = 2^{2n-2}n^{2n}(n+1)^{(2n+3)(n-1)}(2n-1)^n(2n-1)^{2n-1}(2n+3) \cdot (2n^2+2n-1)^n(2n^2+2n+1)^n.$
- (5)  $\kappa(L(\overline{K_{n,n+1} \equiv K_{n+1,n}})) = 2^{(2n-1)(3n+1)}3^{6n^2-3}n^{6n^2-2}(3n+1)^{2n-2}(3n+2)^n \cdot (4n+1).$

**Proof.** It is easy to check the correctness by Theorems 1, 3, 4 and Lemmas 4, 5, 6, 7 and 8.  $\square$

**Theorem 6.** For the complement of the regular graph  $K_{1,p}[(p-1)K_p]$ , the characteristic polynomial of  $\overline{K_{1,p}[(p-1)K_p]}$  is

$$P[\overline{K_{1,p}[(p-1)K_p]}, x] = x^{(p-1)(p-2)}(x+p)^{p-2}(x-p^2+p)(x^2+x-p+1)^p.$$

**Proof.** It is easy to check the correctness by Theorem 2 and Lemmas 4 and 5.  $\square$

**Theorem 7.** For the line graph, the complement of the line graph and the line graph of the complement of the regular graph  $K_{1,p}[(p-1)K_p]$ , we have the the following results.



(1) The characteristic polynomial of  $L(K_{1,p}[(p-1)K_p])$  is

$$P[L(K_{1,p}[(p-1)K_p]), x] = (x+2)^{\frac{1}{2}(p-2)(p^2+1)}(x-p+3)^{(p-1)(p-2)}(x-2p+3)^{p-2} \cdot (x-2p+2)[(x-p+2)^2 + (x-p+2) - (p-1)]^p.$$

(2) The characteristic polynomial of  $\overline{L(K_{1,p}[(p-1)K_p])}$  is

$$P[\overline{L(K_{1,p}[(p-1)K_p])}, x] = (x-1)^{\frac{1}{2}(p-2)(p^2+1)}(x+p-2)^{(p-1)(p-2)}(x+2p-2)^{p-2} \cdot [x - \frac{1}{2}(p-1)^2(p+2)][(x+p-1)^2 - (x+p-1) - (p-1)]^p.$$

(3) The characteristic polynomial of  $L(\overline{K_{1,p}[(p-1)K_p]})$  is

$$P[L(\overline{K_{1,p}[(p-1)K_p]}), x] = [(x-p^2+p+2)^2 + (x-p^2+p+2) - (p-1)]^p \cdot (x-p^2+2p+2)^{p-2}(x-2p^2+2p+2)(x+2)^{\frac{1}{2}(p+1)(p-2)(p^2+1)} \cdot (x-p^2+p+2)^{(p-1)(p-2)}.$$

**Proof.** It is easy to check the correctness by Theorems 2, 6 and Lemmas 4, 5, 6.  $\square$

**Corollary 2.** For the regular graphs  $K_{1,p}[(p-1)K_p]$ ,  $\overline{K_{1,p}[(p-1)K_p]}$ ,  $L(K_{1,p}[(p-1)K_p])$ ,  $\overline{L(K_{1,p}[(p-1)K_p])}$  and  $L(\overline{K_{1,p}[(p-1)K_p]})$ , let  $n$  be any positive integer, then any one of these graphs is integral if and only if  $p = n^2 + n + 1$ .

**Proof.** It is easy to check the correctness by Theorems 2, 6 and 7.  $\square$

**Corollary 3.** For the regular graphs  $K_{1,p}[(p-1)K_p]$ , the line graph, the complement of the line graph and the line graph of the complement of the regular graph  $K_{1,p}[(p-1)K_p]$ , let  $p = n^2 + n + 1$  and  $n$  be any positive integer, then we have the following results.

- (1)  $P(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}], x) = (x+1)^{n(n+1)(n^2+n-1)}[x - n(n+1)]^{n^2+n-1} \cdot (x - n^2 - n - 1)(x + n + 1)^{n^2+n+1}(x - n)^{n^2+n+1}.$
- (2)  $P(\overline{K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]}, x) = x^{n(n+1)(n^2+n-1)}(x + n^2 + n + 1)^{n^2+n-1} [x - n(n+1)(n^2 + n + 1)](x + n + 1)^{n^2+n+1}(x - n)^{n^2+n+1}.$
- (3)  $P[L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]), x] = (x+2)^{\frac{1}{2}(n^2+n-1)[(n^2+n+1)^2+1]} \cdot [x - (n+2)(n-1)]^{n(n+1)(n^2+n-1)}(x - 2n^2 - 2n + 1)^{n^2+n-1} \cdot [x - 2n(n+1)](x - n^2 + 2)^{n^2+n+1}(x - n^2 - 2n + 1)^{n^2+n+1}.$
- (4)  $P(\overline{L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}])}, x) = (x-1)^{\frac{1}{2}(n^2+n-1)[(n^2+n+1)^2+1]} \cdot (x + n^2 + n - 1)^{n(n+1)(n^2+n-1)}(x + 2n^2 + 2n)^{n^2+n-1} \cdot [x - \frac{1}{2}n^2(n+1)^2(n^2 + n + 3)](x + n^2 + 2n)^{n^2+n+1}(x + n^2 - 1)^{n^2+n+1}.$
- (5)  $P(L(\overline{K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]}), x) = (x+2)^{\frac{1}{2}(n^2+n-1)(n^2+n+2)[(n^2+n+1)^2+1]} \cdot (x - 2n^4 - 4n^3 - 4n^2 - 2n + 2)[x - (n^2 + n - 1)(n^2 + n + 2)]^{n(n+1)(n^2+n-1)} \cdot (x - n^4 - 2n^3 - n^2 + 3)^{n^2+n-1}(x - n^4 - 2n^3 - 2n^2 + 3)^{n^2+n+1} \cdot (x - n^4 - 2n^3 - 2n^2 - 2n + 2)^{n^2+n+1}.$

**Proof.** It is easy to check the correctness by Theorems 2, 6 and 7.  $\square$

**Theorem 8.** For the numbers of spanning trees in the graphs  $K_{1,p}[(p-1)K_p]$ ,  $\overline{K_{1,p}[(p-1)K_p]}$ ,  $L(K_{1,p}[(p-1)K_p])$ ,  $\overline{L(K_{1,p}[(p-1)K_p])}$  and  $L(\overline{K_{1,p}[(p-1)K_p]})$ , we have the the following results.

- (1)  $\kappa(K_{1,p}[(p-1)K_p]) = (p+1)^{(p-1)(p-2)}(p^2+1)^{p-1}$ .
- (2)  $\kappa(\overline{K_{1,p}[(p-1)K_p]}) = p^{(p+1)(p-2)}(p-1)^{p^2-p+2}(p^2+1)^{p-1}$ .
- (3)  $\kappa(L(K_{1,p}[(p-1)K_p])) = 2^{\frac{1}{2}(p-2)(p^2+1)+1}p^{\frac{1}{2}(p-2)(p^2+1)-1}(p+1)^{(p-1)(p-2)} \cdot (p^2+1)^{p-1}$ .
- (4)  $\kappa(\overline{L(K_{1,p}[(p-1)K_p])}) = (\frac{1}{2})^{\frac{1}{2}(p^3+p-4)}p^{\frac{1}{2}(p^3-2p^2+p-4)}(p^2-3)^{\frac{1}{2}(p-2)(p^2+1)} \cdot (p^3-p-2)^{(p-1)(p-2)}(p-1)^{2p-2}(p^2+p+2)^{p-2}(p^2+1)^{p-1}(p^3+p^2-2p-4)^p$ .
- (5)  $\kappa(L(\overline{K_{1,p}[(p-1)K_p]})) = 2^{\frac{1}{2}p(p^3-p^2-p-1)}p^{\frac{1}{2}(p^4-p^3+p^2-3p-8)}(p-1)^{\frac{1}{2}p(p^3-p^2+p-3)} \cdot (p^2+1)^{p-1}$ .

**Proof.** It is easy to check the correctness by Theorems 2, 6, 7 and Lemmas 4, 5, 6, 7 and 8.  $\square$

The following Lemmas 9 and 10 can be found in [1] or [27, 28].

**Lemma 9.** If a regular graph  $G$  is integral, then so is  $\overline{G}$ .

**Lemma 10.** If a regular graph  $G$  is integral, then so is its line graph  $L(G)$ .

**Theorem 9.** For integers  $n \geq 0$  and  $m \geq 0$ , if a regular graph  $G$  is integral, then the graphs  $L^m(G)$  and  $L^n(\overline{L^m(G)})$  are integral.

**Proof.** It is easy to check the correctness by Lemmas 4, 9 and 10.  $\square$

**Corollary 4.** For integers  $p \geq 0$ ,  $n \geq 1$  and  $m \geq 0$ , the regular graphs  $L^m(K_{n,n+1} \equiv K_{n+1,n})$ ,  $L^p(\overline{L^m(K_{n,n+1} \equiv K_{n+1,n})})$ ,  $L^p(\overline{L^m(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]})$  and  $L^m(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}])$  are integral.

**Proof.** It is easy to check the correctness by Lemma 4 and Corollaries 1, 2, 3 and Theorems 1, 2, 3, 4, 6, 7 and 9.  $\square$

In the remainder of the paper, we shall consider the Laplacian integral graphs on the regular graphs. Mohar [29] argues that, because of its importance in various physical and chemical theories, the spectrum of  $Lap(G) = D(G) - A(G)$  is more natural and important than the more widely studied adjacency spectrum. For background knowledge, see [29, 30, 31]. The characteristic polynomial of  $Lap(G)$  is the polynomial  $\sigma(G) = \sigma(G, \mu) = \det(\mu I_n - Lap(G))$ . Let  $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_n(G)$  (or simple  $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$ ) be all the eigenvalues of the Laplacian matrix  $Lap(G)$  of  $G$ , the multiplicity of  $\mu$  as an eigenvalue of  $Lap(G)$  will be denoted by  $m_G(\mu)$ .

The following Lemmas 11, 12 and 13 can be found in [27, 30].

**Lemma 11.** If  $G$  is a regular graph of degree  $k$  with  $n$  vertices, then

$$\sigma(G, \mu) = (-1)^n P(G, k - \mu).$$

or  $\mu_i(G) = \mu_i = k - x_i$  ( $i = 1, 2, \dots, n$ ), where the  $x_i$ 's are the eigenvalues of  $A(G)$ , ordered in weakly decreasing manner.

**Lemma 12.** Let  $G$  be a graph on  $n$  vertices, then the eigenvalues of the Laplacian matrix  $Lap(\overline{G})$  are

$$\mu_i(\overline{G}) = n - \mu_{n-i}(G) \quad (1 \leq i < n) \text{ and } 0.$$

**Lemma 13.**  $G$  is Laplacian integral if and only if  $\overline{G}$  is Laplacian integral.

**Theorem 10.** For the  $\sigma$ -polynomials of the regular graphs  $K_{n,n+1} \equiv K_{n+1,n}$ ,  $\overline{K_{n,n+1} \equiv K_{n+1,n}}$ ,  $L(K_{n,n+1} \equiv K_{n+1,n})$ ,  $\overline{L(K_{n,n+1} \equiv K_{n+1,n})}$  and  $L(\overline{K_{n,n+1} \equiv K_{n+1,n}})$ , we have the the following results.

(1) The characteristic polynomial of  $Lap(K_{n,n+1} \equiv K_{n+1,n})$  is

$$\sigma[K_{n,n+1} \equiv K_{n+1,n}, x] = x(x-1)(x-n)^n(x-n-1)^{2n-2}(x-n-2)^n(x-2n-1) \cdot (x-2n-2).$$

(2) The characteristic polynomial of  $Lap(\overline{K_{n,n+1} \equiv K_{n+1,n}})$  is

$$\sigma[\overline{K_{n,n+1} \equiv K_{n+1,n}}, x] = x(x-2n)(x-2n-1)(x-3n)^n(x-3n-1)^{2n-2} \cdot (x-3n-2)^n(x-4n-1).$$

(3) The characteristic polynomial of  $Lap(L(K_{n,n+1} \equiv K_{n+1,n}))$  is

$$\sigma[L(K_{n,n+1} \equiv K_{n+1,n}), x] = x(x-1)(x-n)^n(x-n-1)^{2n-2}(x-n-2)^n \cdot (x-2n-1)(x-2n-2)^{n(2n-1)}.$$

(4) The characteristic polynomial of  $Lap(\overline{L(K_{n,n+1} \equiv K_{n+1,n})})$  is

$$\sigma[\overline{L(K_{n,n+1} \equiv K_{n+1,n})}, x] = x(x-2n^2-n+1)^{n(2n-1)}(x-2n^2-n) \cdot (x-2n^2-2n-1)^n(x-2n^2-2n)^{2n-2}(x-2n^2-2n+1)^n(x-2n^2-3n).$$

(5) The characteristic polynomial of  $Lap(L(\overline{K_{n,n+1} \equiv K_{n+1,n}}))$  is

$$\sigma(L(\overline{K_{n,n+1} \equiv K_{n+1,n}}), x) = x(x-2n)(x-2n-1)(x-3n)^n(x-3n-1)^{2n-2} \cdot (x-3n-2)^n(x-4n-1)(x-6n)^{(2n+1)(3n-2)}.$$

**Proof.** It is easy to check the correctness by Theorems 1, 3, 4 and Lemmas 11, 12.  $\square$

**Corollary 5.** The graphs  $K_{n,n+1} \equiv K_{n+1,n}$ ,  $\overline{K_{n,n+1} \equiv K_{n+1,n}}$ ,  $L(K_{n,n+1} \equiv K_{n+1,n})$ ,  $\overline{L(K_{n,n+1} \equiv K_{n+1,n})}$  and  $L(\overline{K_{n,n+1} \equiv K_{n+1,n}})$  are Laplacian integral.

**Proof.** It is easy to check the correctness by Theorem 10 and Lemmas 11, 12, 13.  $\square$

**Theorem 11.** For the  $\sigma$ -polynomials of the graphs  $K_{1,p}[(p-1)K_p]$ ,  $\overline{K_{1,p}[(p-1)K_p]}$ ,  $L(K_{1,p}[(p-1)K_p])$ ,  $\overline{L(K_{1,p}[(p-1)K_p])}$  and  $L(\overline{K_{1,p}[(p-1)K_p]})$ , we have the the following results.

(1) The characteristic polynomial of  $Lap(K_{1,p}[(p-1)K_p])$  is

$$\sigma[K_{1,p}[(p-1)K_p], x] = x(x-1)^{p-2}(x-p-1)^{(p-1)(p-2)}[(x-p)^2 - (x-p) - (p-1)]^p.$$

(2) The characteristic polynomial of  $Lap(\overline{K_{1,p}[(p-1)K_p]})$  is

$$\sigma[\overline{K_{1,p}[(p-1)K_p]}, x] = x(x-p^2+p)^{(p-1)(p-2)}(x-p^2)^{p-2} \cdot [(x-p^2+p)^2 - (x-p^2+p) - (p-1)]^p.$$

(3) The characteristic polynomial of  $Lap(L(K_{1,p}[(p-1)K_p]))$  is

$$\sigma[L(K_{1,p}[(p-1)K_p]), x] = x(x-p-1)^{(p-1)(p-2)}(x-1)^{p-2}(x-2p)^{\frac{1}{2}(p-2)(p^2+1)} \cdot [(x-p)^2 - (x-p) - (p-1)]^p.$$

(4) The characteristic polynomial of  $Lap(\overline{L(K_{1,p}[(p-1)K_p])})$  is

$$\sigma(\overline{L(K_{1,p}[(p-1)K_p])}, x) = x[x - \frac{1}{2}p(p^2-3)]^{\frac{1}{2}(p-2)(p^2+1)}[x - \frac{1}{2}(p^3-p-2)]^{(p-1)(p-2)} \cdot \{[x - \frac{1}{2}p(p-1)(p+1)]^2 - [x - \frac{1}{2}p(p-1)(p+1)] - (p-1)\}^p \cdot [x - \frac{1}{2}(p-1)(p^2+p+2)]^{p-2}$$

(5) The characteristic polynomial of  $Lap(L(\overline{K_{1,p}[(p-1)K_p]}))$  is

$$\sigma(L(\overline{K_{1,p}[(p-1)K_p]}), x) = x(x-p^2+p)^{(p-1)(p-2)}(x-p^2)^{p-2} \cdot (x-2p^2+2p)^{\frac{1}{2}(p+1)(p-2)(p^2+1)}[(x-p^2+p)^2 - (x-p^2+p) - (p-1)]^p.$$

**Proof.** It is easy to check the correctness by Theorems 2, 6, 7 and Lemmas 11, 12.  $\square$

**Corollary 6.** Any one of the graphs  $K_{1,p}[(p-1)K_p]$ ,  $\overline{K_{1,p}[(p-1)K_p]}$ ,  $L(K_{1,p}[(p-1)K_p])$ ,  $\overline{L(K_{1,p}[(p-1)K_p])}$  and  $L(\overline{K_{1,p}[(p-1)K_p]})$  is Laplacian integral if and only if  $p = n^2 + n + 1$ .

**Proof.** It is easy to check the correctness by Theorems 2, 6, 7, 11 and Lemmas 11, 12, 13.  $\square$

**Corollary 7.** For the  $\sigma$ -polynomials of the graphs  $K_{1,p}[(p-1)K_p]$ ,  $\overline{K_{1,p}[(p-1)K_p]}$ ,  $L(K_{1,p}[(p-1)K_p])$ ,  $\overline{L(K_{1,p}[(p-1)K_p])}$  and  $L(\overline{K_{1,p}[(p-1)K_p]})$ , if  $p = n^2 + n + 1$ , then we have the the following results.

(1) The characteristic polynomial of  $Lap(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}])$  is

$$\sigma[K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}], x] = x(x-1)^{n^2+n-1}(x-n^2-n-2)^{n(n+1)(n^2+n-1)} \cdot (x-n^2-1)^{n^2+n+1}(x-n^2-2n-2)^{n^2+n+1}.$$

(2) The characteristic polynomial of  $Lap(\overline{K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]})$  is

$$\sigma[\overline{K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]}, x] = x[x - n(n+1)(n^2+n+1)]^{n(n+1)(n^2+n-1)} \cdot [x - (n^2+n+1)^2]^{n^2+n-1}[x - (n+1)^2(n^2+1)]^{n^2+n+1} \cdot [x - n^2(n^2+2n+2)]^{n^2+n+1}.$$

(3) The characteristic polynomial of  $Lap(L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]))$  is

$$\sigma[L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]), x] = x(x-n^2-n-2)^{n(n+1)(n^2+n-1)} \cdot (x-1)^{n^2+n-1}[x - 2(n^2+n+1)]^{\frac{1}{2}(n^2+n-1)[(n^2+n+1)^2+1]}(x-n^2-2n-2)^{n^2+n+1} \cdot (x-n^2-1)^{n^2+n+1}.$$

(4) The characteristic polynomial of  $Lap(\overline{L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]})$  is

$$\begin{aligned} \sigma(\overline{L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]}, x) &= x\{x - \frac{1}{2}[(n^2 + n + 1)^3 \\ &- (n^2 + n + 3)]\}^{n(n+1)(n^2+n+1)} \cdot [x - \frac{1}{2}n(n+1)(n^4 + 2n^3 + 4n^2 + 3n + 3)]^{n^2+n-1} \\ &\cdot \{x - \frac{1}{2}(n+1)[n(n^2 + n + 1)(n^2 + n + 2) - 2]\}^{n^2+n+1} \\ &\cdot \{x - \frac{1}{2}n[(n+1)(n^2 + n + 1)(n^2 + n + 2) + 2]\}^{n^2+n+1} \\ &\cdot \{x - \frac{1}{2}(n^2 + n + 1)[(n^2 + n + 1)^2 - 3]\}^{\frac{1}{2}(n^2+n-1)[(n^2+n+1)^2+1]}. \end{aligned}$$

(5) The characteristic polynomial of  $Lap(\overline{L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]})$  is

$$\begin{aligned} \sigma(\overline{L(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]}, x) &= x[x - n(n+1)(n^2 + n + 1)]^{n(n+1)(n^2+n-1)} \\ &\cdot [x - (n^2 + n + 1)^2]^{n^2+n-1} [x - 2n(n+1)(n^2 + n + 1)]^{\frac{1}{2}(n^2+n-1)(n^2+n+2)[(n^2+n+1)^2+1]} \\ &\cdot [x - (n+1)^2(n^2 + 1)]^{n^2+n+1} [x - n^2(n^2 + 2n + 2)]^{n^2+n+1}. \end{aligned}$$

**Proof.** It is easy to check the correctness by Theorems 2, 6, 7, 11 and Lemmas 11, 12.  $\square$

**Theorem 12.** For integers  $n \geq 0$  and  $m \geq 0$ , if a regular graph  $G$  is integral, then the graphs  $L^m(G)$  and  $L^n(\overline{L^m(G)})$  are Laplacian integral.

**Proof.** It is easy to check the correctness by Lemmas 4, 11, 12, 13 and Theorem 9.  $\square$

**Corollary 8.** For integers  $p \geq 0$ ,  $n \geq 1$  and  $m \geq 0$ , the regular graphs  $L^m(K_{n,n+1} \equiv K_{n+1,n})$ ,  $L^p(\overline{L^m(K_{n,n+1} \equiv K_{n+1,n})})$ ,  $L^p(\overline{L^m(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}]})$  and  $L^m(K_{1,n^2+n+1}[n(n+1)K_{n^2+n+1}])$  are Laplacian integral.

**Proof.** It is easy to check the correctness by Lemmas 4, 11, 12, 13 and Corollary 7 as well as Theorems 1, 2 and 12.  $\square$

**Theorem 13.** If  $G$  is a regular graph of degree  $k$  with  $n$  vertices and  $m = \frac{1}{2}nk$  edges, let  $t \geq 2$  be an integer, then the characteristic polynomial of the  $(2^t k - 2^{t+1} + 2)$ -regular graph  $L^t(G)$  with  $n \prod_{i=0}^{t-1} (2^{i-1}k - 2^i + 1)$  vertices and  $n \prod_{i=0}^t (2^{i-1}k - 2^i + 1)$  edges is

$$\begin{aligned} P(L^t(G), x) &= P[G, x + (2 - k)(2^t - 1)](x + 2)^{2^{t-2}n(k-2) \prod_{i=0}^{t-2} (2^{i-1}k - 2^i + 1)} \\ &\cdot \prod_{j=2}^{t-1} [x + 2 + (2 - k) \sum_{i=j}^{t-1} 2^i]^{2^{j-2}n(k-2) \prod_{i=0}^{j-2} (2^{i-1}k - 2^i + 1)} [x + 2 + (2 - k) \\ &\cdot (2^t - 2)]^{\frac{1}{2}n(k-2)}. \end{aligned}$$

**Proof.** By induction on  $t \geq 2$ , we are easy to check the correctness from Lemmas 4 and 6.  $\square$

## References

- [1] F. Harary and A.J. Schwenk, Which graphs have integral spectra? Graphs and Combinatorics, Lectuer Notes in Mathematics 406, Springer-Verlag, Berlin (1974) 45-51.

- [2] F.C. Bussemaker and D. Cvetković, There are exactly 13 connected cubic integral graphs, *Publ. Elektrotech. Fak. Ser. Mat. Fiz.* 544(1976) 43-48.
- [3] A.J. Schwenk, Exactly thirteen connected cubic graphs have integral spectra, *Lecture Notes in Math. Ser.A 642*, Springer-Verlag, Berlin (1978) 516-533.
- [4] M. Roitman, An infinite family of integral graphs, *Discrete Math.* 52(2-3)(1984) 313-315.
- [5] Mikhail Muzychuk and Mikhail Klin, On graphs with three eigenvalues, *Discrete Math.* 189(1998) 191-207.
- [6] K.T. Balińska and S.K. Simić, The nonregular, bipartite, integral graphs with maximum degree 4– Part I: basic properties, *Discrete Math.* 236(2001) 13-24.
- [7] M. Watanabe and A.J. Schwenk, Integral starlike trees, *J. Austral. Math. Soc. Ser.A* 28(1979) 120-128.
- [8] M. Capobianco, S. Maurer, D. McCarthy and J. Molluzzo, A collection of open problems, *Annals New York Academy of Sciences* (1980) 582-583.
- [9] Xueliang Li and Guoning Lin, On the problem of integral trees, *Kexue Tongbao* 32:11(1987) 813-816.
- [10] Zhenfu Cao, On the integral trees of diameter  $R$  when  $3 \leq R \leq 6$ , *J. Heilongjiang University* 2(1988) 1-3.
- [11] Ruying Liu, Integral trees of diameter 5, *J. Systems Sci. & Math. Sci.* 8:4(1988) 357-360.
- [12] Zhenfu Cao, Some new classes of integral trees with diameter 5 or 6, *J. Systems Sci. & Math. Sci.* 11:1(1991) 20-26.
- [13] Pavol Híc, On balanced integral trees of diameter 6, *CO-MAT-TECH '97*, 5. vedecká konferencia smedzinárodnou účasťou, Trnava 14-15 (1997) 125-129.
- [14] Pavol Híc and Roman Nedela, Balanced integral trees, *Math. Slovaca* 48:5(1998) 429-445.
- [15] Pingzhi Yuan, Integral trees of diameter 4, *J. Systems Sci. & Math. Sci.* 18:2(1998) 177-181.
- [16] Ligong Wang and Xueliang Li, Some new classes of integral trees with diameters 4 and 6, *Australasian J. Combinatorics* 21(2000) 237-243.
- [17] Maosheng Li, Wensheng Yang and Jiabao Wang, Notes on the spectra of trees with small diameters, *J. Changsha Railway University* 18:2 (2000) 84-87.
- [18] Xueliang Li and Ligong Wang, Integral trees —A survey, *Chinese Journal of Engineering Math.* 17:5(2000) 91-93.

- [19] Ligong Wang, Xueliang Li and Xiangjuan Yao, Integral trees with diameters 4, 6 and 8, *Australasian J. Combinatorics* 25 (2002) 29-44.
- [20] Ligong Wang, Xueliang Li and Shenggui Zhang, Some new families of integral trees with diameters 4 and 6, *Electronic Notes in Discrete Mathematics* Vol.8 (2001)(or Ligong Wang, Xueliang Li and Shenggui Zhang, Families of integral trees with diameters 4, 6 and 8, *Discrete Appl. Math.* 136:(2-3)(2004) 349-362.
- [21] Ligong Wang, Xueliang Li and Shenggui Zhang, Construction of integral graphs, *Appl. Math. J. Chinese Univ. Ser.B* 15:3(2000) 239-246.
- [22] D. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs—Theory and Application*, Academic Press, New York, Francisco, London (1980) 59-60, 72-77, 266-267, 312-317.
- [23] D. Cvetković, M. Doob, I. Gutman and A. Torgašev, *Recent Results in the Theory of Graph Spectra*, North-Holland, Amsterdam (1988) 96-101.
- [24] Ligong Wang, Xueliang Li and C. Hoede, Integral complete  $r$ -partite Graphs, *Discrete Math.* 283:(1-3) (2004) 231-241.
- [25] Gangyuan Mao, *Circulant Matrices and Their Applications on Molecular Vibration*, Huazhang University of Technology Press 1995.
- [26] Zhihao Cao, A note on symmetric block circulant matrix, *J. Math. Res. Exposition* 10:3(1990) 469-473.
- [27] Norman Biggs, *Algebra Graph Theory*, Cambridge University Press 1993.
- [28] A.J. Schwenk and R.J. Wilson, On the eigenvalues of a graph, In: *Selected Topics in Graph Theory Algebra Graph Theory*, L.W. Beineke and R.J.Wilson Eds., Academic Press, London (1978) 307-336.
- [29] Bojan Mohar, The Laplacian spectrum of graphs, *Graph Theory, Combinatorics and Applications*, Y. Alavi et al. eds., John Wiley, New York (1991) 871-898.
- [30] Robert Grone and Russell Merris, The Laplacian spectrum of a graph II, *SIAM J. Discrete Math.* 7:2(1994) 221-229.
- [31] Russell Merris, Degree maximal graphs are Laplacian integral, *Linear Algebra Appl.* 199(1994) 381-389.
- [32] Philip J. Davis, *Circulant Matrices*, John Wiley & Sons, New York (1979).
- [33] F. Harary, *Graph Theory*, Addison-Wesley, Reading (1969).
- [34] Ligong Wang, Xueliang Li and C. Hoede, The spectrum of the regular graph  $K_p[K_{p-1}]$  of degree  $(p - 1)$  with  $p(p - 1)$  vertices, submitted.