

Trees with Small Randić Connectivity Indices

Haixing Zhao^{1,3} and Xueliang Li²

¹ Department of Computer Science and Engineering

Northwestern Polytechnical University

Xian, Shaanxi 710072, P.R. China

² Center for Combinatorics

Nankai University

Tianjin 300071, P.R. China

x.li@eyou.com

³ Department of Mathematics

Qinghai Normal University

Xining, Qinghai 810008, P.R. China

haixingzhao@yahoo.com.cn

Received August 1, 2003

Abstract

Let T be a tree and $d(v)$ the degree of its vertex v . Then the connectivity index, or Randić index, of T is defined as $\chi(T) = \sum_{uv} \frac{1}{\sqrt{d(u)d(v)}}$, where the summation goes over all edges uv of T . In the existing literature, trees of order n with m pending vertices and with the smallest connectivity index were determined by Hansen et al, whereas the unique tree of order n with the smallest connectivity index was determined by Bollobás et al. In this paper, we determine all trees of order n with m pending vertices and with the second smallest connectivity index and all trees of order n with diameter r and with the smallest and the second smallest connectivity indices. The unique tree of order n with, respectively, the second, the third and the fourth smallest connectivity index is also determined.

1 Introduction

The connectivity index, or Randić index, χ is a topological index proposed by Randić in 1975. Randić himself demonstrated that the Randić index χ was well correlated with a variety of physico-chemical properties of alkanes, such as boiling point, surface area and solubility in water. Eventually, χ became one of the most popular molecular graph-based structure-descriptors, used in QSPR and QSAR studies. On its applications for predicting physico-chemical and pharmacologic properties of organic compounds two books (Kier and Hall, 1976, 1986) and scores of papers were published; details and further bibliography can be found in [1,2] and [5-14].

Let G be a graph with the vertex set $V(G)$ and the edge set $E(G)$. Then the connectivity index is defined as

$$\chi(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_G(u)d_G(v)}},$$

where $d_G(x)$, or simply $d(x)$, denotes the degree of a vertex x in G .

For a graph G and $u \in V(G)$, we denote by $N_G(u)$ the set of all neighbors of u in G and by $n(G)$ the number of vertices of G . We denote respectively by S_n and P_n the star and the path with n vertices. By $P_{n,m}$ we denote the graph obtained from S_{n+1} and P_m by identifying the center of S_{n+1} with a vertex of degree 1 of P_m . By $S_{n,m}$ we denote the graph obtained from S_{n+2} and S_{m+1} by identifying a vertex of degree 1 of S_{n+2} with the center of S_{m+1} . We denote by $D(G)$ the diameter of G , which is defined as $D(G) = \max\{d(u,v) | u, v \in V(G)\}$ where $d(u,v)$ denotes the distance between the vertices u and v in G . We denote by $\mathcal{T}(n,r)$ the set of all trees with n vertices such that $D(T) = r$. Undefined notations and terminology will conform to those in [3].

Bollobás and Erdős [4] in 1998 got the fundamental result that among all connected graphs G with n vertices the star S_n has the smallest connectivity index χ , that is, $\chi(G) \geq \sqrt{n-1}$ for any such a graph G . Caporossi et al [5] in 1999 proved that $\chi(T) \leq \frac{n-3}{2} + \sqrt{2}$ for all trees T of order n and the equality holds if and only if $G \cong P_n$. Very recently, Hansen et al [10] obtained the following result.

Theorem 1.([10]) Let T be a tree of order $n \geq 3$ with m pending vertices. Then if $m < n - 1$,

$$\chi(T) \geq \sqrt{m} + \left(\frac{1}{\sqrt{2}} - 1 \right) \frac{1}{\sqrt{m}} + \frac{n-m-3}{2} + \frac{1}{\sqrt{2}},$$

and the equality holds if and only if T is the comet $T_{n,m}$, where $T_{n,m} \cong P_{m-1,n-m+1}$.

In this paper, we determine all trees of order n with m pending vertices and with the second smallest connectivity index and the unique tree of order n with, respectively, the second, the third and the fourth smallest connectivity index. Trees of order n with diameter r and with the smallest and the second smallest connectivity indices are also determined.

2 Lower Bounds for the Connectivity Index of Trees

In this section, we determine all trees with the second smallest connectivity index among all trees T of order n with m pending vertices. Let T be a tree with $n(T)$ vertices and u a vertex of T . Denote by $T + uv$ the tree obtained from T by adding a pendent edge uv . For convenience, we write $f(k) = \frac{1}{\sqrt{k+1}+\sqrt{k}}$ and $g(k) = \frac{1}{\sqrt{k(k+1)(\sqrt{k+1}+\sqrt{k})}}$.

Lemma 1. Let T a tree with a vertex u such that $d_T(u) = k$. Suppose that $N_T(u) = \{1, 2, 3, \dots, k\}$ and $v \in V(T)$. Then

$$\chi(T + uv) = \chi(T) + f(k) + g(k) \left(k - \sum_{i \in N_T(u)} \frac{1}{\sqrt{d_T(i)}} \right).$$

Proof. Suppose that $Q = \{ui | i \in N_T(u)\}$ and $\Omega = \sum_{xy \in E(T) - Q} \frac{1}{\sqrt{d_T(x)d_T(y)}}$. Then we have

$$\begin{aligned} \chi(T) &= \sum_{xy \in E(T)} \frac{1}{\sqrt{d_T(x)d_T(y)}} \\ &= \Omega + \sum_{i \in N_T(u)} \frac{1}{\sqrt{k d_T(i)}} \end{aligned}$$

and

$$\begin{aligned} \chi(T + uv) &= \sum_{xy \in E(T+uv)} \frac{1}{\sqrt{d_{T+uv}(x)d_{T+uv}(y)}} \\ &= \Omega + \sum_{i \in N_T(u)} \frac{1}{\sqrt{(k+1)d_T(i)}} + \frac{1}{\sqrt{k+1}}. \end{aligned}$$

So, we get that

$$\begin{aligned} \chi(T + uv) - \chi(T) &= \frac{1}{\sqrt{k+1}} - \frac{1}{\sqrt{k(k+1)(\sqrt{k+1}+\sqrt{k})}} \left(\sum_{i \in N_T(u)} \frac{1}{\sqrt{d_T(i)}} \right) \\ &= f(k) + g(k) \left(k - \sum_{i \in N_T(u)} \frac{1}{\sqrt{d_T(i)}} \right). \end{aligned}$$

Clearly, the lemma holds. \square

Let v be a vertex of T . One can see that there is a vertex $w \in N_T(v)$ such that $d_T(w) \geq 2$ except if v is the center of a star. So, we have

$$k - \sum_{i \in N_T(v)} \frac{1}{\sqrt{d_T(i)}} \geq 1 - \frac{1}{\sqrt{2}} \quad (1)$$

if $T + uv$ is not a star.

Denote by Q_{n_1, n_2} and P_{n_1, n_2, n_3} the two graphs shown in Figure 1, where G is a connected graph.



Figure 1. Graphs Q_{n_1, n_2} and P_{n_1, n_2, n_3}

Theorem 2. Let $n_1 \geq n_2 + 2 \geq 2$ and $d(v_2) \geq d(u_2)$. Then

$$\chi(Q_{n_1, n_2}) < \chi(Q_{n_1-1, n_2+1}).$$

Proof. By Lemma 1 it follows that

$$\chi(Q_{n_1, n_2}) = \chi(Q_{n_1-1, n_2}) + f(n_1) + g(n_1) \left(1 - \frac{1}{\sqrt{d(u_2)}}\right)$$

and

$$\chi(Q_{n_1-1, n_2+1}) = \chi(Q_{n_1-1, n_2}) + f(n_2 + 1) + g(n_2 + 1) \left(1 - \frac{1}{\sqrt{d(v_2)}}\right).$$

Since $n_1 \geq n_2 + 2$ and $d(v_2) \geq d(u_2)$, the theorem holds. \square

It is not hard to see that the proof of Theorem 2 is valid for the graph P_{n_1, n_2, n_3} , where $n_1 \geq n_3 + 2 \geq 2$ and $n_2 \geq 2$. So, we have the following lemma.

Lemma 2. Let $n_1 \geq n_3 + 2 \geq 2$ and $n_2 \geq 2$. Then

$$\chi(P_{n_1, n_2, n_3}) < \chi(P_{n_1-1, n_2, n_3+1}). \quad \square$$

Lemma 3. Let $n_1 \geq n_2 \geq 2$ and G be a tree. If Q_{n_1, n_2} has n vertices and m pending vertices, then $\chi(Q_{n_1, n_2}) \geq \chi(P_{m-2, n-m, 2})$.

Proof. By induction on m . Clearly, $m \geq 4$. When $m = 4$, $Q_{n_1, n_2} \cong P_{2, n-4, 2}$. So, the lemma is true for $m = 4$ and all $n \geq m + 2$.

Suppose that $m \geq 5$ and the lemma holds for every Q_{s_1, s_2} of order n with $m - 1$ pending vertices, where $s_1 \geq s_2 \geq 2$ and $s_1 + s_2 = n_1 + n_2$. We distinguish the following cases:

Case 1. $n_1 = 2$ or $n_2 = 2$.

Let $n_2 = 2$ and $T' = Q_{n_1,1}$. By Lemma 1, we have

$$\chi(Q_{n_1,n_2}) = \chi(T') + f(2) + g(2) \left(1 - \frac{1}{\sqrt{d_{T'}(v_2)}}\right)$$

and

$$\chi(P_{m-2,n-m,2}) = \chi(P_{m-2,n-m+1}) + f(2) + g(2) \left(1 - \frac{1}{\sqrt{2}}\right).$$

Note that T' has $n - 1$ vertices and $m - 1$ pending vertices. From Theorem 1 we have that $\chi(T') \geq \chi(P_{m-2,n-m+1})$ and the equality holds if and only if $T' \cong P_{m-2,n-m+1}$. So, we have that $\chi(Q_{n_1,n_2}) \geq \chi(P_{m-2,n-m,2})$ and the equality holds if and only if $Q_{n_1,n_2} \cong P_{m-2,n-m,2}$.

Case 2. $n_1 \geq 3$ and $n_2 \geq 3$.

Let $T' = Q_{n_1-1,n_2}$. By Lemma 1, we have

$$\chi(Q_{n_1,n_2}) = \chi(T') + f(n_1) + g(n_1) \left(1 - \frac{1}{\sqrt{d_{T'}(u_2)}}\right) \quad (2)$$

and

$$\chi(P_{m-2,n-m,2}) = \chi(P_{m-3,n-m,2}) + f(m-2) + g(m-2) \left(1 - \frac{1}{\sqrt{2}}\right). \quad (3)$$

Since $T' \cong Q_{n_1-1,n_2}$ and T' has $m-1$ pending vertices, by the induction hypothesis, $\chi(P_{m-3,n-m,2}) \leq \chi(T')$. Note that $n_1 \leq m-3$ and Q_{n_1,n_2} is not a star. Thus from (1), (2) and (3) we have

$$\chi(Q_{n_1,n_2}) > \chi(P_{m-2,n-m,2}).$$

This completes the proof. \square

Let $v_1v_2v_3 \cdots v_k$ be a path P_k and $T_{k,v_i,m}$ be a graph shown in Figure 2, where $k \geq 5$ and $m \geq 1$.




Figure 2. Graph $T_{k,v_i,m}$

Lemma 4. Let $r \geq 4$ and $n \geq r + 3$. Then

$$\chi(P_{n-r-1,r-1,2}) > \chi(T_{r+1,v_3,n-r-1}).$$

Proof. By the definition of χ , we have

$$\chi(P_{n-r-1,r-1,2}) = \frac{n-r-1}{\sqrt{n-r}} + \frac{1}{\sqrt{2(n-r)}} + \frac{r-4}{2} + \frac{2}{\sqrt{3}} + \frac{1}{\sqrt{6}}$$

and

$$\chi(T_{r+1,v_3,n-r-1}) = \frac{n-r-1}{\sqrt{n-r+1}} + \frac{\sqrt{2}}{\sqrt{n-r+1}} + \frac{r-4}{2} + \sqrt{2}.$$

Let $x = n - r$. Obviously, x is an integer and $x \geq 3$. So, we get that

$$\chi(P_{n-r-1,r-1,2}) - \chi(T_{r+1,v_3,n-r-1}) = \phi(x), \quad (4)$$

where $\phi(x) = (\sqrt{2} - 1) \left(\frac{\sqrt{2}}{\sqrt{x+1}} - \frac{1}{\sqrt{2x}} \right) + \frac{2\sqrt{2+1-\sqrt{12}}}{\sqrt{6}} - \frac{1}{\sqrt{x+1+\sqrt{x}}}$. By calculation, one can check that $\phi(x) > 0$ for each $x \in \{3, 4, 5, 6, 7, 8, 9, 10\}$ and $\phi(x) > \frac{2\sqrt{2+1-\sqrt{12}}}{\sqrt{6}} - \frac{1}{\sqrt{11+\sqrt{12}}} > 0$ for $x \geq 11$.

Thus by formula (4) the lemma is true.

Lemma 5. If $T \in \mathcal{T}(n, 4) - \{P_{n-4,4}\}$, then $\chi(T) \geq \frac{n-5}{\sqrt{n-3}} + \frac{\sqrt{2}}{\sqrt{n-3}} + \sqrt{2}$ and the equality holds if and only if $T \cong T_{5,v_3,n-5}$.

Proof. Clearly, $T \in \mathcal{T}(n, 4) - \{P_{n-5,5}\}$ if and only if T has $n - 3$ pending vertices. So, we have the following cases:

Case 1. There is a path $u_1u_2u_3u_4u_5$ in T such that $d(u_2) \geq 3$ and $d(u_4) \geq 3$. By Lemmas 3 and 4,

$$\chi(T) \geq \chi(P_{n-5,3,2}) > \chi(T_{5,v_3,n-5}).$$

Case 2. For each path $u_1u_2u_3u_4u_5$ in T , we must have $d(u_2) = 2$ or $d(u_4) = 2$. Recalling that $D(T) = 4$, one can see that T must be the graph $U_k(n_1, t)$ shown in Figure 2, where $k \geq 0$, $n_1 \geq 1$, $t \geq 2$ and $n_1 + 2t + k = n$.




Figure 3. Graph $U_k(n_1, t)$

By Lemma 1,

$$\chi(T_{5,v_3,n-5}) = \chi(T_{5,v_3,n-6}) + f(n-4) + g(n-4)(2 - \sqrt{2}). \quad (5)$$

Subcase 2.1. $k \geq 1$ in $U_k(n_1, t)$.

By Lemma 1, we have

$$\chi(U_k(n_1, t)) \geq \chi(U_{k-1}(n_1, t)) + f(k+t-1) + g(k+t-1)(2 - \sqrt{2}) \quad (6)$$

and the equality holds if and only if $n_1 = 1$ and $t = 2$, that is, $U_k(n_1, t) \cong T_{5, v_3, n-5}$. Since $k+t-1 \leq n-4$, by (5) and (6) we have

$$\chi(U_k(n_1, t)) \geq \chi(T_{5, v_3, n-5})$$

and the equality holds if and only if $U_k(n_1, t) \cong T_{5, v_3, n-5}$.

Subcase 2.2. $k = 0$ and $t \geq 3$ in $U_k(n_1, t)$.

By Lemma 1, we have that

$$\chi(U_0(n_1, t)) = \chi(U_1(n_1, t-1)) + f(1) + g(1)\left(1 - \frac{1}{\sqrt{t}}\right)$$

and

$$\chi(U_1(n_1+1, t-1)) = \chi(U_1(n_1, t-1)) + f(n_1+1) + g(n_1+1)\left(1 - \frac{1}{\sqrt{t}}\right).$$

Clearly, $n_1+1 > 1$. So, $\chi(U_0(n_1, t)) > \chi(U_1(n_1+1, t-1))$. From Subcase 2.1, we have

$$\chi(U_0(n_1, t)) > \chi(U_1(n_1+1, t-1)) \geq \chi(T_{5, v_3, n-5}).$$

Subcase 2.3. $k = 0$ and $t = 2$ in $U_k(n_1, t)$.

So, $U_0(n_1, 2) \cong P_{5, n-5}$, which contradicts to the condition of the lemma.

By calculation, we have

$$\chi(T_{5, v_3, n-5}) = \frac{n-5}{\sqrt{n-3}} + \frac{\sqrt{2}}{\sqrt{n-3}} + \sqrt{2}.$$

This completes the proof. \square

Let T be a tree of order $n \geq 3$ with m pending vertices. Obviously, $m \leq n-1$ and the equality holds if and only if $T = S_n$. When $m = n-2$, one can see that $T \in \{S_{n_1, n_2} | n_1 + n_2 = n-2, n_1 \geq n_2\}$. By Lemma 2, for all $t \geq 5$ we have

$$\chi(S_{n-3, 1}) > \chi(S_{n-4, 2}) > \chi(S_{n-t, t-2}).$$

So, $S_{n-4, 2}$ has the second smallest connectivity index among all trees of order n with $n-2$ pending vertices. For $m < n-2$, we have

Theorem 3. Let T be a tree of order $n \geq 3$ with m pending vertices. If $m < n - 2$ and $T \not\cong P_{m-1, n-m+1}$, then

$$\chi(T) \geq \sqrt{m} + (\sqrt{2} - 2) \frac{1}{\sqrt{m}} + \frac{n - m - 3}{2} + \sqrt{2}$$

and the equality holds if and only if $T \in \{T_{n-m+2, v_i, m-2} | 3 \leq i \leq n - m\}$.

Proof. Let T be a tree of order $n \geq 3$ with m pending vertices. By the condition of the theorem, $m \geq 3$ and $n \geq m + 3$. By calculation, it is not difficult to obtain that for $3 \leq i \leq r - 1$,

$$\chi(T_{n-m+2, v_3, m-2}) = \chi(T_{n-m+2, v_i, m-2}) = \sqrt{m} + (\sqrt{2} - 2) \frac{1}{\sqrt{m}} + \frac{n - m - 3}{2} + \sqrt{2}.$$

So, we need only to prove that $\chi(T) \geq \chi(T_{n-m+2, v_3, m-2})$. We prove it by induction on n .

For any $m \geq 3$, one can see that $n = m + 3$ if and only if $D(T) = 4$. By Lemma 5, the theorem holds for $n = m + 3$ with $m \geq 3$.

Suppose that $n \geq m + 4$ and that the theorem is true for all trees T of order $n - 1$ with m pending vertices. Then for a tree T of order n with m pending vertices, we consider the following cases:

Case 1. T is a tree of form Q_{n_1, n_2} such that $n_1 \geq n_2 \geq 2$. Then, from Lemmas 3 and 4, it follows that

$$\chi(T) \geq \chi(P_{m-2, n-m, 2}) > \chi(T_{n-m+2, v_3, m-2}).$$

Case 2. There is a path $u_1 u_2 u_3$ in T such that $d(u_1) = 1$, $d(u_2) = 2$ and $d(u_3) \geq 2$. Let $T' = T - u_1$. Then, by Lemma 1 we have that

$$\chi(T) \geq \chi(T') + f(1) + g(1) \left(1 - \frac{1}{\sqrt{d_{T'}(u_3)}}\right) \quad (7)$$

and

$$\chi(T_{n-m+2, v_3, m-2}) = \chi(T_{n-m+1, v_3, m-2}) + f(1) + g(1) \left(1 - \frac{1}{\sqrt{2}}\right). \quad (8)$$

Clearly, T' has $n - 1$ vertices and m pending vertices. Since $T \not\cong P_{m-1, n-m+1}$, we have $T \cong T_{n-m+2, v_3, m-2}$ if $T' \cong P_{m-1, n-m}$. For $T' \not\cong P_{m-1, n-m}$, by the induction hypothesis we have $\chi(T') \geq \chi(T_{n-m+1, v_3, m-2})$. So, from (7) and (8),

$$\chi(T) \geq \chi(T_{n-m+2, v_3, m-2})$$

and the equality holds if and only if $T' \in \{T_{n-m+1, v_i, m-2} | 3 \leq i \leq n - m - 1\}$ and $d_{T'}(u_3) = 2$. Furthermore, the equality holds if and only if $d_T(u_3) = 2$ and $T \in \{T_{n-m+2, v_i, m-2} | 3 \leq i \leq n - m\}$.

This completes the proof. \square

In the following, using Theorems 1 and 3 we find the smallest value of the connectivity index of trees in $\mathcal{T}(n, r)$ and determine the corresponding trees. Let $T \in \mathcal{T}(n, r)$ and $r \geq 3$. Then, there is a path $u_1 u_2 \cdots u_{r+1}$ in T such that $d(u_1) = d(u_{r+1}) = 1$ and $d(u_i) \geq 2$ for all $2 \leq i \leq r$. So, T has at most $n - r + 1$ pending vertices. By Theorem 1, it is not difficult to see that $\chi(T) \geq \chi(P_{m-1, n-m+1})$ if T has m pending vertices. By Lemma 2, for $m \geq 3$ we have

$$\chi(P_{m-2, n-m+1, 1}) > \chi(P_{m-1, n-m+1, 0}),$$

that is,

$$\chi(P_{m-2, n-m+2}) > \chi(P_{m-1, n-m+1}). \quad (9)$$

Thus, we have $\chi(T) \geq \chi(P_{n-r, r})$ and the equality holds if and only if $T \cong P_{n-r, r}$.

For $r = 3$, we have that $S_{n-4, 2}$ is the unique tree with the second smallest connectivity index among all trees of order n with diameter 3. For $r \geq 4$, by (9) and Theorems 1 and 3 we have that if $T \not\cong P_{n-r, r}$, then $\chi(T) \geq \chi(T_{r+1, v_3, n-r-1})$ when T has $n - r + 1$ pending vertices and $\chi(T) \geq \chi(P_{n-r-1, r+1})$ when T has no more than $n - r$ pending vertices. By calculation, we have

$$\chi(T_{r+1, v_3, n-r-1}) = \frac{n-r-1}{\sqrt{n-r+1}} + \frac{\sqrt{2}}{\sqrt{n-r+1}} + \frac{r-4}{2} + \sqrt{2}. \quad (10)$$

and

$$\chi(P_{n-r-1, r+1}) = \frac{n-r-1}{\sqrt{n-r}} + \frac{1}{\sqrt{2(n-r)}} + \frac{r-2}{2} + \frac{1}{\sqrt{2}}. \quad (11)$$

From (10) and (11), we have

$$\chi(P_{n-r-1, r+1}) - \chi(T_{r+1, v_3, n-r-1}) = \psi(x),$$

where $\psi(x) = \frac{x-1}{\sqrt{x}} - \frac{x-1}{\sqrt{x+1}} + \frac{1}{\sqrt{2x}} - \frac{\sqrt{2}}{\sqrt{x+1}} + 1 - \frac{1}{\sqrt{2}}$ and $x = n - r \geq 2$. It is easy to check that $\psi(x) > 0$ for $x = 2, 3, 4$. For $x \geq 5$, we have

$$\psi(x) > 1 - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2(x+1)}} - \frac{\sqrt{2}}{\sqrt{x+1}} \geq 1 - \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{12}} > 0.$$

So, from the above arguments and Theorems 1 and 3, we have

Theorem 4. (i) For $T \in \mathcal{T}(n, r)$ and $r \geq 3$, we have

$$\chi(T) \geq \frac{n-r}{\sqrt{n-r+1}} + \frac{1}{\sqrt{2(n-r+1)}} + \frac{r-3}{2} + \frac{1}{\sqrt{2}},$$

and the equality holds if and only if $T \cong P_{n-r,r}$.

(ii) For $r \geq 4$ and $T \in \mathcal{T}(n, r) - \{P_{n-r,r}\}$, we have

$$\chi(T) \geq \frac{n-r-1}{\sqrt{n-r+1}} + \frac{\sqrt{2}}{\sqrt{n-r+1}} + \frac{r-4}{2} + \sqrt{2},$$

and the equality holds if and only if $T \in \{T_{r+1,v_i,n-r-1} | 3 \leq i \leq r-1\}$. \square

3 Trees with Small Connectivity Indices

In this section, we determine the unique tree of order n with, respectively, the second, the third and the fourth smallest connectivity index.

Let T be a tree of order n . Then, for smaller n we have the following: for $n = 1, 2, 3$, we have $T \cong S_n$; for $n \geq 4$, we can easily check that

- (a) for $n = 4$, $\chi(P_3) > \chi(S_3)$;
- (b) for $n = 5$, $\chi(P_5) > \chi(S_{2,1}) > \chi(S_4)$;
- (c) for $n = 6, 7$, $\chi(T) > \chi(P_{n-4,4}) > \chi(S_{n-4,2}) > \chi(S_{n-3,1}) > \chi(S_n)$ if $T \notin \{P_{n-4,4}, S_{n-4,2}, S_{n-3,1}, S_n\}$.

Now we consider the case $n \geq 8$. By Lemma 2, we have

$$\chi(S_{n_1,n_2}) > \chi(S_{n_1+1,n_2-1}) \text{ for } n_1 \geq n_2 \geq 2. \quad (12)$$

By (9) and Theorem 4, we have

$$\chi(T) > \chi(P_{n-4,4}) \text{ if } T \in \mathcal{T}(n, r) - P_{n-4,4} \text{ and } r \geq 4. \quad (13)$$

By calculation, we obtain the following:

- (i) $\chi(S_{n-3,1}) = \frac{n-3}{\sqrt{n-2}} + \frac{1}{\sqrt{2(n-2)}} + \frac{1}{\sqrt{2}}$,
- (ii) $\chi(S_{n-4,2}) = \frac{n-4}{\sqrt{n-3}} + \frac{1}{\sqrt{3(n-3)}} + \frac{2}{\sqrt{3}}$,
- (iii) $\chi(S_{n-5,3}) = \frac{n-5}{\sqrt{n-4}} + \frac{1}{2\sqrt{n-4}} + \frac{3}{2}$,
- (iv) $\chi(P_{n-4,4}) = \frac{n-4}{\sqrt{n-3}} + \frac{1}{\sqrt{2(n-3)}} + \frac{1+\sqrt{2}}{2}$.

From (i) to (iv), it is not difficult to obtain that $\chi(S_{n-5,3}) > \chi(S_{n-4,2}) > \chi(S_{n-3,1})$ and $\chi(P_{n-4,4}) > \chi(S_{n-4,2}) > \chi(S_{n-3,1})$ for $n \geq 8$. On the other hand, we have

$$\chi(S_{n-5,3}) - \chi(P_{n-4,4}) = z(x), \quad (14)$$

where $z(x) = 1 - \frac{\sqrt{2}}{2} - \frac{1}{2\sqrt{x}} - \frac{1}{\sqrt{2(x+1)}} + \frac{x}{\sqrt{x}} - \frac{x}{\sqrt{x+1}}$ and $x = n - 4 \geq 4$. By calculation, one obtains that $z(x) < 0$ for $n = 8, 9$ and $z(x) > 0$ for $10 \leq n \leq 20$. For $n \geq 21$, it follows that

$$z(x) > 1 - \frac{\sqrt{2}}{2} - \frac{1}{2\sqrt{17}} - \frac{1}{\sqrt{36}} > 0.$$

So, by (14) we have that $\chi(S_{n-5,3}) < \chi(P_{n-4,4})$ for $n = 8, 9$ and $\chi(S_{n-5,3}) > \chi(P_{n-4,4})$ for $n \geq 10$. Thus, by (c), (12) and (13), we have the following theorem.

Theorem 5. Let T be a tree of order n and let $T \notin \{S_{n-4,2}, S_{n-3,1}, S_n\}$. Then,

(i) $\chi(T) \geq \chi(S_{n-5,3}) > \chi(S_{n-4,2}) > \chi(S_{n-3,1}) > \chi(S_n)$ for $n = 8, 9$ and the equality holds if and only if $T \cong S_{n-5,3}$.

(ii) $\chi(T) \geq \chi(P_{n-4,4}) > \chi(S_{n-4,2}) > \chi(S_{n-3,1}) > \chi(S_n)$ for $n = 6, 7$ or $n \geq 10$ and the equality holds if and only if $T \cong P_{n-4,4}$. \square

Acknowledgements

We would like to express our thanks to the referees for the valuable suggestions which helped to improve the paper. The work is supported by National Science Foundation of China and the Science Foundation of the State Education Ministry of China.

References

- [1] O. Araujo and J. Rada, Randić index and lexicographic order, *J. Math. Chem.* 27 (2000) 19-30.
- [2] O. Araujo and J.A. de la Pena, The connectivity index of a weighted graph, *Lin. Alg. Appl.* 283 (1998) 171-177.
- [3] N. Biggs, *Algebraic Graph Theory*, Cambridge University Press, London, 1993.
- [4] B. Bollobás and P. Erdős, Graphs of extremal weights, *Ars Combin.* 50(1998) 223-225.
- [5] G. Caporossi, I. Gutman and P. Hansen, Variable neighborhood search for extremal graphs IV: Chemical trees with extremal connectivity index, *Computers and Chemistry* 23(1999) 469-477.
- [6] L.H. Clark and J.W. Moon, On the general Randić index for certain families of trees, *Ars Combin.* 54(2000) 223-235.

- [7] I. Gutman and M. Randić, Algebraic characterization of skeletal branching, *Chem. Phys. Lett.* 47 (1977) 15-19.
- [8] I. Gutman, O. Miljković, G. Caporossi and P. Hansen, Alkanes with small and large Randić connectivity indices, *Chem. Phys. Lett.* 306(1999) 366-372.
- [9] I. Gutman, O. Araujo and D.A. Morales, Bounds for the Randić connectivity index, *J. Chem. Inf. Comput. Sci.* 43(2000) 593-598.
- [10] P. Hansen and H. Mélot, Variable neighborhood search for extremal graphs 6. Analyzing bounds for the connectivity index, *J. Chem. Inf. Comput. Sci.* 43(2003) 1-14.
- [11] L.B. Kier and L.H. Hall, *Molecular Connectivity in Chemistry and Drug Research*, Academic Press, New York, 1976.
- [12] L.B. Kier and L.H. Hall, *Molecular Connectivity in Structure-Activity Analysis*, Wiley, New York, 1986.
- [13] M. Randić, On the characterization of molecular branching, *J. Am. Chem. Soc.* 97(1975) 6609-6615.
- [14] M. Randić, Representation of molecular graphs by basic graphs, *J. Chem. Inf. Comput. Sci.* 32(1992) 57-69.