

ON MOLECULAR GRAPHS WITH SMALLEST AND GREATEST ZEROth-ORDER GENERAL RANDIĆ INDEX*

Yumei Hu, Xueliang Li, Yongtang Shi, Tianyi Xu

*Center for Combinatorics and LPMC
Nankai University, Tianjin 300071, P.R. China*

Ivan Gutman

*Faculty of Science, University of Kragujevac, Kragujevac
Serbia and Montenegro*

(Received April 8, 2005)

Abstract

A molecular (n, m) -graph G is a connected simple graph with n vertices, m edges and vertex degrees not exceeding 4. If $d(v)$ denotes the degree of the vertex v , then the zeroth-order general Randić index ${}^0R_\alpha$ of the graph G is defined as $\sum_{v \in V(G)} d(v)^\alpha$, where α is a pertinently chosen real number. We characterize, for any α , the molecular (n, m) -graphs with smallest and greatest ${}^0R_\alpha$.

*Supported by the National Science Foundation of China.

1 Introduction

The Randić (or connectivity) index was introduced by Randić in 1975 and is defined as [25]

$$R = R(G) = \sum_{uv \in E(G)} (d(u)d(v))^{-1/2}$$

where $d(u)$ denotes the degree of the vertex u of the graph G , and $E(G)$ is the edge set of G . Randić himself demonstrated [25] that this index is well correlated with a variety of physico-chemical properties of various classes of organic compounds. Eventually, this structure-descriptor became one of the most popular topological indices to which two books [15, 17], several reviews [7, 24, 26] and countless research papers are devoted.

Like other successful structure-descriptors, the Randić index received considerable attention also from mathematical chemists and mathematicians. In particular, bounds for R and graphs extremal with regard to R were extensively studied [1, 2, 4, 8, 9, 10, 11, 12].

In 1998 Bollobás and Erdős [3] generalized $R(G)$ by replacing the exponent $-1/2$ by an arbitrary real number α . This graph invariant is called *the general Randić index* and will be denoted by $R_\alpha = R_\alpha(G)$. Li and Yang [20] studied R_α for general n -vertex graphs, obtained lower and upper bounds for it, and characterized the corresponding extremal graphs. Later Hu, Li and Yuan [13, 14] determined the trees extremal with regard to R_α , whereas Li, Wang and Wei [19] gave lower and upper bounds for R_α of molecular (n, m) -graphs. Other mathematical studies of the general Randić index are found in [5, 6].

The zeroth-order Randić index, conceived by Kier and Hall [16], is

$${}^0R = {}^0R(G) = \sum_{v \in V(G)} d(v)^{-1/2}$$

Eventually, Li and Zheng [22] defined the *zeroth-order general Randić index* of a graph G as

$${}^0R_\alpha = {}^0R_\alpha(G) = \sum_{v \in V(G)} d(v)^\alpha$$

for any real number α .

Pavlović determined the graphs with maximum 0R -index [23]. Li et al. [18] investigated the same problem for the topological index $M_1(G)$, one of the Zagreb indices, that is defined as $M_1(G) = \sum_{v \in V(G)} d(v)^2$. (Evidently, $M_1 \equiv {}^0R_\alpha$ for $\alpha = +2$.) They obtained sufficient and necessary conditions under which (n, m) -graphs have minimum M_1 , and a necessary condition for an (n, m) -graph having maximum $M_1(G)$. Li and Zhao [21] characterized trees with the first three smallest and largest zeroth-order general Randić index, with the exponent α being equal to k , $-k$, $1/k$, and $-1/k$, where $k \geq 2$ is an integer.

In this paper we investigate the zeroth-order general Randić index of molecular (n, m) -graphs, i. e., connected simple graphs with n vertices, m edges and maximum vertex degree at most 4. We characterize the molecular (n, m) -graphs with extremal (maximum or minimum) zeroth-order general Randić index.

First we need to introduce some notation.

Denote by $D(G) = [d_1, d_2, \dots, d_n]$ the degree sequence of the graph G , where d_i stands the degree of the i -th vertex of G , and $d_1 \geq d_2 \geq \dots \geq d_n$.

If there is a graph G , such that $d_i \geq d_j + 2$, let G' be the graph obtained from G by replacing the pair (d_i, d_j) by the pair $(d_i - 1, d_j + 1)$. In other words, if $D(G) = [d_1, d_2, \dots, d_{i-1}, d_i, d_{i+1}, \dots, d_{j-1}, d_j, d_{j+1}, \dots, d_n]$, then $D(G') = [d_1, d_2, \dots, d_{i-1}, d_i - 1, d_{i+1}, \dots, d_{j-1}, d_j + 1, d_{j+1}, \dots, d_n]$.

Note that if $\alpha = 0$ then ${}^0R_\alpha(G) = n$, and if $\alpha = 1$ then ${}^0R_\alpha(G) = 2m$. Therefore, in the following we always assume that $\alpha \neq 0, 1$.

Lemma 1.1 *For the two graphs G and G' , specified above, we have*

$$(i) \quad {}^0R_\alpha(G) > {}^0R_\alpha(G') \quad \text{for } \alpha < 0 \text{ or } \alpha > 1$$

$$(ii) \quad {}^0R_\alpha(G) < {}^0R_\alpha(G') \quad \text{for } 0 < \alpha < 1.$$

Proof. Since ${}^0R_\alpha(G) = \sum_{v \in V(G)} d(v)^\alpha$, we have

$$\begin{aligned} {}^0R_\alpha(G) - {}^0R_\alpha(G') &= d_i^\alpha + d_j^\alpha - (d_i - 1)^\alpha - (d_j + 1)^\alpha \\ &= [d_i^\alpha - (d_i - 1)^\alpha] - [(d_j + 1)^\alpha - d_j^\alpha] \\ &= \alpha (\xi_1^{\alpha-1} - \xi_2^{\alpha-1}) \end{aligned}$$

where $\xi_1 \in (d_i - 1, d_i)$, and $\xi_2 \in (d_j, d_j + 1)$. So, by $d_i \geq d_j + 2$, we have $\xi_1 > \xi_2$. Then ${}^0R_\alpha(G) > {}^0R_\alpha(G')$ for $\alpha < 0$ or $\alpha > 1$, whereas ${}^0R_\alpha(G) < {}^0R_\alpha(G')$ for $0 < \alpha < 1$.

■

2 Extremal molecular (n, m) -graphs

Denote by n_i the number of vertices of degree i in a molecular (n, m) -graph G . Then we have

$${}^0R_\alpha(G) = n_1 + 2^\alpha n_2 + 3^\alpha n_3 + 4^\alpha n_4 \quad (2.1)$$

Theorem 2.1 *Let C^* be a molecular (n, m) -graph with degree sequence $[d_1, d_2, \dots, d_n]$, such that $|d_i - d_j| \leq 1$ for any $i \neq j$. Then for $\alpha < 0$ or $\alpha > 1$, C^* has the minimum zeroth-order general Randić index among all molecular (n, m) -graphs, whereas for $0 < \alpha < 1$, C^* has the maximum zeroth-order general Randić index among all molecular (n, m) -graphs. Moreover,*

$${}^0R_\alpha(C^*) = \begin{cases} 2 + 2^\alpha (n - 2) & \text{if } m = n - 1 \\ 2^\alpha (3n - 2m) + 3^\alpha (2m - 2n) & \text{if } n \leq m \leq \lfloor 3n/2 \rfloor \\ 3^\alpha (4n - 2m) + 4^\alpha (2m - 3n) & \text{if } \lfloor 3n/2 \rfloor < m \leq 2n \end{cases}$$

Proof. We only consider the case $0 < \alpha < 1$, because the proof for the other case is fully analogous. Let G be a molecular graph and $D(G) = [d_1, d_2, \dots, d_n]$. If $G \not\cong C^*$, then there must exist a pair (d_i, d_j) such that $d_i \geq d_j + 2$. By Lemma 1.1, the graph G' , obtained by replacing the pair (d_i, d_j) by the pair $(d_i - 1, d_j + 1)$, has a greater ${}^0R_\alpha$ -value than G . Consequently, G is not a molecular (n, m) -graph with maximum zeroth-order general Randić index.

To show the existence, we construct the extremal (n, m) -graph C^* (with minimum ${}^0R_\alpha$ for $\alpha < 0$ or $\alpha > 1$, and with maximum ${}^0R_\alpha$ for $0 < \alpha < 1$) by adding edges one by one. First, we start from a tree. There must be at least two 1-degree vertices in a tree. By Lemma 1.1, there does not exist any 3-degree vertex, and so the extremal tree must be the path P_n . Next we add an edge joining the two leaves of the path. In this way the degrees of all vertices become equal to two, and then we get a cycle. We continue by adding edges one by one, so as to maximize the number of 3-degree vertices, until either there remain no 2-degree vertices, or remains exactly one. If more edges need to be added, then we first connect the 2-degree vertex (if such does exist) with a non-adjacent 3-degree vertex, and continue by connecting pairs of nonadjacent 3-degree vertices. The construction is shown in Figure 2.1. ■




Figure 2.1 Constructing molecular graphs with extremal zeroth-order general Randić index, according to Theorem 2.1.

Theorem 2.2 *Let G^* be a molecular (n, m) -graph with at most one vertex of degree 2 or 3. If one of the following conditions holds:*

(I) $m = n - 1$

(II) $m \geq n \geq 6$, for $n = 6$, $m \geq 10$, and for $n = 7$, $m \neq 8$

then for $\alpha < 0$ or $\alpha > 1$, G^* has the maximum zeroth-order general Randić index among all molecular (n, m) -graphs, whereas for $0 < \alpha < 1$, the same graph has the minimum zeroth-order general Randić index among all molecular (n, m) -graphs. Moreover,

$${}^0R_\alpha(G^*) = \begin{cases} (4n - 2m)/3 + 4^\alpha (2m - n)/3 & \text{if } 2m - n \equiv 0 \pmod{3} \\ (4n - 2m - 2)/3 + 2^\alpha + 4^\alpha (2m - n - 1)/3 & \text{if } 2m - n \equiv 1 \pmod{3} \\ (4n - 2m - 1)/3 + 3^\alpha + 4^\alpha (2m - n - 2)/3 & \text{if } 2m - n \equiv 2 \pmod{3} \end{cases}$$

Proof. Again, we only consider the case $0 < \alpha < 1$, because the proof for the other case is similar. Let G' be a molecular (n, m) -graph and $D(G') = [d_1, d_2, \dots, d_n]$. Let G' possess two vertices of degree 2 or 3, i. e., let there be a pair (d_i, d_j) , such that $3 \geq d_i \geq d_j \geq 2$. Then by Lemma 1.1, there is a graph G , obtained by replacing the pair (d_i, d_j) by the pair $(d_i + 1, d_j - 1)$, that has a smaller ${}^0R_\alpha$ -value than G' . Repeating the above operation until there is no pair (d_i, d_j) , such that $3 \geq d_i \geq d_j \geq 2$, we arrive at G^* with minimum zeroth-order general Randić index. In view of (2.1), for G^* we have

$$\begin{cases} n_1 + n_2 + n_3 + n_4 & = n \\ n_1 + 2n_2 + 3n_3 + 4n_4 & = 2m \\ n_2 + n_3 & \leq 1 \end{cases}$$

From the above equations, we have one of the following three options:

- (1) $n_2 = n_3 = 0$, implying $n_1 = (4n - 2m)/3$, $n_4 = (2m - n)/3$, and $2m - n \equiv 0 \pmod{3}$
- (2) $n_2 = 1$, $n_3 = 0$, implying $n_1 = (4n - 2m - 2)/3$, $n_4 = (2m - n - 1)/3$, and $2m - n \equiv 1 \pmod{3}$
- (3) $n_2 = 0$, $n_3 = 1$, implying $n_1 = (4n - 2m - 1)/3$, $n_4 = (2m - n - 2)/3$, and $2m - n \equiv 2 \pmod{3}$.

In order to show the existence, we construct G^* by distinguishing the following cases:

- (I) $m = n - 1$, i. e., G^* is a tree.
 - (I.1) If $2m - n \equiv 0 \pmod{3}$, we first construct a path with n_4 vertices, and then add n_1 pendent vertices, taking care that no vertex gets degree greater than 4.
 - (I.2) If $2m - n \equiv 1 \pmod{3}$, we first construct a path with n_4 vertices, then add n_1 pendent vertices, taking care that no vertex gets degree greater than 4, and finally subdivide an edge by inserting to it a vertex of degree 2.
 - (I.3) If $2m - n \equiv 2 \pmod{3}$, we first construct a path with $n_4 + 1$ vertices, and then add n_1 pendent vertices, taking care that no vertex gets degree greater than 4.

(II) $m \geq n \geq 6$, for $n = 6$ and $m \geq 10$, or for $n = 7$ and $m \neq 8$.

In Figure 2.2 we show one of the possible graphs G^* for $n_4 \leq 4$, that is for

$\lfloor (2m - n)/3 \rfloor \leq 4$. For $n_4 \geq 5$, we construct G^* as follows:

- (II.1) If $2m - n \equiv 0 \pmod{3}$, we first construct a 4-regular graph on n_4 vertices, then delete $n_1/2$ edges from it, and then add n_1 pendent vertices, taking care that no vertex gets degree greater than 4.
- (II.2) If $2m - n \equiv 1 \pmod{3}$, we first construct a 4-regular graph on n_4 vertices, then delete $n_1/2$ edges from it, then add n_1 pendent vertices, taking care that no vertex gets degree greater than 4, and finally subdivide an edge inserting to it a vertex of degree 2.
- (II.3) If $2m - n \equiv 2 \pmod{3}$, we first construct a 4-regular graph on $n_4 + 1$ vertices, then delete $(n_1 + 1)/2$ edges from it, and then add n_1 pendent vertices, taking care that no vertex gets degree greater than 4.

This completes the proof. ■

Note that Theorem 2.2 holds under the conditions $m = n - 1$, or $n = 6$ and $m \geq 10$, or $n = 7$ and $m \neq 8$, or $m \geq n \geq 8$, since for the other pairs of n and m the extremal degree sequences obtained in Theorem 2.2 are not graphic. It is easy to check that for $n = 1, 2, 3$, and for $n \geq 4$ and $m = \binom{n}{2} - 1$ or $m = \binom{n}{2}$ the (n, m) -graph is unique. For $n = 4$ and $m = 4$, or $n = 5$ and $5 \leq m \leq 8$, or $n = 6$ and $6 \leq m \leq 9$, or $n = 7$ and $m = 8$ we can characterize the extremal graphs by examining all possible degree sequences. These extremal graphs are depicted in Figure 2.3 (minimum ones for $0 < \alpha < 1$, maximum ones for $\alpha < 0$ or $\alpha > 1$, except for $n = 5$ and $m = 7$, in which case (a) is the minimum graph for $0 < \alpha < 1$ and maximum graph for $\alpha < 0$ or $1 < \alpha < 2$, and (b) is the maximum graph for $\alpha \geq 2$).




Figure 2.2 Molecular graphs with extremal zeroth-order general Randić index, having four or fewer vertices of degree 4.




Figure 2.3 Some graphs with extremal zeroth-order general Randić index; for details see text.

References

- [1] O. Araujo, J. A. de la Peña, Some bounds for the connectivity index of a chemical graph, *J. Chem. Inf. Comput. Sci.* **38** (1998) 827–831.
- [2] O. Araujo, J. Rada, Randić index and lexicographic order, *J. Math. Chem.* **27** (2000) 201–212.
- [3] B. Bollobás, P. Erdős, Graphs of extremal weights, *Ars Combin.* **50** (1998) 225–233.
- [4] G. Caporossi, I. Gutman, P. Hansen, L. Pavlović, Graphs with maximum connectivity index, *Comput. Biol. Chem.* **27** (2003) 85–90.
- [5] L. H. Clark, J.W. Moon, On the general Randić index for certain families of trees, *Ars Combin.* **54** (2000) 223–235.
- [6] M. Fischermann, A. Hoffmann, D. Rautenbach, L. Volkmann, A linear-programming approach to the generalized Randić index, *Discr. Appl. Math.* **128** (2003) 375–385.
- [7] S. P. Gupta, Quantitative structure–activity relationship studies on local anesthetics, *Chem. Rev.* **91** (1991) 1109–1119.
- [8] I. Gutman, O. Araujo, D. A. Morales, Bounds for the Randić connectivity index, *J. Chem. Inf. Comput. Sci.* **40** (2000) 572–579.
- [9] I. Gutman, O. Araujo, D. A. Morales, Estimating the connectivity index of a saturated hydrocarbon, *Indian J. Chem.* **39A** (2000) 381–385.
- [10] I. Gutman, O. Miljković, Molecules with smallest connectivity indices, *MATCH Commun. Math. Comput. Chem.* **41** (2000) 57–70.

- [11] I. Gutman, O. Miljković, G. Caporossi, P. Hansen, Alkanes with small and large Randić connectivity index *Chem. Phys. Lett.* **306** (1999) 366–372.
- [12] P. Hansen, H. Mélot, Variable neighborhood search for extremal graphs 6: Analyzing bounds for the connectivity index, *J. Chem. Inf. Comput. Sci.* **43** (2003) 1–14.
- [13] Y. Hu, X. Li, Y. Yuan, Trees with minimum general Randić index, *MATCH Commun. Math. Comput. Chem.* **52** (2004) 119–128.
- [14] Y. Hu, X. Li, Y. Yuan, Trees with maximum general Randić index, *MATCH Commun. Math. Comput. Chem.* **52** (2004) 129–146.
- [15] L. B. Kier, L. H. Hall, *Molecular Connectivity in Chemistry and Drug Research*, Academic Press, New York, 1976.
- [16] L. B. Kier, L. H. Hall, The nature of structure-activity relationships and their relation to molecular connectivity, *Europ. J. Med. Chem.* **12** (1977) 307–312.
- [17] L. B. Kier, L. H. Hall, *Molecular Connectivity in Structure-Analysis*, Research Studies Press, Wiley, Chichester, UK, 1986.
- [18] R. Lang, X. Li, S. Zhang, Inverse problem for Zagreb index of molecular graphs, *Appl. Math. J. Chinese Univ.* **A18** (2003) 487–493 (in Chinese).
- [19] X. Li, X. Q. Wang, B. Wei, On the lower and upper bounds for general Randić index of chemical (n, m) -graphs, *MATCH Commun. Math. Comput. Chem.* **52** (2004) 157–166.
- [20] X. Li, Y. Yang, Sharp bounds for the general Randić index, *MATCH Commun. Math. Comput. Chem.* **51** (2004) 155–166.
- [21] X. Li, H. Zhao, Trees with the first three smallest and largest generalized topological indices, *MATCH Commun. Math. Comput. Chem.* **50** (2004) 57–62.
- [22] X. Li, J. Zheng, A unified approach to the extremal trees for different indices, *MATCH Commun. Math. Comput. Chem.* **54** (2005) 195–208.
- [23] L. Pavlović, Maximal value of the zeroth-order Randić index, *Discr. Appl. Math.* **127** (2003) 615–626.
- [24] L. Pogliani, From molecular connectivity indices to semiempirical connectivity terms: Recent trends in graph theoretical descriptors, *Chem. Rev.* **100** (2000) 3827–3858.
- [25] M. Randić, On characterization of molecular branching, *J. Am. Chem. Soc.* **97** (1975) 6609–6615.
- [26] M. Randić, The connectivity index 25 years after, *J. Mol. Graphics Modell.* **20** (2001) 19–35.