

ON SOME CONTINUED FRACTION EXPANSIONS OF THE ROGERS-RAMANUJAN TYPE

NANCY S. S. GU[†] AND HELMUT PRODINGER^{*}

ABSTRACT. By guessing the relative quantities and proving the recursive relation, we present some continued fraction expansions of the Rogers-Ramanujan type. Meanwhile, we also give some J -fraction expansions for the q -tangent and q -cotangent functions.

1. INTRODUCTION

In [27], the second author studied the functions

$$F(z) = \sum_{n \geq 0} \frac{(-1)^n z^n}{[2n+1]_q!} q^{dn^2},$$

$$G(z) = \sum_{n \geq 0} \frac{(-1)^n z^n}{[2n]_q!} q^{dn^2},$$

where

$$[n]_q := \frac{1 - q^n}{1 - q}, \quad [n]_q! := [1]_q [2]_q \cdots [n]_q,$$

and gave some continued fraction expansions of the q -tangent and q -cotangent functions for $d = 0, 1, 2$ in the following forms:

$$\frac{zF(z)}{G(z)} = \frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{a_2 + \frac{z}{\ddots}}}}, \tag{1.1}$$

Date: April 4, 2011.

1991 Mathematics Subject Classification. 33D15, 11A55.

Key words and phrases. continued fraction, Rogers-Ramanujan type, q -tangent function, q -cotangent function.

[†] The first author was supported by the National Natural Science Foundation of China, the PCSIRT Project of the Ministry of Education, and the Specialized Research Fund for the Doctoral Program of Higher Education of China (200800551042).

^{*} The second author was supported by International Science and Technology Agreement (Grant 67215).

$$\frac{zG(z)}{F(z)} = \frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{a_2 + \ddots}}}, \quad (1.2)$$

which are variants of Jackson's [17]. Some special cases were considered in [11, 25, 26].

In [27], the second author also discussed a continued fraction identity of Ramanujan and the celebrated Rogers-Ramanujan continued fraction expansion and companions. Some related kinds of the continued fraction identities of Ramanujan were widely studied in the literature, see [2–8, 13–16, 22, 23]. Recently, in [9], Bowman et al. used two q -continued fractions given in [8] to derive two general transformations, and then obtained m -versions of several identities of Rogers-Ramanujan-Slater type. Inspired by the idea in [27], we find that the method which resembles the method due to Viskovatoff [20, Pages 27-28] can be used for some Rogers-Ramanujan type functions, and we find nice continued fraction expansions for them. By applying a sequence of similarity transformations, the continued fraction expansions we derived through this method can be transformed into much more elegant continued fractions. Meanwhile, some special cases of these continued fraction expansions may lead to representations of infinite q -products as q -continued fractions. For the Rogers-Ramanujan type identities, Sills [34] gave an annotated and cross-referenced version of Slater's list [35] of the Rogers-Ramanujan type identities. When we refer to Slater's list in this paper, we are referring to Sills' version.

In Section 2, we focus on the continued fraction expansions in the following form:

$$\frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{\ddots}}},$$

and discuss this kind of continued fraction expansions for $zF(z)/G(z)$ and its companion $zG(z)/F(z)$ of the Rogers-Ramanujan type, as in (1.1) and (1.2).

In Section 3, we find that our method can be used to give an elementary proof for a continued fraction identity due to Ramanujan which was proved by Andrews in [3].

Section 4 is devoted to expansions of the form:

$$\frac{z^2}{a_0 + b_0z + \frac{z^2}{a_1 + b_1z + \frac{z^2}{\ddots}}},$$

and we study this kind (“ J -fraction”) of continued fraction identities for $z^2F(z)/G(z)$ and $z^2G(z)/F(z)$ of the Rogers-Ramanujan type. We also give some expansions for the q -tangent and q -cotangent functions. Recently, Shin and Zeng in [33] proved a similar

ON SOME CONTINUED FRACTION EXPANSIONS OF THE ROGERS-RAMANUJAN TYPE 3
kind of the continued fraction expansions which were used to give a unified proof of Josuat-Vergès' recent q -analogues of two identities due to Euler and Roselle.

As usual, we follow the notation and terminology in [12]. For $|q| < 1$, the q -shifted factorial is defined by

$$(a; q)_\infty = \prod_{k=0}^{\infty} (1 - aq^k) \quad \text{and} \quad (a; q)_n = \frac{(a; q)_\infty}{(aq^n; q)_\infty}, \text{ for } n \in \mathbb{C}.$$

The main results in this paper are summarized in Table 1 and Table 2.

We don't claim that neither the method nor all the results are original. However, we try to be systematic within the context of the Rogers-Ramanujan type identities and we are confident that a fair share of our results are indeed new.

2. $zF(z)/G(z)$ AND $zG(z)/F(z)$

For the continued fraction of the form

$$\frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{\ddots}}},$$

we briefly state the approach as follows.

Let

$$\frac{zF(z)}{G(z)} = \frac{z}{N_0} = \frac{z}{a_0 + \frac{z}{N_1}} = \frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{N_2}}} = \dots$$

and

$$N_i = \frac{r_i}{s_i}.$$

Then we have

$$N_i = a_i + \frac{z}{N_{i+1}}.$$

That is to say,

$$\frac{r_i}{s_i} = a_i + \frac{zs_{i+1}}{r_{i+1}}.$$

Setting $r_{i+1} = s_i$, we have

$$zs_{i+1} = s_{i-1} - a_i s_i, \tag{2.1}$$

where the initial conditions are

$$s_{-1} = G(z) \quad \text{and} \quad s_0 = F(z).$$

Therefore, if we guess the number a_k and the power series s_k , and prove the recurrence relation (2.1) by induction, then we prove the continued fraction identities. Since the proof is a routine computation, we only show the proof for the first example.

$F(z)$ & $G(z)$	Expansions	Theorems
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n(n+1)/2} (-u; q)_n}{(q; q)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n(n-1)/2} (-u; q)_n}{(q; q)_n}$	$zF(z)/G(z)$	Theorem 2.1
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q^4; q^4)_n}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.2
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+4n}}{(q^4; q^4)_n}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.3
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2} (-q; q^2)_n}{(q^2; q^2)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n} (-q; q^2)_n}{(q^2; q^2)_n}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.4
$F(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n+1}}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.5
$F(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n}}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.6
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n+1}}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.7
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n}}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.8
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n+1}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n}}$	$zF(z)/G(z)$ $zG(z)/F(z)$	Theorem 2.9
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n} (-q; q^2)_n}{(q^4; q^4)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2} (-q; q^2)_n}{(q^4; q^4)_n}$	$zF(z)/G(z)$	Theorem 2.10

TABLE 1. Continued fraction expansions of $zF(z)/G(z)$ and $zG(z)/F(z)$

2.1. **A.8/A.13 type.** First, we consider the continued fraction expansions of the Rogers-Ramanujan type functions in the identities A.8 and A.13 in Slater's list which are stated as follows.

Identity A.8 (Gauss-Lebesgue [21]):

$$\sum_{n=0}^{\infty} \frac{(-q; q)_n q^{n(n+1)/2}}{(q; q)_n} = \frac{(q^4; q^4)_{\infty}}{(q; q)_{\infty}}.$$

$F(z)$ & $G(z)$	Expansions	Theorems
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_n}$	$z^2 F(z)/G(z)$ $z^2 G(z)/F(z)$	Theorem 4.1
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_n}$	$z^2 G(z)/F(z)$	Theorem 4.2
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q^4; q^4)_n}$	$z^2 F(z)/G(z)$ $z^2 G(z)/F(z)$	Theorem 4.3
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+4n}}{(q^4; q^4)_n}$	$z^2 F(z)/G(z)$ $z^2 G(z)/F(z)$	Theorem 4.4
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2} (-q; q^2)_n}{(q^2; q^2)_n}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n} (-q; q^2)_n}{(q^2; q^2)_n}$	$z^2 G(z)/F(z)$	Theorem 4.5
$F(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n+1}}$	$z^2 G(z)/F(z)$ $z^2 F(z)/G(z)$	Theorem 4.6 Theorem 4.13
$F(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n}}$	$z^2 G(z)/F(z)$	Theorem 4.7
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n+1}}$	$z^2 G(z)/F(z)$	Theorem 4.8
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n}}$	$z^2 F(z)/G(z)$ $z^2 G(z)/F(z)$	Theorem 4.9
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n+1}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n}}$	$z^2 F(z)/G(z)$	Theorem 4.10
$F(z) = \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n+1}}$ $G(z) = \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n}}$	$z^2 F(z)/G(z)$ $z^2 G(z)/F(z)$	Theorem 4.11 Theorem 4.14
$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n+1}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}$	$z^2 F(z)/G(z)$	Theorem 4.12
$F(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n+1}}$ $G(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}$	$z^2 G(z)/F(z)$	Theorem 4.15
$F(z) = \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n+1}}$ $G(z) = \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n}}$	$z^2 G(z)/F(z)$	Theorem 4.16

 TABLE 2. Continued fraction expansions of $z^2 F(z)/G(z)$ and $z^2 G(z)/F(z)$

Identity A.13 (Slater [35]):

$$\sum_{n=0}^{\infty} \frac{(-q; q)_n q^{n(n-1)/2}}{(q; q)_n} = \frac{(q^4; q^4)_{\infty}}{(q; q)_{\infty}} + \frac{(-q; q^2)_{\infty}}{(q; q^2)_{\infty}}.$$

In this paper, we mainly focus on the Rogers-Ramanujan type identities. But we claim that the method used in this paper may lead to more general results, and we give an example in the following theorem.

Theorem 2.1. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n(n+1)/2} (-u; q)_n}{(q; q)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n(n-1)/2} (-u; q)_n}{(q; q)_n}.$$

For the continued fraction expansion

$$\frac{zF(z)}{G(z)} = \frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{\ddots}}}, \quad (2.2)$$

we have

$$a_{2k} = \frac{(-u; q)_k}{u^k q^{k(k+1)/2}}, \quad a_{2k+1} = \frac{u^k q^{k(k-1)/2}}{(-u; q)_{k+1}},$$

$$s_{2k} = q^{k^2} \sum_{n \geq 0} \frac{z^n u^k q^{n(n+2k+1)/2} (-uq^k; q)_n}{(q; q)_n},$$

$$s_{2k+1} = q^{k(k+1)/2} \sum_{n \geq 0} \frac{z^n q^{n(n+2k+1)/2} (-u; q)_{n+k+1}}{(q; q)_n}.$$

The continued fraction expansion (2.2) can be transformed into the following forms:

$$\frac{zF(z)}{G(z)} = \frac{z}{1 + \frac{z(1+u)}{1 + \frac{zuq}{1 + \frac{zq(1+uq)}{1 + \frac{zuq^3}{1 + \dots + \frac{zuq^{2n-1}}{1 + \frac{zq^n(1+uq^n)}{1 + \dots}}}}}}}. \quad (2.3)$$

This continued fraction can be obtained from [5, p. 144, Entry 6.2.1] by setting $a = z/q$, $\lambda = zu/q$, and $b = 0$. It was also studied (in more general form) in [7, 14].

Setting $u = q$ and $z = 1$ in (2.3), we have

$$\frac{(q^4; q^4)_{\infty}}{(q^4; q^4)_{\infty} + (-q; q^2)_{\infty} (q^2; q^2)_{\infty}} = \frac{1}{1 + \frac{1+q}{1 + \frac{q^2}{1 + \frac{q(1+q^2)}{1 + \frac{q^4}{1 + \dots + \frac{q^{2n}}{1 + \frac{q^n(1+q^{n+1})}{1 + \dots}}}}}}}$$

Now we give the proof by induction.

Proof. According to the expansion (2.2), we know that $F(z) = s_0$ and $G(z) = s_{-1}$. Therefore, we have

$$\begin{aligned}
 s_1 &= \frac{1}{z}(s_{-1} - a_0 s_0) \\
 &= \frac{1}{z} \left(\sum_{n \geq 0} \frac{z^n q^{n(n-1)/2} (-u; q)_n}{(q; q)_n} - \sum_{n \geq 0} \frac{z^n q^{n(n+1)/2} (-u; q)_n}{(q; q)_n} \right) \\
 &= \sum_{n \geq 0} \frac{z^{n-1} q^{n(n-1)/2} (-u; q)_n}{(q; q)_n} (1 - q^n) \\
 &= \sum_{n \geq 1} \frac{z^{n-1} q^{n(n-1)/2} (-u; q)_n}{(q; q)_{n-1}} \\
 &= \sum_{n \geq 0} \frac{z^n q^{n(n+1)/2} (-u; q)_{n+1}}{(q; q)_n}.
 \end{aligned}$$

$$\begin{aligned}
 s_2 &= \frac{1}{z}(s_0 - a_1 s_1) \\
 &= \frac{1}{z} \left(\sum_{n \geq 0} \frac{z^n q^{n(n+1)/2} (-u; q)_n}{(q; q)_n} - \frac{1}{1+u} \sum_{n \geq 0} \frac{z^n q^{n(n+1)/2} (-u; q)_{n+1}}{(q; q)_n} \right) \\
 &= \sum_{n \geq 0} \frac{z^{n-1} q^{n(n+1)/2} (-uq; q)_{n-1}}{(q; q)_n} (1 + u - 1 - uq^n) \\
 &= \sum_{n \geq 1} \frac{z^{n-1} uq^{n(n+1)/2} (-uq; q)_{n-1}}{(q; q)_{n-1}} \\
 &= \sum_{n \geq 0} \frac{z^n uq^{(n+1)(n+2)/2} (-uq; q)_n}{(q; q)_n}.
 \end{aligned}$$

Next, we show for all n that the recurrence relation (2.1) holds. In this case, we need to prove the following two relations:

$$[z^n](s_{2k-1} - a_{2k} s_{2k}) = [z^{n-1}] s_{2k+1}, \quad (2.4)$$

$$[z^n](s_{2k} - a_{2k+1} s_{2k+1}) = [z^{n-1}] s_{2k+2}. \quad (2.5)$$

For the first relation (2.4):

$$\begin{aligned}
 [z^n](s_{2k-1} - a_{2k} s_{2k}) &= q^{k(k-1)/2} \frac{q^{n(n+2k-1)/2} (-u; q)_{n+k}}{(q; q)_n} \\
 &\quad - \frac{(-u; q)_k}{u^k q^{k(k+1)/2}} q^{k^2} \frac{u^k q^{n(n+2k+1)/2} (-uq^k; q)_n}{(q; q)_n} \\
 &= \frac{q^{n(n-1)/2 + nk + k(k-1)/2} (-u; q)_{n+k}}{(q; q)_{n-1}}.
 \end{aligned}$$

On the other hand,

$$[z^{n-1}]s_{2k+1} = \frac{q^{n(n-1)/2+nk+k(k-1)/2}(-u; q)_{n+k}}{(q; q)_{n-1}},$$

which is the same.

For the second relation (2.5):

$$\begin{aligned} [z^n](s_{2k} - a_{2k+1}s_{2k+1}) &= q^{k^2} \frac{u^k q^{n(n+2k+1)/2}(-uq^k; q)_n}{(q; q)_n} \\ &\quad - \frac{u^k q^{k(k-1)/2}}{(-u; q)_{k+1}} q^{k(k+1)/2} \frac{q^{n(n+2k+1)/2}(-u; q)_{n+k+1}}{(q; q)_n} \\ &= \frac{u^{k+1} q^{n(n+1)/2+nk+k^2+k}(-uq^{k+1}; q)_{n-1}}{(q; q)_{n-1}}. \end{aligned}$$

On the other hand,

$$[z^{n-1}]s_{2k+2} = \frac{u^{k+1} q^{n(n+1)/2+kn+k^2+k}(-uq^{k+1}; q)_{n-1}}{(q; q)_{n-1}},$$

which is the same. □

2.2. A.16/A.20 type. The identities A.16 and A.20 in Slater's list are stated as follows.

Identity A.16 (Rogers [30]):

$$\sum_{n=0}^{\infty} \frac{q^{n^2+2n}}{(q^4; q^4)_n} = \frac{1}{(q^2, q^3; q^5)_{\infty} (-q^2; q^2)_{\infty}}.$$

Identity A.20 (Rogers [30]):

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q^4; q^4)_n} = \frac{1}{(q, q^4; q^5)_{\infty} (-q^2; q^2)_{\infty}}.$$

According to the recurrence relation (2.1), we have the following theorem.

Theorem 2.2. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q^4; q^4)_n}.$$

(1) *For the continued fraction expansion*

$$\frac{zF(z)}{G(z)} = \frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{\ddots}}},$$

we have

$$a_{2k} = (-1)^k (1 + q^{4k}) q^{2k^2 - 2k}, \quad k \geq 1, \quad a_0 = 1,$$

$$a_{2k+1} = (-1)^{k-1}(1 + q^{4k+2})q^{-2k^2-4k-1}.$$

$$s_{2k} = \sum_{n \geq 0} \frac{z^n q^{(n+k)^2}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k}},$$

$$s_{2k+1} = (-1)^{k+1} q^{3k^2+4k+1} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k+1}}.$$

$$\begin{aligned} \frac{zF(z)}{G(z)} &= \frac{z}{1} - \frac{zq}{1+q^2} + \frac{zq}{1+q^4} - \frac{zq^7}{1+q^6} + \frac{zq^3}{1+q^8} - \frac{zq^{13}}{1+q^{10}} \\ &\quad + \cdots + \frac{zq^{2n-1}}{1+q^{4n}} - \frac{zq^{6n+1}}{1+q^{4n+2}} + \cdots. \end{aligned} \quad (2.6)$$

Setting $z = 1$ in (2.6), we have

$$\begin{aligned} \frac{(q^2, q^3; q^5)}{(q, q^4; q^5)} &= \frac{1}{1} - \frac{q}{1+q^2} + \frac{q}{1+q^4} - \frac{q^7}{1+q^6} + \frac{q^3}{1+q^8} - \frac{q^{13}}{1+q^{10}} \\ &\quad + \cdots + \frac{q^{2n-1}}{1+q^{4n}} - \frac{q^{6n+1}}{1+q^{4n+2}} + \cdots. \end{aligned}$$

(2) For the continued fraction expansion

$$\frac{zG(z)}{F(z)} = \frac{z}{a_0 + \frac{z}{a_1 + \frac{z}{\ddots}}},$$

we have

$$\begin{aligned} a_{2k} &= (-1)^k (1 + q^{4k}) q^{-2k^2-2k}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= (-1)^k (1 + q^{4k+2}) q^{2k^2-1}. \end{aligned}$$

$$s_{2k} = (-1)^k q^{3k^2+2k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k}},$$

$$s_{2k+1} = q^{(k+1)^2} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k+1}}.$$

$$\begin{aligned} \frac{zG(z)}{F(z)} &= \frac{z}{1} + \frac{zq}{1+q^2} - \frac{zq^5}{1+q^4} + \frac{zq^3}{1+q^6} - \frac{zq^{11}}{1+q^8} + \cdots + \frac{zq^{2n+1}}{1+q^{4n+2}} - \frac{zq^{6n+5}}{1+q^{4n+4}} + \cdots. \end{aligned} \quad (2.7)$$

This continued fraction can be obtained from [6, p. 41, Eq. (14.1)].

Setting $z = 1$ in (2.7), we have

$$\frac{(q, q^4; q^5)}{(q^2, q^3; q^5)} = \frac{1}{1} + \frac{q}{1+q^2} - \frac{q^5}{1+q^4} + \frac{q^3}{1+q^6} - \frac{q^{11}}{1+q^8} + \cdots + \frac{q^{2n+1}}{1+q^{4n+2}} - \frac{q^{6n+5}}{1+q^{4n+4}} + \cdots.$$

We also find a variant of the above theorem.

Theorem 2.3. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+4n}}{(q^4; q^4)_n}.$$

(1) *For the continued fraction expansion of $zF(z)/G(z)$, we have*

$$\begin{aligned} a_{2k} &= (-1)^k (1 + q^{4k-2}) q^{2k^2-4k+2}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= (-1)^{k-1} (1 + q^{4k}) q^{-2k^2-2k-1}, \quad k \geq 1, \quad a_1 = -\frac{1}{q}. \end{aligned}$$

$$\begin{aligned} s_{2k} &= \sum_{n \geq 0} \frac{z^n q^{(n+k)^2}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k-1}}, \quad k \geq 1, \\ s_{2k+1} &= (-1)^{k+1} q^{3k^2+2k+1} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k}}. \end{aligned}$$

$$\begin{aligned} \frac{zF(z)}{G(z)} &= \frac{z}{1} - \frac{zq}{1+q^2} + \frac{zq}{1+q^4} - \frac{zq^5}{1+q^6} + \frac{zq^3}{1+q^8} - \frac{zq^{11}}{1+q^{10}} \\ &\quad + \dots + \frac{zq^{2n+1}}{1+q^{4n+2}} - \frac{zq^{6n+5}}{1+q^{4n+4}} + \dots \end{aligned}$$

(2) *For the continued fraction expansion of $zG(z)/F(z)$, we have*

$$\begin{aligned} a_{2k} &= (-1)^k (1 + q^{4k-2}) q^{-2k^2}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= (-1)^k (1 + q^{4k}) q^{2k^2-2k-1}, \quad k \geq 1, \quad a_1 = \frac{1}{q}. \end{aligned}$$

$$\begin{aligned} s_{2k} &= (-1)^k q^{3k^2} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k-1}}, \quad k \geq 1, \\ s_{2k+1} &= \sum_{n \geq 0} \frac{z^n q^{(n+k+1)^2}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k}}. \end{aligned}$$

$$\frac{zG(z)}{F(z)} = \frac{z}{1} + \frac{zq}{1+q^2} + \frac{zq^3}{1+q^4} - \frac{zq^9}{1+q^6} + \dots + \frac{zq^{2n+1}}{1+q^{4n}} - \frac{zq^{6n+3}}{1+q^{4n+2}} + \dots$$

2.3. A.34/A.36 type. The identities A.34 and A.36 in Slater's list are stated as follows.

Identity A.34 (Slater [35]): The analytic version of the second Göllnitz-Gordon partition identity.

$$\sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^{n^2+2n}}{(q^2; q^2)_n} = \frac{1}{(q^3, q^4, q^5; q^8)_{\infty}}.$$

Identity A.36 (Slater [35]): The analytic version of the first Göllnitz-Gordon partition identity.

$$\sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^{n^2}}{(q^2; q^2)_n} = \frac{1}{(q, q^4, q^7; q^8)_{\infty}}.$$

Theorem 2.4. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2} (-q; q^2)_n}{(q^2; q^2)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n} (-q; q^2)_n}{(q^2; q^2)_n}.$$

(1) *For the continued fraction expansion of $zF(z)/G(z)$, we have*

$$\begin{aligned} a_{2k} &= -\frac{(-q; q^2)_k q^{k^2-1}}{m_k m_{k-1}}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= -\frac{m_k^2}{(-q; q^2)_{k+1} q^{k^2+4k+1}}. \\ s_{2k} &= \sum_{n \geq 0} \frac{z^n q^{(n+k)^2} (-q^{2k+1}; q^2)_n (m_k - (1 - q^{2n}) q^{2k+1} m_{k-1})}{(q^2; q^2)_n}, \\ s_{2k+1} &= -q^{2k^2+4k+1} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)} (-q; q^2)_{n+k+1}}{(q^2; q^2)_n m_k}, \end{aligned}$$

where

$$m_k = (-q^3; q^2)_k \sum_{i=0}^k \frac{q^{i^2}}{(-q^3; q^2)_i},$$

which satisfy the following recurrence relation:

$$m_k = (1 + q^{2k+1}) m_{k-1} + q^{k^2}.$$

$$\begin{aligned} \frac{zF(z)}{G(z)} &= \frac{z}{1} - \frac{zq(1+q)}{1} + \frac{zqm_1}{1} + \frac{zq^6(1+q^3)}{m_1} + \frac{zq^3m_2}{1} + \frac{zq^{10}(1+q^5)m_1}{m_2} \\ &\quad + \cdots + \frac{zq^{4n+2}(1+q^{2n+1})m_{n-1}}{m_n} + \frac{zq^{2n+1}m_{n+1}}{1} + \cdots. \end{aligned} \quad (2.8)$$

Setting $z = 1$ in (2.8), we have

$$\begin{aligned} \frac{(q^3, q^5; q^8)_{\infty}}{(q, q^7; q^8)_{\infty}} &= \frac{1}{1} - \frac{q(1+q)}{1} + \frac{qm_1}{1} + \frac{q^6(1+q^3)}{m_1} + \frac{q^3m_2}{1} + \frac{q^{10}(1+q^5)m_1}{m_2} \\ &\quad + \cdots + \frac{q^{4n+2}(1+q^{2n+1})m_{n-1}}{m_n} + \frac{q^{2n+1}m_{n+1}}{1} + \cdots. \end{aligned}$$

(2) *For the continued fraction expansion of $zG(z)/F(z)$, we have*

$$\begin{aligned} a_{2k} &= \frac{(-q; q^2)_k}{q^{k^2+2k}}, \quad a_{2k+1} = \frac{q^{k^2-1}}{(-q; q^2)_{k+1}}. \\ s_{2k} &= q^{2k^2+2k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)} (-q^{2k+1}; q^2)_n}{(q^2; q^2)_n}, \quad s_{2k+1} = \sum_{n \geq 0} \frac{z^n q^{(n+k+1)^2} (-q; q^2)_{n+k+1}}{(q^2; q^2)_n}. \end{aligned}$$

$$\frac{zG(z)}{F(z)} = \frac{z}{1 + \frac{zq(1+q)}{1 + \frac{zq^4}{1 + \frac{zq^3(1+q^3)}{1 + \frac{zq^8}{1 + \frac{zq^5(1+q^5)}{1 + \frac{zq^{4n}}{1 + \frac{zq^{2n+1}(1+q^{2n+1})}{1 + \dots}}}}}}}}}} \quad (2.9)$$

The above continued fraction can be obtained from [5, p. 144, Entry 6.2.1] by setting $q \rightarrow q^2$, $a = z/q$, $\lambda = z$, and $b = 0$.

Setting $z = 1$ in (2.9), we have

$$\frac{(q, q^7; q^8)_\infty}{(q^3, q^5; q^8)_\infty} = \frac{1}{1 + \frac{q(1+q)}{1 + \frac{q^4}{1 + \frac{q^3(1+q^3)}{1 + \frac{q^8}{1 + \frac{q^5(1+q^5)}{1 + \frac{q^{4n}}{1 + \frac{q^{2n+1}(1+q^{2n+1})}{1 + \dots}}}}}}}}}} \quad (2.9)$$

This continued fraction which is called the Ramanujan-Göllnitz-Gordon continued fraction appears in [5, p. 154, Corollary 6.2.8], [6, p. 50, Entry 22], and [29]. It was widely studied in the literature, see [1, 2, 10, 13, 28, 31, 32].

2.4. A.38/A.39 type. The identities A.38 and A.39 in Slater's list are stated as follows.

Identity A.38 (Slater [35]):

$$\sum_{n=0}^{\infty} \frac{q^{2n^2+2n}}{(q; q)_{2n+1}} = \frac{(q^3, q^5, q^8; q^8)_\infty (q^2, q^{14}; q^{16})_\infty}{(q; q)_\infty}.$$

Identity A.39 (Jackson [18]):

$$\sum_{n=0}^{\infty} \frac{q^{2n^2}}{(q; q)_{2n}} = \frac{(q, q^7, q^8; q^8)_\infty (q^6, q^{10}; q^{16})_\infty}{(q; q)_\infty}.$$

Theorem 2.5. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n+1}}.$$

(1) *For the continued fraction expansion of $zF(z)/G(z)$, we have*

$$a_{2k} = -\frac{(1 - q^{4k+1})(1 - q^2)^2 q^{2k-2}}{(1 - q^{2k-1})(1 - q^{2k})(1 - q^{2k+1})(1 - q^{2k+2})}, \quad k \geq 1, \quad a_0 = \frac{1}{1 - q},$$

$$a_{2k+1} = -\frac{(1 - q^{4k+3})(1 - q^{2k+1})^2 (1 - q^{2k+2})^2}{(1 - q^2)^2 q^{6k+2}}.$$

$$s_{2k} = \sum_{n \geq 0} \frac{z^n q^{2(n+k)^2} (1 - q^2 + q^{2n+2} - q^{2n+2k+1} - q^{2n+2k+2} + q^{2n+4k+3})}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k} (1 - q^2)},$$

$$s_{2k+1} = -q^{2k^2+6k+2} \sum_{n \geq 0} \frac{z^n q^{2n^2+4n(k+1)} (1 - q^2)}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k+1} (1 - q^{2k+1})(1 - q^{2k+2})}.$$

$$\frac{zF(z)}{G(z)} = \frac{z(1-q)}{1 - (1-q)(1-q^2)(1-q^3) + \frac{zq^2(1-q^2)}{1-q^5} + (1-q^3)(1-q^4)(1-q^7)} + \dots + \frac{zq^{4n-2}(1-q^{2n+1})(1-q^{2n+2})}{1-q^{4n+1}} + \frac{zq^{4n+4}(1-q^{2n-1})(1-q^{2n})}{(1-q^{2n+1})(1-q^{2n+2})(1-q^{4n+3})} + \dots \quad (2.10)$$

Setting $z = 1$ in (2.10), we have

$$\frac{(q, q^7; q^8)_\infty (q^6, q^{10}; q^{16})_\infty}{(q^3, q^5; q^8)_\infty (q^2, q^{14}; q^{16})_\infty} = \frac{1-q}{1 - (1-q)(1-q^2)(1-q^3) + \frac{q^2(1-q^2)}{1-q^5} + \frac{q^2(1-q^3)(1-q^4)}{1-q^5}} + \frac{q^8(1-q)(1-q^2)}{(1-q^3)(1-q^4)(1-q^7)} + \dots + \frac{q^{4n-2}(1-q^{2n+1})(1-q^{2n+2})}{1-q^{4n+1}} + \frac{q^{4n+4}(1-q^{2n-1})(1-q^{2n})}{(1-q^{2n+1})(1-q^{2n+2})(1-q^{4n+3})} + \dots$$

(2) For the continued fraction expansion of $zG(z)/F(z)$, we have

$$a_{2k} = \frac{1-q^{4k+1}}{q^{2k}}, \quad a_{2k+1} = \frac{1-q^{4k+3}}{q^{2k+2}}.$$

$$s_{2k} = q^{2k^2+2k} \sum_{n \geq 0} \frac{z^n q^{2n^2+2n(2k+1)}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k}}, \quad s_{2k+1} = \sum_{n \geq 0} \frac{z^n q^{2(n+k+1)^2}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k+1}}.$$

$$\frac{zG(z)}{F(z)} = \frac{z}{1-q} + \frac{zq^2}{1-q^3} + \frac{zq^4}{1-q^5} + \frac{zq^6}{1-q^7} + \dots + \frac{zq^{2n}}{1-q^{2n+1}} + \dots \quad (2.11)$$

Setting $z = 1$ in (2.11), we have

$$\frac{(q^3, q^5; q^8)_\infty (q^2, q^{14}; q^{16})_\infty}{(q, q^7; q^8)_\infty (q^6, q^{10}; q^{16})_\infty} = \frac{1}{1-q} + \frac{q^2}{1-q^3} + \frac{q^4}{1-q^5} + \frac{q^6}{1-q^7} + \dots + \frac{q^{2n}}{1-q^{2n+1}} + \dots$$

Theorem 2.6. Let

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n}}.$$

(1) For the continued fraction expansion of $zF(z)/G(z)$, we have

$$a_{2k} = -\frac{(1-q^{4k-1})(1-q^2)^2 q^{2k-2}}{(1-q^{2k-1}-q^{2k}+q^{4k-3})(1-q^{2k+1}-q^{2k+2}+q^{4k+1})}, \quad k \geq 1, \quad a_0 = 1,$$

$$a_{2k+1} = -\frac{(1-q^{4k+1})(1-q^{2k+1}-q^{2k+2}+q^{4k+1})^2}{(1-q^2)^2 q^{6k+2}}.$$

$$s_{2k} = \sum_{n \geq 0} \frac{z^n q^{2(n+k)^2} (1-q^2 + q^{2n+2} - q^{2n+2k+1} - q^{2n+2k+2} + q^{2n+4k+1})}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k-1} (1-q^2)},$$

$$s_{2k+1} = -q^{2k^2+6k+2} \sum_{n \geq 0} \frac{z^n q^{2n^2+4n(k+1)} (1-q^2)}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k} (1-q^{2k+1}-q^{2k+2}+q^{4k+1})}.$$

$$\frac{zF(z)}{G(z)} = \frac{z}{1 - (1-q)(1-q^2) + \frac{zq^2(1-q^2)}{1-q^3} + (1-q^5)(1-q^3-q^4+q^5) + \dots} + \frac{zq^8(1-q^2)}{(1-q^5)(1-q^3-q^4+q^5) + \dots} + \frac{zq^{4n-2}(1-q^{2n+1}-q^{2n+2}+q^{4n+1})}{1-q^{4n-1}} + \frac{zq^{4n+4}(1-q^{2n-1}-q^{2n}+q^{4n-3})}{(1-q^{4n+1})(1-q^{2n+1}-q^{2n+2}+q^{4n+1})} + \dots$$

(2) For the continued fraction expansion of $zG(z)/F(z)$, we have

$$a_{2k} = \frac{1 - q^{4k-1}}{q^{2k}}, \quad k \geq 1, \quad a_0 = 1, \quad a_{2k+1} = \frac{1 - q^{4k+1}}{q^{2k+2}}.$$

$$s_{2k} = q^{2k^2+2k} \sum_{n \geq 0} \frac{z^n q^{2n^2+2n(2k+1)}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k-1}}, \quad s_{2k+1} = \sum_{n \geq 0} \frac{z^n q^{2(n+k+1)^2}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k}}.$$

$$\frac{zG(z)}{F(z)} = \frac{z}{1} + \frac{zq^2}{1 - q} + \frac{zq^4}{1 - q^3} + \frac{zq^6}{1 - q^5} + \frac{zq^8}{1 - q^7} + \cdots + \frac{zq^{2n}}{1 - q^{2n-1}} + \cdots.$$

This continued fraction can be obtained from [5, p. 160, Theorem 6.4.1] by setting $q \rightarrow q^2$, $a = 0$, $\lambda = z$, $b = -1/q$. Some more general forms were studied in [14, 15], and [8, Theorem 2].

2.5. A.79/A.96 type. The identities A.79 and A.96 in Slater's list are stated as follows.

Identity A.79 (Rogers [30]):

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_{2n}} = \frac{(q^8, q^{12}, q^{20}; q^{20})_{\infty} (-q; q^2)_{\infty}}{(q^2; q^2)_{\infty}}.$$

Identity A.96 (Rogers [30]):

$$\sum_{n=0}^{\infty} \frac{q^{n^2+2n}}{(q; q)_{2n+1}} = \frac{(q^4, q^6, q^{10}; q^{10})_{\infty} (q^2, q^{18}; q^{20})_{\infty}}{(q; q)_{\infty}}.$$

Theorem 2.7. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n+1}}.$$

(1) For the continued fraction expansion of $zF(z)/G(z)$, we have

$$a_{2k} = -\frac{(1 - q^{4k+1})q^{2k^2-k-1}}{(1 - q^{2k^2-k})(1 - q^{2k^2+3k+1})}, \quad k \geq 1, \quad a_0 = \frac{1}{1 - q},$$

$$a_{2k+1} = -\frac{(1 - q^{4k+3})(1 - q^{2k^2+3k+1})^2}{q^{2k^2+5k+1}}.$$

$$s_{2k} = \sum_{n \geq 0} \frac{z^n q^{(n+k)^2} (1 - q^{2n+2k^2+3k+1})}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k}},$$

$$s_{2k+1} = -q^{3k^2+5k+1} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k+1} (1 - q^{2k^2+3k+1})}.$$

$$\begin{aligned} \frac{zF(z)}{G(z)} = & \frac{z(1-q)}{1} - \frac{zq}{(1-q)(1-q^3)} + \frac{zq(1-q^6)}{1-q^5} + \frac{zq^8(1-q)}{(1-q^6)(1-q^7)} \\ & + \cdots + \frac{zq^{2n-1}(1-q^{2n^2+3n+1})}{1-q^{4n+1}} + \frac{zq^{6n+2}(1-q^{2n^2-n})}{(1-q^{4n+3})(1-q^{2n^2+3n+1})} + \cdots \end{aligned} \quad (2.12)$$

Setting $z = 1$ in (2.12), we have

$$\begin{aligned} \frac{(q^8, q^{12}; q^{20})_\infty}{(q^4, q^{16}; q^{20})_\infty} &= \frac{1-q}{1} - \frac{q}{(1-q)(1-q^3)} + \frac{q(1-q^6)}{1-q^5} + \frac{q^8(1-q)}{(1-q^6)(1-q^7)} \\ &\quad + \dots + \frac{q^{2n-1}(1-q^{2n^2+3n+1})}{1-q^{4n+1}} + \frac{q^{6n+2}(1-q^{2n^2-n})}{(1-q^{4n+3})(1-q^{2n^2+3n+1})} + \dots \end{aligned}$$

(2) For the continued fraction expansion of $zG(z)/F(z)$, we have

$$a_{2k} = \frac{1-q^{4k+1}}{q^{2k^2+3k}}, \quad a_{2k+1} = (1-q^{4k+3})q^{2k^2+k-1}.$$

$$s_{2k} = q^{3k^2+3k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q; q)_{2n+1}(q^{2n+3}; q^2)_{2k}}, \quad s_{2k+1} = \sum_{n \geq 0} \frac{z^n q^{(n+k+1)^2}}{(q; q)_{2n+1}(q^{2n+3}; q^2)_{2k+1}}.$$

$$\frac{zG(z)}{F(z)} = \frac{z}{1-q} + \frac{zq}{1-q^3} + \frac{zq^6}{1-q^5} + \frac{zq^3}{1-q^7} + \dots + \frac{zq^{6n}}{1-q^{4n+1}} + \frac{zq^{2n+1}}{1-q^{4n+3}} + \dots \quad (2.13)$$

Setting $z = 1$ in (2.13), we have

$$\frac{(q^4, q^{16}; q^{20})_\infty}{(q^8, q^{12}; q^{20})_\infty} = \frac{1}{1-q} + \frac{q}{1-q^3} + \frac{q^6}{1-q^5} + \frac{q^3}{1-q^7} + \dots + \frac{q^{6n}}{1-q^{4n+1}} + \frac{q^{2n+1}}{1-q^{4n+3}} + \dots.$$

Theorem 2.8. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n}}.$$

(1) For the continued fraction expansion of $zF(z)/G(z)$, we have

$$a_{2k} = -(1-q^{4k-1})q^{2k^2-3k+1}, \quad k \geq 1, \quad a_0 = 1, \quad a_{2k+1} = -\frac{1-q^{4k+1}}{q^{2k^2+3k+1}}.$$

$$\begin{aligned} s_{2k} &= \sum_{n \geq 0} \frac{z^n q^{(n+k)^2}}{(q; q)_{2n+1}(q^{2n+3}; q^2)_{2k-1}}, \\ s_{2k+1} &= -q^{3k^2+3k+1} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q; q)_{2n+1}(q^{2n+3}; q^2)_{2k}}. \end{aligned}$$

$$\frac{zF(z)}{G(z)} = \frac{z}{1} - \frac{zq}{1-q} + \frac{zq}{1-q^3} + \frac{zq^6}{1-q^5} + \frac{zq^3}{1-q^7} + \dots + \frac{zq^{2n-1}}{1-q^{4n-1}} + \frac{zq^{6n}}{1-q^{4n+1}} + \dots.$$

(2) For the continued fraction expansion of $zG(z)/F(z)$, we have

$$a_{2k} = \frac{1-q^{4k-1}}{q^{2k^2+k}}, \quad k \geq 1, \quad a_0 = 1, \quad a_{2k+1} = (1-q^{4k+1})q^{2k^2-k-1}.$$

$$\begin{aligned} s_{2k} &= q^{3k^2+k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)}}{(q; q)_{2n+1}(q^{2n+3}; q^2)_{2k-1}}, \\ s_{2k+1} &= \sum_{n \geq 0} \frac{z^n q^{(n+k+1)^2}}{(q; q)_{2n+1}(q^{2n+3}; q^2)_{2k}}. \end{aligned}$$

$$\frac{zG(z)}{F(z)} = \frac{z}{1} + \frac{zq}{1-q} + \frac{zq^4}{1-q^3} + \frac{zq^3}{1-q^5} + \frac{zq^{10}}{1-q^7} + \dots + \frac{zq^{6n-2}}{1-q^{4n-1}} + \frac{zq^{2n+1}}{1-q^{4n+1}} + \dots.$$

2.6. **A.94/A.99 type.** The identities A.94 and A.99 in Slater's list are stated as follows.

Identity A.94 (Rogers [30]):

$$\sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(q; q)_{2n+1}} = \frac{(q^3, q^7, q^{10}; q^{10})_{\infty} (q^4, q^{16}; q^{20})_{\infty}}{(q; q)_{\infty}}.$$

Identity A.99 (Rogers [30]):

$$\sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(q; q)_{2n}} = \frac{(q, q^9, q^{10}; q^{10})_{\infty} (q^8, q^{12}; q^{20})_{\infty}}{(q; q)_{\infty}}.$$

Theorem 2.9. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n+1}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n}}.$$

(1) *For the continued fraction expansion of $zF(z)/G(z)$, we have*

$$a_{2k} = (1 - q^{4k+1})q^{2k^2-k}, \quad a_{2k+1} = \frac{1 - q^{4k+3}}{q^{2k^2+5k+3}}.$$

$$s_{2k} = q^{k^2+k} \sum_{n \geq 0} \frac{z^n q^{n^2+n(2k+1)}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k}},$$

$$s_{2k+1} = q^{3(k+1)^2} \sum_{n \geq 0} \frac{z^n q^{n^2+n(2k+3)}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k+1}}.$$

$$\frac{zF(z)}{G(z)} = \frac{z}{1-q} + \frac{zq^3}{1-q^3} + \frac{zq^2}{1-q^5} + \frac{zq^9}{1-q^7} + \dots + \frac{zq^{2n}}{1-q^{4n+1}} + \frac{zq^{6n+3}}{1-q^{4n+3}} + \dots. \quad (2.14)$$

Setting $z = 1$ in (2.14), we have

$$\frac{(q^3, q^7; q^{10})_{\infty} (q^4, q^{16}; q^{20})_{\infty}}{(q, q^9; q^{10})_{\infty} (q^8, q^{12}; q^{20})_{\infty}} = \frac{1}{1-q} + \frac{q^3}{1-q^3} + \frac{q^2}{1-q^5} + \frac{q^9}{1-q^7} + \dots + \frac{q^{2n}}{1-q^{4n+1}} + \frac{q^{6n+3}}{1-q^{4n+3}} + \dots.$$

(2) *For the continued fraction expansion of $zG(z)/F(z)$, we have*

$$a_{2k} = -\frac{(1 - q^{4k+1})q^{2k^2-k}}{(1 - q^{2k^2-k})(1 - q^{2k^2+3k+1})}, \quad k \geq 1, \quad a_0 = \frac{1}{1-q},$$

$$a_{2k+1} = -\frac{(1 - q^{4k+3})(1 - q^{2k^2+3k+1})^2}{q^{2k^2+5k+3}}.$$

$$s_{2k} = q^{k^2+k} \sum_{n \geq 0} \frac{z^n q^{n^2+n(2k+1)}(1 - q^{2n+2k^2+3k+1})}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k}},$$

$$s_{2k+1} = -q^{3(k+1)^2} \sum_{n \geq 0} \frac{z^n q^{n^2+n(2k+3)}}{(q; q)_{2n+1} (q^{2n+3}; q^2)_{2k+1} (1 - q^{2k^2+3k+1})}.$$

$$\begin{aligned} \frac{zG(z)}{F(z)} = & \frac{z(1-q)}{1} - \frac{zq^3}{(1-q)(1-q^3)} + \frac{zq^2(1-q^6)}{1-q^5} + \frac{zq^9(1-q)}{(1-q^6)(1-q^7)} \\ & + \dots + \frac{zq^{2n}(1-q^{2n^2+3n+1})}{1-q^{4n+1}} + \frac{zq^{6n+3}(1-q^{2n^2-n})}{(1-q^{4n+3})(1-q^{2n^2+3n+1})} + \dots. \end{aligned} \quad (2.15)$$

Setting $z = 1$ in (2.15), we have

$$\begin{aligned} \frac{(q, q^9; q^{10})_\infty (q^8, q^{12}; q^{20})_\infty}{(q^3, q^7; q^{10})_\infty (q^4, q^{16}; q^{20})_\infty} = & \frac{1-q}{1} - \frac{q^3}{(1-q)(1-q^3)} + \frac{q^2(1-q^6)}{1-q^5} + \frac{q^9(1-q)}{(1-q^6)(1-q^7)} \\ & + \dots + \frac{q^{2n}(1-q^{2n^2+3n+1})}{1-q^{4n+1}} + \frac{q^{6n+3}(1-q^{2n^2-n})}{(1-q^{4n+3})(1-q^{2n^2+3n+1})} + \dots. \end{aligned}$$

2.7. A.25 type. The identity A.25 in Slater's list is stated as follows.

Identity A.25 (Slater [35]):

$$\sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^{n^2}}{(q^4; q^4)_n} = \frac{(q^3, q^3, q^6; q^6)_\infty (-q; q^2)_\infty}{(q^2; q^2)_\infty}. \quad (2.16)$$

Recently, in [24], McLaughlin et al. found a partner to Equation (2.16).

An identity (McLaughlin et al. [24, Eq. (2.7)]):

$$\sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^{n^2+2n}}{(q^4; q^4)_n} = \frac{(q^6; q^6)_\infty}{(q^4; q^4)_\infty (q^3, q^9; q^{12})_\infty}.$$

Theorem 2.10. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n} (-q; q^2)_n}{(q^4; q^4)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2} (-q; q^2)_n}{(q^4; q^4)_n}.$$

For the continued fraction expansion of $zF(z)/G(z)$, we have

$$\begin{aligned} a_{2k} &= \frac{(-q; q^2)_k (1 + q^{4k})}{(q; q^2)_k q^{k^2+2k}}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= \frac{(1 + q^{4k+2})(q; q^2)_k q^{k^2-1}}{(-q; q^2)_{k+1}}, \\ s_{2k} &= q^{2k^2+2k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(k+1)} (-q; q^2)_{n+k} (q; q^2)_k}{(q^2; q^2)_n (-q^2; q^2)_{n+2k} (-q; q^2)_k}, \\ s_{2k+1} &= \sum_{n \geq 0} \frac{z^n q^{(n+k+1)^2} (-q; q^2)_{n+k+1}}{(q^2; q^2)_n (-q^2; q^2)_{n+2k+1}}. \end{aligned}$$

$$\frac{zF(z)}{G(z)} = \frac{z}{1} + \frac{zq(1+q)}{1+q^2} + \frac{zq^4(1-q)}{1+q^4} + \frac{zq^3(1+q^3)}{1+q^6} + \frac{zq^8(1-q^3)}{1+q^8}$$

$$+ \dots + \frac{zq^{4n}(1 - q^{2n-1})}{1 + q^{4n}} + \frac{zq^{2n+1}(1 + q^{2n+1})}{1 + q^{4n+2}} + \dots \quad (2.17)$$

Setting $z = 1$ in (2.17), we have

$$\begin{aligned} \frac{(q, q^5; q^6)_\infty}{(q^3, q^3; q^6)_\infty} &= \frac{1}{1 +} \frac{q(1 + q)}{1 + q^2} + \frac{q^4(1 - q)}{1 + q^4} + \frac{q^3(1 + q^3)}{1 + q^6} + \frac{q^8(1 - q^3)}{1 + q^8} \\ &\quad + \dots + \frac{q^{4n}(1 - q^{2n-1})}{1 + q^{4n}} + \frac{q^{2n+1}(1 + q^{2n+1})}{1 + q^{4n+2}} + \dots \end{aligned}$$

3. A CONTINUED FRACTION IDENTITY OF RAMANUJAN

In [3], Andrews gave a proof of the following “slightly tricky” continued fraction of Ramanujan

$$\frac{zF(z)}{G(z)} = \frac{z}{1 + \frac{zq}{1 + bq + \frac{zq^2}{1 + bq^2 + \frac{zq^3}{\ddots}}}}, \quad (3.1)$$

where

$$F(z) = \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_n (-bq; q)_n}, \quad G(z) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_n (-bq; q)_n}.$$

Later, Bhargava and Adiga proved this identity in [7]. In what follows, applying our method, we can give an elementary proof of this continued fraction identity (3.1).

Proof. In fact, this continued fraction (3.1) fits in the following form:

$$\frac{zF(z)}{G(z)} = \frac{z}{a_0 + \frac{zq}{a_1 + \frac{zq^2}{a_2 + \frac{zq^3}{\ddots}}}}. \quad (3.2)$$

We define

$$\frac{zF(z)}{G(z)} = \frac{z}{N_0} = \frac{z}{a_0 + \frac{zq}{N_1}} = \frac{z}{a_0 + \frac{zq}{a_1 + \frac{zq^2}{N_2}}} = \dots$$

and

$$N_i = \frac{r_i}{s_i}.$$

Then we have

$$N_i = a_i + \frac{zq^{i+1}}{N_{i+1}}.$$

That is to say,

$$\frac{r_i}{s_i} = a_i + \frac{zq^{i+1}s_{i+1}}{r_{i+1}}.$$

Setting $r_{i+1} = s_i$, we have

$$zq^{i+1}s_{i+1} = s_{i-1} - a_i s_i, \quad (3.3)$$

where the initial conditions are

$$s_{-1} = G(z) \quad \text{and} \quad s_0 = F(z).$$

Therefore, if we give the number a_k and the power series s_k , and prove the recurrence relation (3.3) by induction, then we prove the continued fraction identity (3.2).

In this case, we have

$$a_k = 1 + bq^k, \quad k \geq 1, \quad a_0 = 1.$$

$$s_k = \sum_{n \geq 0} \frac{z^n q^{n^2 + n(k+1)}}{(q; q)_n (-bq; q)_{n+k}}.$$

Then we can prove the continued fraction identity (3.1) by induction. \square

4. $z^2 F(z)/G(z)$ AND $z^2 G(z)/F(z)$

In this section, we discuss the following continued fraction expansions:

$$\begin{aligned} \frac{z^2 F(z)}{G(z)} &= \frac{z^2}{a_0 + b_0 z + \frac{z^2}{a_1 + b_1 z + \frac{z^2}{a_2 + b_2 z + \frac{z^2}{\ddots}}}}, \\ \frac{z^2 G(z)}{F(z)} &= \frac{z^2}{a_0 + b_0 z + \frac{z^2}{a_1 + b_1 z + \frac{z^2}{a_2 + b_2 z + \frac{z^2}{\ddots}}}}. \end{aligned} \quad (4.1)$$

Here, we take the continued fraction expansion (4.1) as an example to present the method. For the expansion (4.1), let

$$\frac{z^2 F(z)}{G(z)} = \frac{z^2}{N_0} = \frac{z^2}{a_0 + b_0 z + \frac{z^2}{N_1}} = \frac{z^2}{a_0 + b_0 z + \frac{z^2}{a_1 + b_1 z + \frac{z^2}{N_2}}}$$

and

$$N_i = \frac{r_i}{s_i}.$$

Then we have

$$N_i = a_i + b_i z + \frac{z^2}{N_{i+1}}.$$

That is to say,

$$\frac{r_i}{s_i} = a_i + b_i z + \frac{z^2 s_{i+1}}{r_{i+1}}.$$

Setting $r_{i+1} = s_i$, we have

$$z^2 s_{i+1} = s_{i-1} - (a_i + b_i z) s_i, \quad (4.2)$$

where the initial conditions are

$$s_{-1} = G(z) \quad \text{and} \quad s_0 = F(z).$$

Therefore, if we guess the numbers a_k , b_k , and the power series s_k , and prove the recurrence relation (4.2) by induction, then we prove the continued fraction identity (4.1).

Theorem 4.1. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_n}.$$

(1) *For the continued fraction expansion of $z^2 F(z)/G(z)$, we have*

$$\begin{aligned} a_{2k} &= -\frac{(q^2; q^2)_k^4 (1-q)^2}{(q; q^2)_k^4 (1-q^{2k})(1-q^{2k+1}) q^{4k+1}}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= \frac{(q; q^2)_k^4 (1-q^{2k+1})^3}{(q^2; q^2)_k^4 (1-q^{2k+2})(1-q)^2 q^{4k+2}}, \\ b_{2k} &= -\frac{(q^2; q^2)_k^4 (1-q)^2 (1+q^3 - 2q^{2k+1} - 2q^{2k+3} + q^{4k+2} + q^{4k+3})}{(q; q^2)_k^4 (1-q^{2k})^2 (1-q^{2k+1})^2 q^2}, \quad k \geq 1, \quad b_0 = -q, \\ b_{2k+1} &= \frac{(q; q^2)_k^4 (1-q^{2k+1})^2 (1+q^3 - 2q^{2k+2} - 2q^{2k+4} + q^{4k+4} + q^{4k+5})}{(q^2; q^2)_k^4 (1-q^{2k+2})^2 (1-q)^2 q}, \\ s_{2k} &= (-1)^k q^{4k^2+3k} \sum_{n \geq 0} \frac{z^n q^{n^2+4nk} (q; q^2)_k (q^3; q^2)_{k-1} (1-q + q^{n+1} - q^{n+2k+1})}{(q; q)_n (q^2; q^2)_k^2}, \\ s_{2k+1} &= (-1)^k q^{4k^2+7k+2} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)} (q^2; q^2)_k^2 (1-q + q^{n+1} - q^{n+2k+2})}{(q; q)_n (q; q^2)_{k+1} (q^3; q^2)_k}, \\ \frac{z^2 F(z)}{G(z)} &= \frac{z^2}{1-zq + (1-q)^3(1-q^2) + zq(1-q)^2(1-2q^2+q^3-q^4+q^5)} \\ &\quad \frac{z^2 q^2 (1-q)^2 (1-q^2)^2}{z^2 q^7 (1-q)^4 (1-q^3)^2} \\ &\quad - \frac{(1-q^2)(1-q^3) + zq^3(1-q^3 - 2q^5 + q^6 + q^7)}{z^2 q^{11} (1-q^2)^2 (1-q^3)^2 (1-q^4)^2} \\ &\quad - \frac{(1-q^3)^3 (1-q^4) + zq^5 (1-q^3)^2 (1+q^3 - 2q^4 - 2q^6 + q^8 + q^9)}{z^2 q^{8n-1} (1-q^{2n-1})^4 (1-q^{2n+1})^2} \\ &\quad - \dots - \frac{(1-q^{2n})(1-q^{2n+1}) + zq^{4n-1} (1+q^3 - 2q^{2n+1} - 2q^{2n+3} + q^{4n+2} + q^{4n+3})}{z^2 q^{8n-1} (1-q^{2n-1})^4 (1-q^{2n+1})^2} \end{aligned}$$

$$\begin{aligned}
 & \frac{z^2 q^{8n+3} (1-q^{2n})^2 (1-q^{2n+1})^2 (1-q^{2n+2})^2}{(1-q^{2n+1})^3 (1-q^{2n+2}) + z q^{4n+1} (1-q^{2n+1})^2 (1+q^3 - 2q^{2n+2} - 2q^{2n+4} + q^{4n+4} + q^{4n+5})} - \dots \\
 & \hspace{15em} (4.3)
 \end{aligned}$$

Setting $z = 1$ in (4.3), we have

$$\begin{aligned}
 \frac{(q^2, q^3; q^5)_\infty}{(q, q^4; q^5)_\infty} &= \frac{1}{1-q} \frac{q^2(1-q)^2(1-q^2)^2}{(1-q)^3(1-q^2) + q(1-q)^2(1-2q^2+q^3-q^4+q^5)} \\
 & \quad - \frac{q^7(1-q)^4(1-q^3)^2}{(1-q^2)(1-q^3) + q^3(1-q^3-2q^5+q^6+q^7)} \\
 & \quad - \frac{q^{11}(1-q^2)^2(1-q^3)^2(1-q^4)^2}{(1-q^3)^3(1-q^4) + q^5(1-q^3)^2(1+q^3-2q^4-2q^6+q^8+q^9)} \\
 & \quad - \dots - \frac{q^{8n-1}(1-q^{2n-1})^4(1-q^{2n+1})^2}{(1-q^{2n})(1-q^{2n+1}) + q^{4n-1}(1+q^3-2q^{2n+1}-2q^{2n+3}+q^{4n+2}+q^{4n+3})} \\
 & \quad - \frac{q^{8n+3}(1-q^{2n})^2(1-q^{2n+1})^2(1-q^{2n+2})^2}{(1-q^{2n+1})^3(1-q^{2n+2}) + q^{4n+1}(1-q^{2n+1})^2(1+q^3-2q^{2n+2}-2q^{2n+4}+q^{4n+4}+q^{4n+5})} - \dots
 \end{aligned}$$

(2) For the continued fraction expansion of $z^2 G(z)/F(z)$, we have

$$a_{2k} = q^{-4k}, \quad a_{2k+1} = -q^{-4k-3}.$$

$$b_{2k} = 1+q, \quad k \geq 1, \quad b_0 = q, \quad b_{2k+1} = -\frac{1+q}{q}.$$

$$s_{2k} = (-1)^k q^{4k^2+3k} \sum_{n \geq 0} \frac{z^n q^{n^2+n(4k+1)}}{(q; q)_n},$$

$$s_{2k+1} = (-1)^{k-1} q^{4k^2+7k+3} \sum_{n \geq 0} \frac{z^n q^{n^2+n(4k+3)}}{(q; q)_n}.$$

$$\begin{aligned}
 \frac{z^2 G(z)}{F(z)} &= \frac{z^2}{1+zq} - \frac{z^2 q^3}{1+zq^2(1+q)} - \frac{z^2 q^7}{1+zq^4(1+q)} - \frac{z^2 q^{11}}{1+zq^6(1+q)} \\
 & \quad - \frac{z^2 q^{15}}{1+zq^8(1+q)} - \dots - \frac{z^2 q^{4n-1}}{1+zq^{2n}(1+q)} - \dots \hspace{2em} (4.4)
 \end{aligned}$$

Setting $z = 1$ in (4.4), we have

$$\begin{aligned}
 \frac{(q, q^4; q^5)_\infty}{(q^2, q^3; q^5)_\infty} &= \frac{1}{1+q} - \frac{q^3}{1+q^2(1+q)} - \frac{q^7}{1+q^4(1+q)} - \frac{q^{11}}{1+q^6(1+q)} \\
 & \quad - \frac{q^{15}}{1+q^8(1+q)} - \dots - \frac{q^{4n-1}}{1+q^{2n}(1+q)} - \dots
 \end{aligned}$$

We point out that for the functions $F(z)$ and $G(z)$ in the above theorem, the continued fraction expansions of $zF(z)/G(z)$ and $zG(z)/F(z)$ were provided in [27].

Theorem 4.2. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_n}.$$

For the continued fraction expansion of $z^2G(z)/F(z)$, we have

$$a_{2k} = q^{-4k}, \quad a_{2k+1} = -q^{-4k-5}.$$

$$b_{2k} = q(1+q), \quad b_{2k+1} = -\frac{1+q}{q^2}.$$

$$s_{2k} = (-1)^k q^{4k^2+5k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)}}{(q; q)_n},$$

$$s_{2k+1} = (-1)^{k-1} q^{4k^2+9k+5} \sum_{n \geq 0} \frac{z^n q^{n^2+4n(k+1)}}{(q; q)_n}.$$

$$\begin{aligned} \frac{z^2G(z)}{F(z)} = & \frac{z^2}{1+zq(1+q)} - \frac{z^2q^5}{1+zq^3(1+q)} - \frac{z^2q^9}{1+zq^5(1+q)} - \frac{z^2q^{13}}{1+zq^7(1+q)} \\ & - \frac{z^2q^{17}}{1+zq^9(1+q)} - \dots - \frac{z^2q^{4n+1}}{1+zq^{2n+1}(1+q)} - \dots \end{aligned}$$

For the functions $F(z)$ and $G(z)$ in the above theorem, the continued fraction expansion of $zG(z)/F(z)$ was provided in [27].

Theorem 4.3. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q^4; q^4)_n}.$$

(1) *For the continued fraction expansion of $z^2F(z)/G(z)$, we have*

$$a_{2k} = \frac{(-q^6; q^8)_k^2 (1+q^{8k})}{(-q^2; q^8)_k^2 q^{8k}}, \quad k \geq 1, \quad a_0 = 1,$$

$$a_{2k+1} = \frac{(-q^2; q^8)_{k+1}^2 (1+q^{8k+4})}{(-q^6; q^8)_k^2 q^{8k+2}}.$$

$$b_{2k} = \frac{(-q^6; q^8)_k^2 (1-q^{16k})}{(-q^2; q^8)_k^2 (1+q^{8k+2})(1+q^{8k-2})q^{4k+1}}, \quad k \geq 1, \quad b_0 = -\frac{q}{1+q^2},$$

$$b_{2k+1} = \frac{(-q^2; q^8)_{k+1}^2 (1-q^{16k+8})}{(-q^6; q^8)_k^2 (1+q^{8k+6})(1+q^{8k+2})q^{4k+1}}.$$

$$s_{2k} = q^{8k^2+2k} \sum_{n \geq 0} \frac{z^n q^{n^2+4nk} (-q^2; q^8)_k}{(q^2; q^2)_n (-q^2; q^2)_{n+4k} (-q^6; q^8)_k},$$

$$s_{2k+1} = q^{8k^2+10k+2} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)} (-q^6; q^8)_k}{(q^2; q^2)_n (-q^2; q^2)_{n+4k+2} (-q^2; q^8)_{k+1}}.$$

$$\frac{z^2F(z)}{G(z)} = \frac{z^2(1+q^2)}{1+q^2-zq} + \frac{z^2q^2(1+q^6)}{(1+q^2)(1+q^4)(1+q^6)+zq(1-q^8)}$$

$$\begin{aligned}
 & \frac{z^2 q^{10}(1+q^2)(1+q^{10})}{(1+q^6)(1+q^8)(1+q^{10}) + zq^3(1-q^{16})} + \frac{z^2 q^{18}(1+q^6)(1+q^{14})}{(1+q^{10})(1+q^{12})(1+q^{14}) + zq^5(1-q^{24})} \\
 & + \cdots + \frac{z^2 q^{8n+2}(1+q^{4n-2})(1+q^{4n+6})}{(1+q^{4n+2})(1+q^{4n+4})(1+q^{4n+6}) + zq^{2n+1}(1-q^{8n+8})} + \cdots. \quad (4.5)
 \end{aligned}$$

Setting $z = 1$ in (4.5), we have

$$\begin{aligned}
 \frac{(q^2, q^3; q^5)_\infty}{(q, q^4; q^5)_\infty} &= \frac{(1+q^2)}{1+q^2-q} \frac{q^2(1+q^6)}{(1+q^2)(1+q^4)(1+q^6) + q(1-q^8)} \\
 &+ \frac{q^{10}(1+q^2)(1+q^{10})}{(1+q^6)(1+q^8)(1+q^{10}) + q^3(1-q^{16})} + \frac{q^{18}(1+q^6)(1+q^{14})}{(1+q^{10})(1+q^{12})(1+q^{14}) + q^5(1-q^{24})} \\
 &+ \cdots + \frac{q^{8n+2}(1+q^{4n-2})(1+q^{4n+6})}{(1+q^{4n+2})(1+q^{4n+4})(1+q^{4n+6}) + q^{2n+1}(1-q^{8n+8})} + \cdots.
 \end{aligned}$$

(2) For the continued fraction expansion of $z^2 G(z)/F(z)$, we have

$$\begin{aligned}
 a_{2k} &= \frac{(-q^6; q^8)_k^2 (1+q^{8k})}{(-q^2; q^8)_k^2 q^{8k}}, \quad k \geq 1, \quad a_0 = 1, \\
 a_{2k+1} &= \frac{(-q^2; q^8)_{k+1}^2 (1+q^{8k+4})}{(-q^6; q^8)_k^2 q^{8k+6}}. \\
 b_{2k} &= \frac{(-q^6; q^8)_k^2 (1-q^{16k})}{(-q^2; q^8)_k^2 (1+q^{8k+2})(1+q^{8k-2})q^{4k-1}}, \quad k \geq 1, \quad b_0 = \frac{q}{1+q^2}, \\
 b_{2k+1} &= \frac{(-q^2; q^8)_{k+1}^2 (1-q^{16k+8})}{(-q^6; q^8)_k^2 (1+q^{8k+6})(1+q^{8k+2})q^{4k+3}}. \\
 s_{2k} &= q^{8k^2+6k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)} (-q^2; q^8)_k}{(q^2; q^2)_n (-q^2; q^2)_{n+4k} (-q^6; q^8)_k}, \\
 s_{2k+1} &= q^{8k^2+14k+6} \sum_{n \geq 0} \frac{z^n q^{n^2+4n(k+1)} (-q^6; q^8)_k}{(q^2; q^2)_n (-q^2; q^2)_{n+4k+2} (-q^2; q^8)_{k+1}}.
 \end{aligned}$$

$$\begin{aligned}
 \frac{z^2 G(z)}{F(z)} &= \frac{z^2(1+q^2)}{1+q^2+zq} \frac{z^2 q^6(1+q^6)}{(1+q^2)(1+q^4)(1+q^6) + zq^3(1-q^8)} \\
 &+ \frac{z^2 q^{14}(1+q^2)(1+q^{10})}{(1+q^6)(1+q^8)(1+q^{10}) + zq^5(1-q^{16})} + \frac{z^2 q^{22}(1+q^6)(1+q^{14})}{(1+q^{10})(1+q^{12})(1+q^{14}) + zq^7(1-q^{24})} \\
 &+ \cdots + \frac{z^2 q^{8n-2}(1+q^{4n-6})(1+q^{4n+2})}{(1+q^{4n-2})(1+q^{4n})(1+q^{4n+2}) + zq^{2n+1}(1-q^{8n})} + \cdots. \quad (4.6)
 \end{aligned}$$

Setting $z = 1$ in (4.6), we have

$$\begin{aligned}
 \frac{(q, q^4; q^5)_\infty}{(q^2, q^3; q^5)_\infty} &= \frac{(1+q^2)}{1+q^2+q} \frac{q^6(1+q^6)}{(1+q^2)(1+q^4)(1+q^6) + q^3(1-q^8)} \\
 &+ \frac{q^{14}(1+q^2)(1+q^{10})}{(1+q^6)(1+q^8)(1+q^{10}) + q^5(1-q^{16})} + \frac{q^{22}(1+q^6)(1+q^{14})}{(1+q^{10})(1+q^{12})(1+q^{14}) + q^7(1-q^{24})} \\
 &+ \cdots
 \end{aligned}$$

$$+ \dots + \frac{q^{8n-2}(1+q^{4n-6})(1+q^{4n+2})}{(1+q^{4n-2})(1+q^{4n})(1+q^{4n+2}) + q^{2n+1}(1-q^{8n})} + \dots$$

Theorem 4.4. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q^4; q^4)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+4n}}{(q^4; q^4)_n}.$$

(1) *For the continued fraction expansion of $z^2 F(z)/G(z)$, we have*

$$\begin{aligned} a_{2k} &= \frac{(-q^4; q^8)_k^2 (1+q^{8k-2})}{(-q^8; q^8)_{k-1}^2 q^{8k-2}}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= \frac{(-q^8; q^8)_k^2 (1+q^{8k+2})}{(-q^4; q^8)_k^2 q^{8k+2}}. \\ b_{2k} &= \frac{(-q^4; q^8)_{k-1} (-q^4; q^8)_k (1-q^{16k-4})}{(-q^8; q^8)_{k-1} (-q^8; q^8)_k q^{4k-1}}, \quad k \geq 1, \quad b_0 = -q, \\ b_{2k+1} &= \frac{(-q^8; q^8)_{k-1} (-q^8; q^8)_k (1-q^{16k+4})}{(-q^4; q^8)_k (-q^4; q^8)_{k+1} q^{4k+1}}, \quad k \geq 1, \quad b_1 = \frac{1}{(1+q^4)q}. \\ s_{2k} &= \frac{q^{8k^2} (-q^8; q^8)_{k-1}}{(-q^4; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+4nk}}{(q^2; q^2)_n (-q^2; q^2)_{n+4k-1}}, \quad k \geq 1, \\ s_{2k+1} &= \frac{q^{2(2k+1)^2} (-q^4; q^8)_k}{(-q^8; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)}}{(q^2; q^2)_n (-q^2; q^2)_{n+4k+1}}. \end{aligned}$$

$$\begin{aligned} \frac{z^2 F(z)}{G(z)} &= \frac{z^2}{1-zq} + \frac{z^2 q^3 (1+q^4)}{(1+q^2)(1+q^4)q + zq^2} + \frac{z^2 q^9 (1+q^4)(1+q^8)}{(1+q^4)^2 (1+q^6)(1+q^8) + zq^3 (1+q^4)(1-q^{12})} \\ &\quad + \frac{z^2 q^{16} (1+q^4)^2 (1+q^8)(1+q^{12})}{(1+q^8)^2 (1+q^{10})(1+q^{12}) + zq^5 (1+q^8)(1-q^{20})} \\ &\quad + \frac{z^2 q^{24} (1+q^8)^2 (1+q^{12})(1+q^{16})}{(1+q^{12})^2 (1+q^{14})(1+q^{16}) + zq^7 (1+q^{12})(1-q^{28})} \\ &\quad + \frac{z^2 q^{8n} (1+q^{4n-4})^2 (1+q^{4n})(1+q^{4n+4})}{(1+q^{4n})^2 (1+q^{4n+2})(1+q^{4n+4}) + zq^{2n+1} (1+q^{4n})(1-q^{8n+4})} + \dots \end{aligned}$$

(2) *For the continued fraction expansion of $z^2 G(z)/F(z)$, we have*

$$\begin{aligned} a_{2k} &= \frac{(-q^4; q^8)_k^2 (1+q^{8k-2})}{(-q^8; q^8)_{k-1}^2 q^{8k}}, \quad k \geq 1, \quad a_0 = 1, \\ a_{2k+1} &= \frac{(-q^8; q^8)_k^2 (1+q^{8k+2})}{(-q^4; q^8)_k^2 q^{8k+4}}. \\ b_{2k} &= \frac{(-q^4; q^8)_k^2 (1-q^{16k-4})}{(-q^8; q^8)_{k-1}^2 (1+q^{8k})(1+q^{8k-4})q^{4k-1}}, \quad k \geq 1, \quad b_0 = q, \\ b_{2k+1} &= \frac{(-q^8; q^8)_{k-1} (-q^8; q^8)_k (1-q^{16k+4})}{(-q^4; q^8)_{k+1} (-q^4; q^8)_k q^{4k+1}}, \quad k \geq 1, \quad b_1 = -\frac{q^3}{1+q^4}. \end{aligned}$$

$$s_{2k} = q^{8k^2+4k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)} (-q^8; q^8)_{k-1}}{(q^2; q^2)_n (-q^2; q^2)_{n+4k-1} (-q^4; q^8)_k}, \quad k \geq 1,$$

$$s_{2k+1} = q^{8k^2+12k+4} \sum_{n \geq 0} \frac{z^n q^{n^2+4n(k+1)} (-q^4; q^8)_k}{(q^2; q^2)_n (-q^2; q^2)_{n+4k+1} (-q^8; q^8)_k}.$$

$$\begin{aligned} \frac{z^2 G(z)}{F(z)} &= \frac{z^2}{1+zq} + \frac{z^2 q^4 (1+q^4)}{(1+q^2)(1+q^4) - zq^7} + \frac{z^2 q^{12} (1+q^8)}{(1+q^4)(1+q^6)(1+q^8) + zq^5(1-q^{12})} \\ &+ \frac{z^2 q^{20} (1+q^4)(1+q^{12})}{(1+q^8)(1+q^{10})(1+q^{12}) + zq^7(1-q^{20})} + \frac{z^2 q^{28} (1+q^8)(1+q^{16})}{(1+q^{12})(1+q^{14})(1+q^{16}) + zq^9(1-q^{28})} \\ &+ \cdots + \frac{z^2 q^{8n+4} (1+q^{4n-4})(1+q^{4n+4})}{(1+q^{4n})(1+q^{4n+2})(1+q^{4n+4}) + zq^{2n+3}(1-q^{8n+4})} + \cdots \end{aligned}$$

Theorem 4.5. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2} (-q; q^2)_n}{(q^2; q^2)_n}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n} (-q; q^2)_n}{(q^2; q^2)_n}.$$

For the continued fraction expansion of $z^2 G(z)/F(z)$, we have

$$a_{2k} = \frac{(-q; q^4)_k}{(-q^3; q^4)_k q^{6k}}, \quad a_{2k+1} = -\frac{(-q^3; q^4)_k}{(-q; q^4)_{k+1} q^{6k+5}},$$

$$b_{2k} = \frac{(-q; q^4)_k (1+q^{4k+1} + q^{4k-1})}{(-q^3; q^4)_k q^{2k-1}}, \quad k \geq 1, \quad b_0 = q(1+q),$$

$$b_{2k+1} = -\frac{(-q^3; q^4)_k (1+q^{4k+1} + q^{4k+3})}{(-q; q^4)_{k+1} q^{2k+2}}.$$

$$s_{2k} = \frac{(-1)^k q^{6k^2+5k}}{(-q; q^4)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)} (-q; q^2)_{n+2k}}{(q^2; q^2)_n},$$

$$s_{2k+1} = \frac{(-1)^{k-1} q^{6k^2+11k+5}}{(-q^3; q^4)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+4n(k+1)} (-q; q^2)_{n+2k+1}}{(q^2; q^2)_n}.$$

$$\begin{aligned} \frac{z^2 G(z)}{F(z)} &= \frac{z^2}{1+zq(1+q)} - \frac{z^2 q^5 (1+q)}{1+zq^3(1+q+q^3)} - \frac{z^2 q^{11} (1+q^3)}{1+zq^5(1+q^3+q^5)} \\ &- \frac{z^2 q^{17} (1+q^5)}{1+zq^7(1+q^5+q^7)} - \cdots - \frac{z^2 q^{6n-1} (1+q^{2n-1})}{1+zq^{2n+1}(1+q^{2n-1}+q^{2n+1})} - \cdots \end{aligned} \quad (4.7)$$

Setting $z = 1$ in (4.7), we have

$$\begin{aligned} \frac{(q, q^7; q^8)_\infty}{(q^3, q^5; q^8)_\infty} &= \frac{1}{1+q(1+q)} - \frac{q^5(1+q)}{1+q^3(1+q+q^3)} - \frac{q^{11}(1+q^3)}{1+q^5(1+q^3+q^5)} \\ &- \frac{q^{17}(1+q^5)}{1+q^7(1+q^5+q^7)} - \cdots - \frac{q^{6n-1}(1+q^{2n-1})}{1+q^{2n+1}(1+q^{2n-1}+q^{2n+1})} - \cdots \end{aligned}$$

Theorem 4.6. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n+1}}.$$

For the continued fraction expansion of $z^2 G(z)/F(z)$, we have

$$\begin{aligned} a_{2k} &= \frac{(q^7; q^8)_k^2 (1 - q^{8k+1})}{(q^3; q^8)_k^2 q^{8k}}, & a_{2k+1} &= -\frac{(q^3; q^8)_{k+1}^2 (1 - q^{8k+5})}{(q^7; q^8)_k^2 q^{8k+6}}. \\ b_{2k} &= \frac{(q^7; q^8)_k^2 (1 - q^{8k+1})(1 + q^2)}{(q^3; q^8)_k^2 (1 - q^{8k+3})(1 - q^{8k-1})}, \quad k \geq 1, & b_0 &= \frac{q^2}{1 - q^3}, \\ b_{2k+1} &= -\frac{(q^3; q^8)_{k+1}^2 (1 - q^{8k+5})(1 + q^2)}{(q^7; q^8)_k^2 (1 - q^{8k+3})(1 - q^{8k+7})q^2}. \\ s_{2k} &= \frac{(-1)^k q^{8k^2+6k} (q^3; q^8)_k}{(q^7; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{2n^2+2n(4k+1)}}{(q; q)_{2n} (q^{2n+1}; q^2)_{4k+1}}, \\ s_{2k+1} &= \frac{(-1)^{k-1} q^{8k^2+14k+6} (q^7; q^8)_k}{(q^3; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n q^{2n^2+2n(4k+3)}}{(q; q)_{2n} (q^{2n+1}; q^2)_{4k+3}}. \\ \frac{z^2 G(z)}{F(z)} &= \frac{z^2(1 - q^3)}{(1 - q)(1 - q^3) + zq^2} - \frac{z^2 q^6(1 - q^7)}{(1 - q^3)(1 - q^5)(1 - q^7) + zq^4(1 + q^2)(1 - q^5)} \\ &\quad - \frac{z^2 q^{14}(1 - q^3)(1 - q^{11})}{(1 - q^7)(1 - q^9)(1 - q^{11}) + zq^8(1 + q^2)(1 - q^9)} \\ &\quad - \frac{z^2 q^{22}(1 - q^7)(1 - q^{15})}{(1 - q^{11})(1 - q^{13})(1 - q^{15}) + zq^{12}(1 + q^2)(1 - q^{13})} \\ &\quad - \frac{z^2 q^{30}(1 - q^3)(1 - q^{19})}{(1 - q^{15})(1 - q^{17})(1 - q^{19}) + zq^{16}(1 + q^2)(1 - q^{17})} \\ &\quad - \dots - \frac{z^2 q^{8n-2}(1 - q^{4n-5})(1 - q^{4n+3})}{(1 - q^{4n-1})(1 - q^{4n+1})(1 - q^{4n+3}) + zq^{4n}(1 + q^2)(1 - q^{4n+1})} - \dots. \end{aligned} \quad (4.8)$$

Setting $z = 1$ in (4.8), we have

$$\begin{aligned} \frac{(q^3, q^5; q^8)_\infty (q^2, q^{14}; q^{16})_\infty}{(q, q^7; q^8)_\infty (q^6, q^{10}; q^{16})_\infty} &= \frac{(1 - q^3)}{(1 - q)(1 - q^3) + q^2} - \frac{q^6(1 - q^7)}{(1 - q^3)(1 - q^5)(1 - q^7) + q^4(1 + q^2)(1 - q^5)} \\ &\quad - \frac{q^{14}(1 - q^3)(1 - q^{11})}{(1 - q^7)(1 - q^9)(1 - q^{11}) + q^8(1 + q^2)(1 - q^9)} - \frac{q^{22}(1 - q^7)(1 - q^{15})}{(1 - q^{11})(1 - q^{13})(1 - q^{15}) + q^{12}(1 + q^2)(1 - q^{13})} \\ &\quad - \frac{q^{30}(1 - q^3)(1 - q^{19})}{(1 - q^{15})(1 - q^{17})(1 - q^{19}) + q^{16}(1 + q^2)(1 - q^{17})} - \dots. \end{aligned}$$

Theorem 4.7. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n}}.$$

For the continued fraction expansion of $z^2 G(z)/F(z)$, we have

$$\begin{aligned} a_{2k} &= \frac{(q^5; q^8)_k^2 (1 - q^{8k-1})}{(q; q^8)_k^2 q^{8k}}, \quad k \geq 1, & a_0 &= 1, \\ a_{2k+1} &= -\frac{(q; q^8)_{k+1}^2 (1 - q^{8k+3})}{(q^5; q^8)_k^2 q^{8k+6}}. \end{aligned}$$

$$\begin{aligned}
 b_{2k} &= \frac{(q^5; q^8)_k^2 (1 - q^{8k-1})(1 + q^2)}{(q; q^8)_k^2 (1 - q^{8k+1})(1 - q^{8k-3})}, \quad k \geq 1, \quad b_0 = \frac{q^2}{1 - q}, \\
 b_{2k+1} &= -\frac{(q; q^8)_{k+1}^2 (1 - q^{8k+3})(1 + q^2)}{(q^5; q^8)_k^2 (1 - q^{8k+1})(1 - q^{8k+5})q^2}, \\
 s_{2k} &= \frac{(-1)^k q^{8k^2+6k} (q; q^8)_k}{(q^5; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{2n^2+2n(4k+1)}}{(q; q)_{2n} (q^{2n+1}; q^2)_{4k}}, \\
 s_{2k+1} &= \frac{(-1)^{k-1} q^{8k^2+14k+6} (q^5; q^8)_k}{(q; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n q^{2n^2+2n(4k+3)}}{(q; q)_{2n} (q^{2n+1}; q^2)_{4k+2}}.
 \end{aligned}$$

$$\begin{aligned}
 \frac{z^2 G(z)}{F(z)} &= \frac{z^2(1-q)}{1-q+zq^2} - \frac{z^2 q^6(1-q^5)}{(1-q)(1-q^3)(1-q^5) + zq^4(1+q^2)(1-q^3)} \\
 &\quad - \frac{z^2 q^{14}(1-q)(1-q^9)}{(1-q^5)(1-q^7)(1-q^9) + zq^8(1+q^2)(1-q^7)} - \frac{z^2 q^{22}(1-q^5)(1-q^{13})}{(1-q^9)(1-q^{11})(1-q^{13}) + zq^{12}(1+q^2)(1-q^{11})} \\
 &\quad - \frac{z^2 q^{8n+6}(1-q^{4n-3})(1-q^{4n+5})}{(1-q^{4n+1})(1-q^{4n+3})(1-q^{4n+5}) + zq^{4n+4}(1+q^2)(1-q^{4n+3})} - \dots.
 \end{aligned}$$

Theorem 4.8. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n+1}}.$$

For the continued fraction expansion of $z^2 G(z)/F(z)$, we have

$$\begin{aligned}
 a_{2k} &= \frac{(q^7; q^8)_k^2 (1 - q^{8k+1})}{(q^3; q^8)_k^2 q^{8k}}, \quad a_{2k+1} = -\frac{(q^3; q^8)_{k+1}^2 (1 - q^{8k+5})}{(q^7; q^8)_k^2 q^{8k+7}}, \\
 b_{2k} &= \frac{(q^7; q^8)_k^2 (1 - q^{16k+2})}{(q^3; q^8)_k^2 (1 - q^{8k+3})(1 - q^{8k-1})q^{4k-1}}, \quad k \geq 1, \quad b_0 = \frac{q}{1 - q^3}, \\
 b_{2k+1} &= -\frac{(q^3; q^8)_k (q^3; q^8)_{k+1} (1 - q^{16k+10})}{(q^7; q^8)_k (q^7; q^8)_{k+1} q^{4k+4}}, \\
 s_{2k} &= \frac{(-1)^k q^{8k^2+7k} (q^3; q^8)_k}{(q^7; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)}}{(q^2; q^2)_n (q; q^2)_{n+4k+1}}, \\
 s_{2k+1} &= \frac{(-1)^{k-1} q^{8k^2+15k+7} (q^7; q^8)_k}{(q^3; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n q^{n^2+4n(k+1)}}{(q^2; q^2)_n (q; q^2)_{n+4k+3}}.
 \end{aligned}$$

$$\begin{aligned}
 \frac{z^2 G(z)}{F(z)} &= \frac{z^2(1-q^3)}{(1-q)(1-q^3) + zq} - \frac{z^2 q^7(1-q^7)}{(1-q^3)(1-q^5)(1-q^7) + zq^3(1-q^{10})} \\
 &\quad - \frac{z^2 q^{15}(1-q^3)(1-q^{11})}{(1-q^7)(1-q^9)(1-q^{11}) + zq^5(1-q^{18})} - \frac{z^2 q^{23}(1-q^7)(1-q^{15})}{(1-q^{11})(1-q^{13})(1-q^{15}) + zq^7(1-q^{26})} \\
 &\quad - \frac{z^2 q^{8n-1}(1-q^{4n-5})(1-q^{4n+3})}{(1-q^{4n-1})(1-q^{4n+1})(1-q^{4n+3}) + zq^{2n+1}(1-q^{8n+2})} - \dots. \quad (4.9)
 \end{aligned}$$

Setting $z = 1$ in (4.9), we have

$$\frac{(q^4, q^{16}; q^{20})_\infty}{(q^8, q^{12}; q^{20})_\infty} = \frac{(1-q^3)}{(1-q)(1-q^3) + q} - \frac{q^7(1-q^7)}{(1-q^3)(1-q^5)(1-q^7) + q^3(1-q^{10})}$$

$$\begin{aligned}
& \frac{q^{15}(1-q^3)(1-q^{11})}{(1-q^7)(1-q^9)(1-q^{11}) + q^5(1-q^{18})} - \frac{q^{23}(1-q^7)(1-q^{15})}{(1-q^{11})(1-q^{13})(1-q^{15}) + q^7(1-q^{26})} \\
& - \dots - \frac{q^{8n-1}(1-q^{4n-5})(1-q^{4n+3})}{(1-q^{4n-1})(1-q^{4n+1})(1-q^{4n+3}) + q^{2n+1}(1-q^{8n+2})} - \dots
\end{aligned}$$

Theorem 4.9. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+2n}}{(q; q)_{2n}}.$$

(1) *For the continued fraction expansion of $z^2 F(z)/G(z)$, we have*

$$\begin{aligned}
a_{2k} &= -\frac{(q^5; q^8)_k^2 (1 - q^{8k-1})}{(q; q^8)_k^2 q^{8k-1}}, \quad k \geq 1, \quad a_0 = 1, \\
a_{2k+1} &= \frac{(q; q^8)_{k+1}^2 (1 - q^{8k+3})}{(q^5; q^8)_k^2 q^{8k+2}}. \\
b_{2k} &= -\frac{(q^5; q^8)_k (q^5; q^8)_{k-1} (1 - q^{16k-2})}{(q; q^8)_k (q; q^8)_{k+1} q^{4k}}, \quad k \geq 1, \quad b_0 = -\frac{q}{1-q}, \\
b_{2k+1} &= \frac{(q; q^8)_{k+1}^2 (1 - q^{16k+6})}{(q^5; q^8)_k (q^5; q^8)_{k+1} (1 - q^{8k+1}) q^{4k+1}}. \\
s_{2k} &= \frac{(-1)^k q^{8k^2+k} (q; q^8)_k}{(q^5; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+4nk}}{(q^2; q^2)_n (q; q^2)_{n+4k}}, \\
s_{2k+1} &= \frac{(-1)^k q^{8k^2+9k+2} (q^5; q^8)_k}{(q; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)}}{(q^2; q^2)_n (q; q^2)_{n+4k+2}}.
\end{aligned}$$

$$\begin{aligned}
\frac{z^2 F(z)}{G(z)} &= \frac{z^2(1-q)}{1-q-zq} + \frac{z^2 q^2 (1-q^5)}{(1-q)(1-q^3)(1-q^5) + zq(1-q^6)} \\
& - \frac{z^2 q^9 (1-q)(1-q^9)}{(1-q^5)(1-q^7)(1-q^9) + zq^3(1-q^{14})} - \frac{z^2 q^{17} (1-q^5)(1-q^{13})}{(1-q^9)(1-q^{11})(1-q^{13}) + zq^5(1-q^{22})} \\
& - \dots - \frac{z^2 q^{8n+1} (1-q^{4n-3})(1-q^{4n+5})}{(1-q^{4n+1})(1-q^{4n+3})(1-q^{4n+5}) + zq^{2n+1}(1-q^{8n+6})} - \dots
\end{aligned}$$

(2) *For the continued fraction expansion of $z^2 G(z)/F(z)$, we have*

$$\begin{aligned}
a_{2k} &= \frac{(q^5; q^8)_k^2 (1 - q^{8k-1})}{(q; q^8)_k^2 q^{8k}}, \quad k \geq 1, \quad a_0 = 1, \\
a_{2k+1} &= -\frac{(q; q^8)_{k+1}^2 (1 - q^{8k+3})}{(q^5; q^8)_k^2 q^{8k+5}}. \\
b_{2k} &= \frac{(q^5; q^8)_k (q^5; q^8)_{k-1} (1 - q^{16k-2})}{(q; q^8)_k (q; q^8)_{k+1} q^{4k-1}}, \quad k \geq 1, \quad b_0 = \frac{q}{1-q}, \\
b_{2k+1} &= -\frac{(q; q^8)_k (q; q^8)_{k+1} (1 - q^{16k+6})}{(q^5; q^8)_k (q^5; q^8)_{k+1} q^{4k+2}}.
\end{aligned}$$

$$s_{2k} = \frac{(-1)^k q^{8k^2+5k} (q; q^8)_k}{(q^5; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)}}{(q^2; q^2)_n (q; q^2)_{n+4k}},$$

$$s_{2k+1} = \frac{(-1)^{k-1} q^{8k^2+13k+5} (q^5; q^8)_k}{(q; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n q^{n^2+4n(k+1)}}{(q^2; q^2)_n (q; q^2)_{n+4k+2}}.$$

$$\begin{aligned} \frac{z^2 G(z)}{F(z)} &= \frac{z^2(1-q)}{1-q+zz} - \frac{z^2 q^5(1-q^5)}{(1-q)(1-q^3)(1-q^5) + zq^3(1-q^6)} \\ &\quad - \frac{z^2 q^{13}(1-q)(1-q^9)}{(1-q^5)(1-q^7)(1-q^9) + zq^5(1-q^{14})} - \frac{z^2 q^{21}(1-q^5)(1-q^{13})}{(1-q^9)(1-q^{11})(1-q^{13}) + zq^7(1-q^{22})} \\ &\quad - \dots - \frac{z^2 q^{8n-3}(1-q^{4n-7})(1-q^{4n+1})}{(1-q^{4n-3})(1-q^{4n-1})(1-q^{4n+1}) + zq^{2n+1}(1-q^{8n-2})} - \dots \end{aligned}$$

Theorem 4.10. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n+1}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_{2n}}.$$

For the continued fraction expansion of $z^2 F(z)/G(z)$, we have

$$a_{2k} = \frac{(q^7; q^8)_k^2 (1-q^{8k+1})}{(q^3; q^8)_k^2 q^{8k}}, \quad a_{2k+1} = -\frac{(q^3; q^8)_{k+1}^2 (1-q^{8k+5})}{(q^7; q^8)_k^2 q^{8k+5}}.$$

$$b_{2k} = \frac{(q^7; q^8)_k (q^7; q^8)_{k-1} (1-q^{16k+2})}{(q^3; q^8)_{k+1} (q^3; q^8)_k q^{4k}}, \quad k \geq 1, \quad b_0 = \frac{q^3}{1-q^3},$$

$$b_{2k+1} = -\frac{(q^3; q^8)_{k+1} (q^3; q^8)_k (1-q^{16k+10})}{(q^7; q^8)_{k+1} (q^7; q^8)_k q^{4k+3}}.$$

$$s_{2k} = \frac{(-1)^k q^{8k^2+5k} (q^3; q^8)_k}{(q^7; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+n(4k+1)}}{(q^2; q^2)_n (q; q^2)_{n+4k+1}},$$

$$s_{2k+1} = \frac{(-1)^{k-1} q^{8k^2+13k+5} (q^7; q^8)_k}{(q^3; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n q^{n^2+n(4k+3)}}{(q^2; q^2)_n (q; q^2)_{n+4k+3}}.$$

$$\begin{aligned} \frac{z^2 F(z)}{G(z)} &= \frac{z^2(1-q^3)}{(1-q)(1-q^3) + zq^3} - \frac{z^2 q^5(1-q^7)}{(1-q^3)(1-q^5)(1-q^7) + zq^2(1-q^{10})} \\ &\quad - \frac{z^2 q^{13}(1-q^3)(1-q^{11})}{(1-q^7)(1-q^9)(1-q^{11}) + zq^4(1-q^{18})} - \frac{z^2 q^{21}(1-q^7)(1-q^{15})}{(1-q^{11})(1-q^{13})(1-q^{15}) + zq^6(1-q^{26})} \\ &\quad - \dots - \frac{z^2 q^{8n-3}(1-q^{4n-5})(1-q^{4n+3})}{(1-q^{4n-1})(1-q^{4n+1})(1-q^{4n+3}) + zq^{2n}(1-q^{8n+2})} - \dots \end{aligned} \quad (4.10)$$

Setting $z = 1$ in (4.10), we have

$$\begin{aligned} \frac{(q^3, q^7; q^{10})_\infty (q^4, q^{16}; q^{20})_\infty}{(q, q^9; q^{10})_\infty (q^8, q^{12}; q^{20})_\infty} &= \frac{(1-q^3)}{(1-q)(1-q^3) + q^3} - \frac{q^5(1-q^7)}{(1-q^3)(1-q^5)(1-q^7) + q^2(1-q^{10})} \\ &\quad - \frac{q^{13}(1-q^3)(1-q^{11})}{(1-q^7)(1-q^9)(1-q^{11}) + q^4(1-q^{18})} - \frac{q^{21}(1-q^7)(1-q^{15})}{(1-q^{11})(1-q^{13})(1-q^{15}) + q^6(1-q^{26})} \\ &\quad - \dots - \frac{q^{8n-3}(1-q^{4n-5})(1-q^{4n+3})}{(1-q^{4n-1})(1-q^{4n+1})(1-q^{4n+3}) + q^{2n}(1-q^{8n+2})} - \dots \end{aligned}$$

For the functions:

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{dn^2}}{(q; q)_{2n+1}}, \quad (4.11)$$

$$G(z) := \sum_{n \geq 0} \frac{z^n q^{dn^2}}{(q; q)_{2n}}, \quad (4.12)$$

the second author in [27] studied the continued fraction expansions of the q -tangent functions $zF(z)/G(z)$ for the cases $d = 0, 1, 2$. Here, we consider the expansions of $z^2F(z)/G(z)$ for the cases $d = 0$ and $d = 1$ in the following two theorems, respectively.

Theorem 4.11. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n+1}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n}}.$$

For the continued fraction expansion of $z^2F(z)/G(z)$, we have

$$a_{2k} = \frac{(q^7; q^8)_k^2 (1 - q^{8k+1})}{(q^3; q^8)_k^2 q^{8k}}, \quad a_{2k+1} = -\frac{(q^3; q^8)_{k+1}^2 (1 - q^{8k+5})}{(q^7; q^8)_k^2 q^{8k+4}}.$$

$$b_{2k} = \frac{(q^7; q^8)_k^2 (1 - q^{8k+1})(1 + q^2)}{(q^3; q^8)_k^2 (1 - q^{8k+3})(1 - q^{8k-1})q}, \quad k \geq 1, \quad b_0 = \frac{q}{1 - q^3},$$

$$b_{2k+1} = -\frac{(q^3; q^8)_{k+1}^2 (1 - q^{8k+5})(1 + q^2)}{(q^7; q^8)_k^2 (1 - q^{8k+7})(1 - q^{8k+3})q}.$$

$$s_{2k} = \frac{(-1)^k q^{8k^2+4k} (q^3; q^8)_k}{(q^7; q^8)_k} \sum_{n \geq 0} \frac{z^n}{(q^2; q^2)_n (q; q^2)_{n+4k+1}},$$

$$s_{2k+1} = \frac{(-1)^{k-1} q^{8k^2+12k+4} (q^7; q^8)_k}{(q^3; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n}{(q^2; q^2)_n (q; q^2)_{n+4k+3}}.$$

$$\begin{aligned} \frac{z^2F(z)}{G(z)} &= \frac{z^2(1 - q^3)}{(1 - q)(1 - q^3) + zq} - \frac{z^2q^4(1 - q^7)}{(1 - q^3)(1 - q^5)(1 - q^7) + zq^3(1 + q^2)(1 - q^5)} \\ &\quad - \frac{z^2q^{12}(1 - q^3)(1 - q^{11})}{(1 - q^7)(1 - q^9)(1 - q^{11}) + zq^7(1 + q^2)(1 - q^9)} \\ &\quad - \frac{z^2q^{20}(1 - q^7)(1 - q^{15})}{(1 - q^{11})(1 - q^{13})(1 - q^{15}) + zq^{11}(1 + q^2)(1 - q^{13})} \\ &\quad - \frac{z^2q^{28}(1 - q^3)(1 - q^{19})}{(1 - q^{15})(1 - q^{17})(1 - q^{19}) + zq^{15}(1 + q^2)(1 - q^{17})} \\ &\quad - \dots - \frac{z^2q^{8n-4}(1 - q^{4n-5})(1 - q^{4n+3})}{(1 - q^{4n-1})(1 - q^{4n+1})(1 - q^{4n+3}) + zq^{4n-1}(1 + q^2)(1 - q^{4n+1})} - \dots \end{aligned}$$

Theorem 4.12. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n+1}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_{2n}}.$$

For the continued fraction expansion of $z^2F(z)/G(z)$, we have

$$a_{2k} = \frac{(q^7; q^8)_k^2 (1 - q^{8k+1})}{(q^3; q^8)_k^2 q^{8k}}, \quad a_{2k+1} = -\frac{(q^3; q^8)_{k+1}^2 (1 - q^{8k+5})}{(q^7; q^8)_k^2 q^{8k+3}}.$$

$$\begin{aligned}
 b_{2k} &= \frac{(q^7; q^8)_k (q^7; q^8)_{k-1} (1 - q^{16k+2})}{(q^3; q^8)_{k+1} (q^3; q^8)_k q^{4k+1}}, \quad k \geq 1, \quad b_0 = \frac{q^2}{1 - q^3}, \\
 b_{2k+1} &= -\frac{(q^3; q^8)_{k+1} (q^3; q^8)_k (1 - q^{16k+10})}{(q^7; q^8)_{k+1} (q^7; q^8)_k q^{4k+2}}. \\
 s_{2k} &= \frac{(-1)^k q^{8k^2+3k} (q^3; q^8)_k}{(q^7; q^8)_k} \sum_{n \geq 0} \frac{z^n q^{n^2+4nk}}{(q^2; q^2)_n (q; q^2)_{n+4k+1}}, \\
 s_{2k+1} &= \frac{(-1)^{k-1} q^{8k^2+11k+3} (q^7; q^8)_k}{(q^3; q^8)_{k+1}} \sum_{n \geq 0} \frac{z^n q^{n^2+2n(2k+1)}}{(q^2; q^2)_n (q; q^2)_{n+4k+3}}.
 \end{aligned}$$

$$\begin{aligned}
 \frac{z^2 F(z)}{G(z)} &= \frac{z^2(1 - q^3)}{(1 - q)(1 - q^3) + zq^2} - \frac{z^2 q^3(1 - q^7)}{(1 - q^3)(1 - q^5)(1 - q^7) + zq(1 - q^{10})} \\
 &\quad - \frac{z^2 q^{11}(1 - q^3)(1 - q^{11})}{(1 - q^7)(1 - q^9)(1 - q^{11}) + zq^3(1 - q^{18})} - \frac{z^2 q^{19}(1 - q^7)(1 - q^{15})}{(1 - q^{11})(1 - q^{13})(1 - q^{15}) + zq^5(1 - q^{26})} \\
 &\quad - \dots - \frac{z^2 q^{8n-5}(1 - q^{4n-5})(1 - q^{4n+3})}{(1 - q^{4n-1})(1 - q^{4n+1})(1 - q^{4n+3}) + zq^{2n-1}(1 - q^{8n+2})} - \dots
 \end{aligned}$$

We notice that the above theorem is the same as Theorem 4.10 when replacing $z \rightarrow z/q$ there.

For the continued fraction expansion of $z^2 F(z)/G(z)$ for the functions (4.11) and (4.12) in the case $d = 2$, we cannot find good expressions for b_k .

Finally, we point out that according to the following formula [19]

$$\frac{1}{1 - \frac{c_1 z}{1 - \frac{c_2 z}{\ddots}}} = \frac{1}{1 - c_1 z - \frac{c_1 c_2 z^2}{1 - (c_2 + c_3)z - \frac{c_3 c_4 z^2}{\ddots}}}, \quad (4.13)$$

we can make connections between Section 2 and Section 4. By considering the following formula which is a variant of (4.13):

$$\frac{1}{a_0 + \frac{z}{a_1 + \frac{z}{a_2 + \frac{z}{\ddots}}}} = \frac{1}{A_0 + B_0 z + \frac{z^2}{A_1 + B_1 z + \frac{z^2}{A_2 + B_2 z + \frac{z^2}{\ddots}}}}, \quad (4.14)$$

where

$$\begin{aligned}
 A_{2k} &= a_0 \prod_{i=1}^k \frac{a_{4i-1}^2 a_{4i}}{a_{4i-4} a_{4i-3}^2}, \quad k \geq 1, \quad A_0 = a_0, & A_{2k+1} &= -\frac{1}{a_0} \prod_{i=0}^k \frac{a_{4i+1}^2 a_{4i+2}}{a_{4i-2} a_{4i-1}^2}, \\
 B_{2k} &= A_{2k} \frac{a_{4k-1} + a_{4k+1}}{a_{4k-1} a_{4k} a_{4k+1}}, \quad k \geq 1, \quad B_0 = \frac{1}{a_1}, & B_{2k+1} &= A_{2k+1} \frac{a_{4k+1} + a_{4k+3}}{a_{4k+1} a_{4k+2} a_{4k+3}},
 \end{aligned}$$

and $a_{-2} = 1/a_0$, $a_{-1} = 1$, we can get the continued fraction expansions of $z^2F(z)/G(z)$ (resp. $z^2G(z)/F(z)$) in Section 4 from those of $zF(z)/G(z)$ (resp. $zG(z)/F(z)$) in Section 2 as a corollary. However, we found a direct approach more attractive.

In what follows, according to the formula (4.14), we list some continued fraction expansions which do not have good expressions for s_k .

Theorem 4.13. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2+2n}}{(q; q)_{2n+1}}.$$

For the continued fraction expansion of $z^2F(z)/G(z)$, we have

$$\begin{aligned} a_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1 - q^{8k+1})(1 - q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_k^2 (q^{4k-1}; q)_4 q^{8k+2}}, \quad k \geq 1, \quad a_0 = \frac{1}{1 - q}, \\ a_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1}^2 (1 - q^{8k+5})}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1 - q^2)^2 (q^{4k+1}; q)_4 q^{8k+4}}. \\ b_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k (q^7; q^8)_{k-1} (1 - q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_{k+1} (q^3; q^8)_k (q^{4k-1}; q)_4 q^4} \\ &\quad \times [q^6(1 - q^{8k-1})(1 - q^{4k-1})^2(1 - q^{4k})^2 + (1 - q^{8k+3})(1 - q^{4k+1})^2(1 - q^{4k+2})^2], \\ &\quad k \geq 1, \quad b_0 = -\frac{q^2}{(1 - q)^2(1 - q^3)}, \\ b_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1} (q^3; q^8)_k}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_{k+1} (q^7; q^8)_k (1 - q^2)^2 (q^{4k+1}; q)_4 q^2} \\ &\quad \times [q^6(1 - q^{8k+3})(1 - q^{4k+1})^2(1 - q^{4k+2})^2 + (1 - q^{8k+7})(1 - q^{4k+3})^2(1 - q^{4k+4})^2]. \\ \frac{z^2F(z)}{G(z)} &= \frac{z^2(1 - q)^3(1 - q^3)}{(1 - q)^2(1 - q^3) - zq^2(1 - q)} \\ &\quad + \frac{z^2q^4(1 - q)(1 - q^3)^2(1 - q^4)^2(1 - q^7)}{(q; q)_4(1 - q^3)(1 - q^5)(1 - q^7) + zq^2[q^6(1 - q)^2(1 - q^2)^2(1 - q^3) + (1 - q^3)^2(1 - q^4)^2(1 - q^7)]} \\ &\quad - \frac{z^2q^{14}(1 - q^3)(1 - q^{11})(1 - q^2)^2(1 - q^2)^2(1 - q^5)^2(1 - q^6)^2}{(q^3; q)_4(1 - q^7)(1 - q^9)(1 - q^{11}) + zq^6[q^6(1 - q^3)^2(1 - q^4)^2(1 - q^7) + (1 - q^5)^2(1 - q^6)^2(1 - q^{11})]} \\ &\quad - \frac{z^2q^{22}(1 - q^7)(1 - q^{15})(1 - q^3)^2(1 - q^4)^2(1 - q^7)^2(1 - q^8)^2}{(q^5; q)_4(1 - q^{11})(1 - q^{13})(1 - q^{15}) + zq^{10}[q^6(1 - q^5)^2(1 - q^6)^2(1 - q^{11}) + (1 - q^7)^2(1 - q^8)^2(1 - q^{15})]} \\ &\quad - \dots - \frac{z^2q^{8n-2}(1 - q^{4n-5})(1 - q^{4n+3})(1 - q^{2n-3})^2(1 - q^{2n-2})^2(1 - q^{2n+1})^2(1 - q^{2n+2})^2}{(q^{2n-1}; q)_4(q^{4n-1}; q^2)_3 + zq^{4n-2}[q^6(q^{2n-1}; q)_2^2(1 - q^{4n-1}) + (q^{2n+1}; q)_2^2(1 - q^{4n+3})]} - \dots \end{aligned} \quad (4.15)$$

Setting $z = 1$ in (4.15), we have

$$\begin{aligned} \frac{(q, q^7; q^8)_\infty (q^6, q^{10}; q^{16})_\infty}{(q^3, q^5; q^8)_\infty (q^2, q^{14}; q^{16})_\infty} &= \frac{(1 - q)^3(1 - q^3)}{(1 - q)^2(1 - q^3) - q^2(1 - q)} \\ &\quad + \frac{q^4(1 - q)(1 - q^3)^2(1 - q^4)^2(1 - q^7)}{(q; q)_4(1 - q^3)(1 - q^5)(1 - q^7) + q^2[q^6(1 - q)^2(1 - q^2)^2(1 - q^3) + (1 - q^3)^2(1 - q^4)^2(1 - q^7)]} \\ &\quad - \frac{q^{14}(1 - q^3)(1 - q^{11})(1 - q^2)^2(1 - q^2)^2(1 - q^5)^2(1 - q^6)^2}{(q^3; q)_4(1 - q^7)(1 - q^9)(1 - q^{11}) + q^6[q^6(1 - q^3)^2(1 - q^4)^2(1 - q^7) + (1 - q^5)^2(1 - q^6)^2(1 - q^{11})]} \\ &\quad - \frac{q^{22}(1 - q^7)(1 - q^{15})(1 - q^3)^2(1 - q^4)^2(1 - q^7)^2(1 - q^8)^2}{(q^5; q)_4(1 - q^{11})(1 - q^{13})(1 - q^{15}) + q^{10}[q^6(1 - q^5)^2(1 - q^6)^2(1 - q^{11}) + (1 - q^7)^2(1 - q^8)^2(1 - q^{15})]} \end{aligned}$$

$$- \dots - \frac{q^{8n-2}(1-q^{4n-5})(1-q^{4n+3})(1-q^{2n-3})^2(1-q^{2n-2})^2(1-q^{2n+1})^2(1-q^{2n+2})^2}{(q^{2n-1}; q)_4(q^{4n-1}; q^2)_3 + q^{4n-2}[q^6(q^{2n-1}; q)_2^2(1-q^{4n-1}) + (q^{2n+1}; q)_2^2(1-q^{4n+3})]} - \dots$$

The continued fraction expansion of $zF(z)/G(z)$ for the functions $F(z)$ and $G(z)$ in the above theorem is given in Theorem 2.5.

Theorem 4.14. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n+1}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n}}.$$

For the continued fraction expansion of $z^2G(z)/F(z)$, we have

$$\begin{aligned} a_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1-q^{8k+1})(1-q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_k^2 (q^{4k-1}; q)_4 q^{8k+2}}, \quad k \geq 1, \quad a_0 = \frac{1}{1-q}, \\ a_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1}^2 (1-q^{8k+5})}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1-q^2)^2 (q^{4k+1}; q)_4 q^{8k+2}}. \\ b_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k (q^7; q^8)_{k-1} (1-q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_{k+1} (q^3; q^8)_k (q^{4k-1}; q)_4 q^5} \\ &\quad \times [q^6(1-q^{8k-1})(1-q^{4k-1})^2(1-q^{4k})^2 + (1-q^{8k+3})(1-q^{4k+1})^2(1-q^{4k+2})^2], \\ &\quad k \geq 1, \quad b_0 = -\frac{q}{(1-q)^2(1-q^3)}, \\ b_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1} (q^3; q^8)_k}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_{k+1} (q^7; q^8)_k (1-q^2)^2 (q^{4k+1}; q)_4 q} \\ &\quad \times [q^6(1-q^{8k+3})(1-q^{4k+1})^2(1-q^{4k+2})^2 + (1-q^{8k+7})(1-q^{4k+3})^2(1-q^{4k+4})^2]. \\ \frac{z^2G(z)}{F(z)} &= \frac{z^2(1-q)^3(1-q^3)}{(1-q)^2(1-q^3) - zq(1-q)} \\ &\quad + \frac{z^2q^2(1-q)(1-q^3)^2(1-q^4)^2(1-q^7)}{(q; q)_4(1-q^3)(1-q^5)(1-q^7) + zq[q^6(1-q)^2(1-q^2)^2(1-q^3) + (1-q^3)^2(1-q^4)^2(1-q^7)]} \\ &\quad - \frac{z^2q^{12}(1-q^3)(1-q^{11})(1-q^2)^2(1-q^2)^2(1-q^5)^2(1-q^6)^2}{(q^3; q)_4(1-q^7)(1-q^9)(1-q^{11}) + zq^5[q^6(1-q^3)^2(1-q^4)^2(1-q^7) + (1-q^5)^2(1-q^6)^2(1-q^{11})]} \\ &\quad - \frac{z^2q^{20}(1-q^7)(1-q^{15})(1-q^3)^2(1-q^4)^2(1-q^7)^2(1-q^8)^2}{(q^5; q)_4(1-q^{11})(1-q^{13})(1-q^{15}) + zq^9[q^6(1-q^5)^2(1-q^6)^2(1-q^{11}) + (1-q^7)^2(1-q^8)^2(1-q^{15})]} \\ &\quad - \dots - \frac{z^2q^{8n-4}(1-q^{4n-5})(1-q^{4n+3})(1-q^{2n-3})^2(1-q^{2n-2})^2(1-q^{2n+1})^2(1-q^{2n+2})^2}{(q^{2n-1}; q)_4(q^{4n-1}; q^2)_3 + zq^{4n-3}[q^6(q^{2n-1}; q)_2^2(1-q^{4n-1}) + (q^{2n+1}; q)_2^2(1-q^{4n+3})]} - \dots \end{aligned}$$

Theorem 4.15. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n+1}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n q^{2n^2}}{(q; q)_{2n}}.$$

For the continued fraction expansion of $z^2G(z)/F(z)$, we have

$$\begin{aligned} a_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1-q^{8k+1})(1-q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_k^2 (q^{4k-1}; q)_4 q^{8k+1}}, \quad k \geq 1, \quad a_0 = \frac{1}{1-q}, \\ a_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1}^2 (1-q^{8k+5})}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1-q^2)^2 (q^{4k+1}; q)_4 q^{8k+5}}. \end{aligned}$$

$$\begin{aligned}
b_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k (q^7; q^8)_{k-1} (1-q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_{k+1} (q^3; q^8)_k (q^{4k-1}; q)_4^2 q^3} \\
&\quad \times [q^6(1-q^{8k-1})(1-q^{4k-1})^2(1-q^{4k})^2 + (1-q^{8k+3})(1-q^{4k+1})^2(1-q^{4k+2})^2], \\
&\quad k \geq 1, \quad b_0 = -\frac{q^3}{(1-q)^2(1-q^3)}, \\
b_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1} (q^3; q^8)_k}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_{k+1} (q^7; q^8)_k (1-q^2)^2 (q^{4k+1}; q)_4^2 q^3} \\
&\quad \times [q^6(1-q^{8k+3})(1-q^{4k+1})^2(1-q^{4k+2})^2 + (1-q^{8k+7})(1-q^{4k+3})^2(1-q^{4k+4})^2]. \\
\frac{z^2 G(z)}{F(z)} &= \frac{z^2(1-q)^3(1-q^3)}{(1-q)^2(1-q^3) - zq^3(1-q)} \\
&\quad + \frac{z^2 q^5(1-q)(1-q^3)^2(1-q^4)^2(1-q^7)}{(q; q)_4(1-q^3)(1-q^5)(1-q^7) + zq^2[q^6(1-q)^2(1-q^2)^2(1-q^3) + (1-q^3)^2(1-q^4)^2(1-q^7)]} \\
&\quad - \frac{z^2 q^{14}(1-q^3)(1-q^{11})(1-q)^2(1-q^2)^2(1-q^5)^2(1-q^6)^2}{(q^3; q)_4(1-q^7)(1-q^9)(1-q^{11}) + zq^6[q^6(1-q^3)^2(1-q^4)^2(1-q^7) + (1-q^5)^2(1-q^6)^2(1-q^{11})]} \\
&\quad - \frac{z^2 q^{22}(1-q^7)(1-q^{15})(1-q^3)^2(1-q^4)^2(1-q^7)^2(1-q^8)^2}{(q^5; q)_4(1-q^{11})(1-q^{13})(1-q^{15}) + zq^{10}[q^6(1-q^5)^2(1-q^6)^2(1-q^{11}) + (1-q^7)^2(1-q^8)^2(1-q^{15})]} \\
&\quad - \dots - \frac{z^2 q^{8n-2}(1-q^{4n-5})(1-q^{4n+3})(1-q^{2n-3})^2(1-q^{2n-2})^2(1-q^{2n+1})^2(1-q^{2n+2})^2}{(q^{2n-1}; q)_4(q^{4n-1}; q^2)_3 + zq^{4n-2}[q^6(q^{2n-1}; q)_2^2(1-q^{4n-1}) + (q^{2n+1}; q)_2^2(1-q^{4n+3})]} - \dots
\end{aligned}$$

Theorem 4.16. *Let*

$$F(z) := \sum_{n \geq 0} \frac{z^n q^{2n}}{(q; q)_{2n+1}}, \quad G(z) := \sum_{n \geq 0} \frac{z^n}{(q; q)_{2n}}.$$

For the continued fraction expansion of $z^2 G(z)/F(z)$, we have

$$\begin{aligned}
a_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1-q^{8k+1})(1-q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_k^2 (q^{4k-1}; q)_4 q^{8k+3}}, \quad k \geq 1, \quad a_0 = \frac{1}{1-q}, \\
a_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1}^2 (1-q^{8k+5})}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k^2 (1-q^2)^2 (q^{4k+1}; q)_4 q^{8k+1}}. \\
b_{2k} &= -\frac{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_k (q^7; q^8)_{k-1} (1-q^2)^2}{(q; q^4)_k^4 (q^2; q^4)_k^4 (q^3; q^8)_{k+1} (q^3; q^8)_k (q^{4k-1}; q)_4^2 q^6} \\
&\quad \times [q^6(1-q^{8k-1})(1-q^{4k-1})^2(1-q^{4k})^2 + (1-q^{8k+3})(1-q^{4k+1})^2(1-q^{4k+2})^2], \\
&\quad k \geq 1, \quad b_0 = -\frac{1}{(1-q)^2(1-q^3)}, \\
b_{2k+1} &= \frac{(q; q^4)_{k+1}^4 (q^2; q^4)_{k+1}^4 (q^3; q^8)_{k+1} (q^3; q^8)_k}{(q^3; q^4)_k^4 (q^4; q^4)_k^4 (q^7; q^8)_{k+1} (q^7; q^8)_k (1-q^2)^2 (q^{4k+1}; q)_4^2} \\
&\quad \times [q^6(1-q^{8k+3})(1-q^{4k+1})^2(1-q^{4k+2})^2 + (1-q^{8k+7})(1-q^{4k+3})^2(1-q^{4k+4})^2]. \\
\frac{z^2 G(z)}{F(z)} &= \frac{z^2(1-q)^3(1-q^3)}{(1-q)^2(1-q^3) - z(1-q)} \\
&\quad + \frac{z^2 q(1-q)(1-q^3)^2(1-q^4)^2(1-q^7)}{(q; q)_4(1-q^3)(1-q^5)(1-q^7) + zq[q^6(1-q)^2(1-q^2)^2(1-q^3) + (1-q^3)^2(1-q^4)^2(1-q^7)]} \\
&\quad - \frac{z^2 q^{12}(1-q^3)(1-q^{11})(1-q)^2(1-q^2)^2(1-q^5)^2(1-q^6)^2}{(q^3; q)_4(1-q^7)(1-q^9)(1-q^{11}) + zq^5[q^6(1-q^3)^2(1-q^4)^2(1-q^7) + (1-q^5)^2(1-q^6)^2(1-q^{11})]}
\end{aligned}$$

$$\begin{aligned}
& \frac{z^2 q^{20} (1-q^7)(1-q^{15})(1-q^3)^2(1-q^4)^2(1-q^7)^2(1-q^8)^2}{(q^5; q)_4(1-q^{11})(1-q^{13})(1-q^{15}) + zq^9[q^6(1-q^5)^2(1-q^6)^2(1-q^{11}) + (1-q^7)^2(1-q^8)^2(1-q^{15})]} \\
& - \dots - \frac{z^2 q^{8n-4} (1-q^{4n-5})(1-q^{4n+3})(1-q^{2n-3})^2(1-q^{2n-2})^2(1-q^{2n+1})^2(1-q^{2n+2})^2}{(q^{2n-1}; q)_4(q^{4n-1}; q^2)_3 + zq^{4n-3}[q^6(q^{2n-1}; q)_2^2(1-q^{4n-1}) + (q^{2n+1}; q)_2^2(1-q^{4n+3})]} - \dots
\end{aligned}$$

The continued fraction expansions of the q -cotangent functions $zG(z)/F(z)$ for the functions $F(z)$ and $G(z)$ in Theorem 4.14, Theorem 4.15, and Theorem 4.16 were given in [27].

Acknowledgments. We are very grateful to the referees for their valuable comments and suggestions.

REFERENCES

- [1] G. E. Andrews, On q -difference equations for certain well-poised basic hypergeometric series, *Quart. J. Math. (Oxford)*, **19** (1968), 433–447.
- [2] G. E. Andrews, An introduction to Ramanujan’s “lost” notebook, *Amer. Math. Monthly*, **86** (1979), 89–108.
- [3] G. E. Andrews, Ramanujan’s “lost” notebook III. The Rogers-Ramanujan continued fraction, *Adv. Math.*, **41** (1981), 186–208.
- [4] G. E. Andrews, B. C. Berndt, L. Jacobsen, R. L. Lamphere, *The Continued Fractions Found in the Unorganized Portions of Ramanujan’s Notebooks*, *Memoirs Amer. Math. Soc.*, **477**, Amer. Math. Soc., Providence, 1992.
- [5] G. E. Andrews, B. C. Berndt, *Ramanujan’s Lost Notebook, Part I*, Springer-Verlag, New York, 2005.
- [6] B. C. Berndt, *Ramanujan’s Notebooks, Part V*, Springer-Verlag, New York, 1998.
- [7] S. Bhargava, C. Adiga, On some continued fraction identities of Srinivasa Ramanujan, *Proc. Amer. Math. Soc.*, **92** (1984), 13–18.
- [8] D. Bowman, J. Mc Laughlin, N. J. Wyshinski, A q -continued fraction, *Int. J. Number Theory*, **2** (2006), 523–547.
- [9] D. Bowman, J. Mc Laughlin, N. J. Wyshinski, Continued fraction proofs of m -versions of some identities of Rogers-Ramanujan-Slater type, *Ramanujan J.*, to appear.
- [10] H. H. Chan, S. -S. Huang, On the Ramanujan-Gordon-Göllnitz continued fraction, *Ramanujan J.*, **1** (1997), 75–90.
- [11] M. Fulmek, A continued fraction expansion for a q -tangent function, *Sém. Lothar. Combin.*, **45** (2000), [B45b].
- [12] G. Gasper, M. Rahman, *Basic Hypergeometric Series, Second Ed.*, Cambridge University Press, Cambridge, 2004.
- [13] B. Gordon, Some continued fractions of the Rogers-Ramanujan type, *Duke Math. J.*, **32** (1965), 741–748.
- [14] M. D. Hirschhorn, A continued fraction, *Duke Math. J.*, **41** (1974), 27–33.
- [15] M. D. Hirschhorn, *Developments in Theory of Partitions* PhD Thesis, University of New South Wales, 1980.
- [16] M. D. Hirschhorn, A continued fraction of Ramanujan, *J. Aust. Math. Soc. Ser. A*, **29** (1980), 80–86.
- [17] F. H. Jackson, A basic-sine and cosine with symbolic solutions of certain differential equations, *Proc. Edinb. Math. Soc.*, **22** (1904), 28–39.
- [18] F. H. Jackson, Examples of a generalization of Euler’s transformation for power series, *Messenger of Mathematics*, **57** (1928), 169–187.
- [19] W. B. Jones, W. J. Thron, *Continued Fractions. Analytic Theory and Applications*, Addison-Wesley, Reading, Mass., 1980.

- [20] A. N. Khovanskii, The Application of Continued Fractions and Their Generalizations to Problems in Approximation Theory, P. Noordhoff, Groningen, 1963.
- [21] V. A. Lebesgue, Sommatation de quelques séries, J. Math. Pures Appl., **5** (1840), 42–71.
- [22] T. Matala-aho, On the values of continued fractions: q -series. J. Approx. Theory, **124** (2003), 139–153.
- [23] T. Matala-aho, V. Meril, On the values of continued fractions: q -series. II, Int. J. Number Theory, **2** (2006), 417–430.
- [24] J. Mc Laughlin, A. Sills, P. Zimmer, Rogers-Ramanujan computer searches, J. Symbolic Comput., **44** (2009), 1068–1078.
- [25] H. Prodinger, Combinatorics of geometrically distributed random variables: New q -tangent and q -secant numbers, Int. J. Math. Math. Sci., **24** (2000), 825–838.
- [26] H. Prodinger, A continued fraction expansion for a q -tangent function: An elementary proof, Sémin. Lothar. Combin., **60** (2008), [B60b].
- [27] H. Prodinger, Continued fraction expansions for q -tangent and q -cotangent functions, Discrete Math. Theor. Comput. Sci., **12** (2010), 47–64.
- [28] K. G. Ramanathan, Ramanujan’s continued fraction, Indian J. Pure Appl. Math., **16** (1985), 695–724.
- [29] S. Ramanujan, The Lost Notebook and Other Unpublished Papers, Narosa, New Delhi, 1988.
- [30] L. J. Rogers, Second memoir on the expansion of certain infinite products, Proc. London Math. Soc., **25** (1894), 318–343.
- [31] A. Selberg, Über einige arithmetische Identitäten Avh. Norske Vid.CAkad. Oslo I. Mat.CNaturv. Kl. No. **8** (1936), 3–23.
- [32] A. Selberg, Collected Papers, Vol. I, Springer-Verlag, Berlin, 1989.
- [33] H. Shin, J. Zeng, The q -tangent and q -secant numbers via continued fractions, European J. Combin., **31** (2010), 1689–1705.
- [34] A. Sills, Finite Rogers-Ramanujan type identities, Electron. J. Combin., **10** (2003), R13.
- [35] L. J. Slater, Further identities of the Rogers-Ramanujan type, Proc. London Math. Soc., **54** (1952), 147–167.

(N. S. S. Gu) CENTER FOR COMBINATORICS, LPMC-TJKLC, NANKAI UNIVERSITY, TIANJIN 300071, P. R. CHINA

E-mail address: gu@nankai.edu.cn

(H. Prodinger) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH, 7602 STELLENBOSCH, SOUTH AFRICA

E-mail address: hproding@sun.ac.za