

Complete solution for the rainbow numbers of matchings *

He Chen, Xueliang Li and Jianhua Tu

Center for Combinatorics and LPMC-TJKLC, Nankai University

Tianjin 300071, P.R. China. Email: lxl@nankai.edu.cn

Abstract

For a given graph H and a positive n , the *rainbow number of H* , denoted by $rb(n, H)$, is the minimum integer k so that in any edge-coloring of K_n with k colors there is a copy of H whose edges have distinct colors. In 2004, Schiermeyer determined $rb(n, kK_2)$ for all $n \geq 3k + 3$. The case for smaller values of n (namely, $n \in [2k, 3k + 2]$) remained generally open. In this paper we extend Schiermeyer's result to all plausible n and hence determine the rainbow number of matchings.

Keywords: edge-colored graph, rainbow subgraph, rainbow number.

AMS subject classification 2000: 05C35, 05C55, 05C70.

1 Introduction

In this paper we consider undirected, finite and simple graphs only, and use standard notations in graph theory (see [3] and [8]). Let K_n be an edge-colored complete graph on n vertices. If a subgraph H of K_n contains no two edges of the same color, then H is called a *totally multicolored (TMC)* or *rainbow subgraph* of K_n and we say that K_n contains a TMC or rainbow H . Let $f(n, H)$ denote the maximum number of colors in an edge-coloring of K_n with no TMC H . We now define $rb(n, H)$ as the minimum number of colors such that any edge-coloring of K_n with at least $rb(n, H) = f(n, H) + 1$ colors contains a TMC or rainbow subgraph isomorphic to H . The number $rb(n, H)$ is called the *rainbow number of H* .

$f(n, H)$ is called the anti-Ramsey number of H , which was introduced by Erdős, Simonovits and Sós in the 1970s. They showed that it is closely related to the Turán number. Anti-Ramsey number has been studied in [1, 2, 5, 9, 11, 6, 7] and elsewhere. There are very few graphs whose anti-Ramsey numbers have been determined exactly.

*Supported by NSFC, PCSIRT and the "973" program.

To the best of our knowledge, $f(n, H)$ is known exactly for large n only when H is a complete graph, a path, a star, a cycle or a broom whose maximum degree exceeds its diameter (a broom is obtained by identifying an end of a path with a vertex of a star) (see [10, 9, 11, 6, 7]).

For a given graph H , let $ext(n, H)$ denote the maximum number of edges that a graph G of order n can have with no subgraph isomorphic to H . For $H = kK_2$, the value $ext(n, kK_2)$ has been determined by Erdős and Gallai [4], where $H = kK_2$ is a matching M of size k .

Theorem 1.1 (Erdős and Gallai [4]) $ext(n, kK_2) = \max\{\binom{2k-1}{2}, \binom{k-1}{2} + (k-1)(n-k+1)\}$ for all $n \geq 2k$ and $k \geq 1$, that is, for any given graph G of order n , if $|E(G)| > \max\{\binom{2k-1}{2}, \binom{k-1}{2} + (k-1)(n-k+1)\}$, then G contains a kK_2 , or a matching of size k .

In 2004, Schiermeyer [10] used some counting technique and determined the rainbow numbers $rb(K_n, kK_2)$ for all $k \geq 2$ and $n \geq 3k + 3$.

Theorem 1.2 (Schiermeyer [10]) $rb(n, kK_2) = ext(n, (k-1)K_2) + 2$ for all $k \geq 2$ and $n \geq 3k + 3$.

It is easy to see that n must be at least $2k$. So, for $2k \leq n < 3k + 3$, the rainbow numbers remain not determined. In this paper, we will use a technique deferent from Schiermeyer [10] to determine the exact values of $rb(n, kK_2)$ for all $k \geq 2$ and $n \geq 2k$. Our technique is to use the Gallai-Edmonds structure theorem for matchings.

Theorem 1.3

$$rb(n, kK_2) = \begin{cases} 4, & n = 4 \text{ and } k = 2; \\ ext(n, (k-1)K_2) + 3, & n = 2k \text{ and } k \geq 2; \\ ext(n, (k-1)K_2) + 2, & \text{otherwise.} \end{cases}$$

2 Preliminaries

Let M be a matching in a given graph G . Then the subgraph of G induced by M , denoted by $\langle M \rangle_G$ or $\langle M \rangle$, is the subgraph of G whose edge set is M and whose vertex set consists of the vertices incident with some edges in M . A vertex of G is said to be *saturated* by M if it is incident with an edge of M ; otherwise, it is said to be *unsaturated*. If every vertex of a vertex subset U of G is saturated, then we say that U is saturated by M . A matching with maximum cardinality is called a *maximum matching*.

In a given graph G , $N_G(U)$ denotes the set of vertices of G adjacent to a vertex of U . If $R, T \in V(G)$, we denote $E_G(R, T)$ or $E(R, T)$ as the set of all edges having a vertex from both R and T . Let $G(m, n)$ denote a bipartite graph with bipartition $A \cup B$, and

$|A| = m$ and $|B| = n$. Without loss of generality, in the following we always assume that $m \geq n$.

Let $\text{ext}(m, n, H)$ denote the maximum number of edges that a bipartite graph $G(m, n)$ can have with no subgraph isomorphic to H . The following lemma is due to Ore and can be found in [8].

Lemma 2.1 *Let $G(m, n)$ be a bipartite graph with bipartition $A \cup B$, and M a maximum matching in G . Then the size of M is $m - d$, where*

$$d = \max\{|S| - |N_G(S)| : S \subseteq A\}.$$

We now determine the value $\text{ext}(m, n, H)$ for $H = kK_2$.

Theorem 2.2

$$\text{ext}(m, n, kK_2) = m(k - 1) \text{ for all } n \geq k \geq 1,$$

that is, for any given bipartite graph $G(m, n)$, if $|E(G(m, n))| > m(k - 1)$, then $kK_2 \subset G(m, n)$.

Proof. Suppose that G contains no kK_2 . Let M be a maximum matching of G and the size of M is $k - i$, where $i \geq 1$. By Lemma 2.1, there exists a subset $S \subset A$ such that $|S| - |N_G(S)| = m - k + i$. Thus

$$|E(G)| \leq |S||N_G(S)| + n(m - |S|) = (|N_G(S)| + m - k + i)|N_G(S)| + n(k - i - |N_G(S)|).$$

Since $0 \leq |N_G(S)| \leq k - i \leq k - 1$, we obtain

$$|E(G)| \leq \max\{m(k - 1), n(k - 1)\} = m(k - 1).$$

So, $\text{ext}(m, n, kK_2) = m(k - 1)$. ■

Lemma 2.3

$$\text{ext}(2k, (k - 1)K_2) = \begin{cases} \binom{k-2}{2} + (k - 2)(k + 2), & 2 \leq k \leq 7; \\ \binom{2k-3}{2}, & k = 2 \text{ or } k \geq 7. \end{cases}$$

Proof. From Theorem 1.1, we have that $\text{ext}(2k, (k - 1)K_2) = \max\{\binom{2k-3}{2}, \binom{k-2}{2} + (k - 2)(k + 2)\}$. Since $\binom{2k-3}{2} - (\binom{k-2}{2} + (k - 2)(k + 2)) = \frac{1}{2}(k - 2)(k - 7)$, we have that if $2 \leq k \leq 7$, $\text{ext}(2k, (k - 1)K_2) = \binom{k-2}{2} + (k - 2)(k + 2)$, and if $k = 2$ or $k \geq 7$, $\text{ext}(2k, (k - 1)K_2) = \binom{2k-3}{2}$. ■

Let G be a graph. Denote by $D(G)$ the set of all vertices in G which are not covered by at least one maximum matching of G . Let $A(G)$ be the set of vertices in $V(G) - D(G)$

adjacent to at least one vertex in $D(G)$. Finally let $C(G) = V(G) - A(G) - D(G)$. We denote the $D(G)$, $A(G)$ and $C(G)$ as the *canonical decomposition* of G .

A *near-perfect* matching in a graph G is a matching of G covering all but exactly one vertex of G . A graph G is said to be *factor-critical* if $G - v$ has a perfect matching for every $v \in V(G)$.

Theorem 2.4 (*The Gallai-Edmonds Structure Theorem [8]*) For a graph G , let $D(G)$, $A(G)$ and $C(G)$ be defined as above. Then

- (a) The components of the subgraph induced by $D(G)$ are factor-critical.
- (b) The subgraph induced by $C(G)$ has a perfect matching.
- (c) The bipartite graph obtained from G by deleting the vertices of $C(G)$ and the edges spanned by $A(G)$ and by contracting each component of $D(G)$ to a single vertex has positive surplus (as viewed from $A(G)$).
- (d) Any maximum matching M of G contains a near-perfect matching of each component of $D(G)$, a perfect matching of each component of $C(G)$ and matches all vertices of $A(G)$ with vertices in distinct components of $D(G)$.
- (e) The size of a maximum matching M is $\frac{1}{2}(|V(G)| - c(D(G)) + |A(G)|)$, where $c(D(G))$ denotes the number of components of the graph spanned by $D(G)$. ■

3 Main results

For $k = 1$, it is clear that $rb(n, K_2) = 1$. Now we determine the value of $rb(n, 2K_2)$ (for $k = 2$).

Theorem 3.1

$$rb(4, 2K_2) = 4,$$

and

$$rb(n, 2K_2) = 2 = ext(n, K_2) + 2 \text{ for all } n \geq 5.$$

Proof. It is obvious that $rb(4, 2K_2) \leq 4$. Let $V(K_4) = \{a_1, a_2, a_3, a_4\}$. If K_4 is edge-colored with 3 colors such that $c(a_1a_2) = c(a_3a_4) = 1$, $c(a_1a_3) = c(a_2a_4) = 2$ and $c(a_1a_4) = c(a_2a_3) = 3$, then K_4 contains no TMC $2K_2$. So, $rb(4, 2K_2) = 4$.

For $n \geq 5$, let the edges of $G = K_n$ be colored with at least 2 colors. Suppose that K_n contains no TMC $2K_2$. Let $e_1 = a_1b_1$ be an edge with $c(e_1) = 1$, $T = \{a_1, b_1\}$ and $R = V(K_n) - T$. Then $c(e) = 1$ for all edges $e \in E(G[R])$. Moreover, $c(e) = 1$ for all edges $e \in E(T, R)$, since $|R| \geq 3$. But then K_n is monochromatic, a contradiction. So, $rb(n, 2K_2) = 2$ for all $n \geq 5$. ■

The next proposition provides a lower and upper bound for $rb(n, kK_2)$.

Proposition 3.2 $ext(n, (k-1)K_2) + 2 \leq rb(n, kK_2) \leq ext(n, kK_2) + 1.$

Proof. The upper bound is obvious. For the lower bound, an extremal coloring of K_n can be obtained from an extremal graph S_n for $ext(n, (k-1)K_2)$ by coloring the edges of S_n differently and the edges of $\overline{S_n}$ by one extra color. It is obvious that the coloring does not contain a TMC kK_2 . ■

We will show that the lower bound can be achieved for all $n \geq 2k + 1$ and $k \geq 3$, and thus obtain the exact value of $rb(n, kK_2)$ for all $n \geq 2k + 1$ and $k \geq 3$.

For $n = 2k$, we suppose that $H = K_{2k-3}$ is a subgraph of K_n and $V(K_n) - V(H) = \{a_1, a_2, a_3\}$. If K_n is edge-colored such that $c(a_1a_2) = 1$, $c(a_1a_3) = c(a_2a_3) = 2$, $c(e) = 1$ for all edges $e \in E(a_3, V(H))$, $c(e) = 2$ for all edges $e \in E(a_1, V(H)) \cup E(a_2, V(H))$ and the edges of $H = K_{2k-3}$ is colored differently by $\binom{2k-3}{2}$ extra colors. It is easy to check that the coloring does not contain a TMC kK_2 in K_n . So, $rb(2k, kK_2) \geq \binom{2k-3}{2} + 3$ for all $k \geq 3$. Hence, if $k \geq 7$, then $ext(2k, (k-1)K_2) = \binom{2k-3}{2}$ and $rb(2k, kK_2) \geq ext(2k, (k-1)K_2) + 3$. We will show that the lower bound can be achieved for all $n \geq 2k$ and $k \geq 7$.

Theorem 3.3 *For all $n \geq 2k$ and $k \geq 3$, we have*

$$rb(n, kK_2) = \begin{cases} ext(n, (k-1)K_2) + 3, & n = 2k \text{ and } k \geq 7; \\ ext(n, (k-1)K_2) + 2, & \text{otherwise.} \end{cases}$$

Proof. We shall prove the theorem by contradiction. If $n = 2k$ and $k \geq 7$, let the edges of K_n be colored with $ext(n, (k-1)K_2) + 3$ colors; otherwise, let the edges of K_n be colored with $ext(n, (k-1)K_2) + 2$ colors. Suppose that K_n contains no TMC kK_2 . Now let $G \subset K_n$ be a TMC spanning subgraph which contains all colors in K_n , i.e., if $n = 2k$ and $k \geq 7$, $|E(G)| = ext(n, (k-1)K_2) + 3$; otherwise $|E(G)| = ext(n, (k-1)K_2) + 2$. Since $|E(G)| \geq ext(n, (k-1)K_2) + 2$, there is a TMC $(k-1)K_2$ in G .

We first need to prove the following two lemmas.

Lemma 3.4 *If two components of G consist of a K_{2k-3} and a K_3 , respectively, and the other components are isolated vertices (see Figure 1), then K_n contains a TMC kK_2 .*

Proof. Denote SG_1 as the special graph G and Q as the set of isolated vertices of G . Without loss of generality, we suppose that $c(u_1u_2) = 1, c(u_2u_3) = 2, c(u_1u_3) = 3, c(v_1v_2) = 4, c(v_2v_3) = 5, c(v_1v_3) = 6$ (see Figure 1).

The proof of the lemma is given by distinguishing the following two cases:

Case I. $k \geq 4$.




Figure 1: The special graph SG_1 .




Figure 2: The special graph SG_2 . G' and G'' is a K_{2k-3} and a P_3 , respectively, or G' and G'' is a K_{2k-3}^- and a K_3 , respectively.

We suppose that G contains no TMC kK_2 . We will show $c(u_1v_1) = 5$. If $c(u_1v_1) \neq 5$, then in $G_1 = K_{2k-3} - u_1$ the number of edges whose colors are not $c(u_1v_1)$ is at least $\binom{2k-4}{2} - 1$. Since $k \geq 4$, we have $\binom{2k-4}{2} - 1 > \text{ext}(2k-4, (k-2)K_2) = \binom{2k-5}{2}$. Thus we can obtain a TMC $H = (k-2)K_2$ which contains no color $c(u_1v_1)$ in G_1 , and hence there is a TMC $kK_2 = H \cup \{u_1v_1, v_2v_3\}$ in K_n . So, $c(u_1v_1)$ must be 5. By the same token, $c(u_2v_2)$ and $c(u_3v_3)$ must be 6 and 4, respectively. Now we can obtain a TMC $H' = (k-3)K_2$ in $G_2 = K_{2k-3} - u_1 - u_2 - u_3$, and hence there is a TMC $kK_2 = H' \cup \{u_1v_1, u_2v_2, u_3v_3\}$ in K_n .

Case II. $k = 3$.

We suppose that K_n contains no TMC $3K_2$. Then $c(u_1v_1) \in \{2, 5\}, c(u_2v_2) \in \{3, 6\}, c(u_3v_3) \in \{1, 4\}$. Now we can obtain a TMC $3K_2 = u_1v_1 \cup u_2v_2 \cup u_3v_3$ in K_n . ■

Lemma 3.5 *If $n \geq 2k + 1$ and two components of G are G' and G'' , where G' and G'' is a K_{2k-3} and a P_3 , respectively, or G' and G'' is a K_{2k-3}^- and a K_3 , respectively, and the other components are isolated vertices (see Figure 2), then K_n contains a TMC kK_2 , where P_3 is a path with three vertices and K_{2k-3}^- is obtained from K_{2k-3} by deleting an edge.*




Figure 3: We can obtain a TMC $3K_2 = u_1v_4 \cup u_3v_3 \cup u_2v_1$ in K_n .

Proof. Denote SG_2 as the special graph G and Q as the set of isolated vertices of G . Without loss of generality, we suppose that $c(u_1u_2) = 1, c(u_2u_3) = 2, c(u_1u_3) = 3, c(v_1v_2) = 4, c(v_2v_3) = 5$ (see Figure 2). The proof of the lemma is given by distinguishing the following two cases:

Case I. $k \geq 4$.

Since $n \geq 2k + 1$, we suppose that $v_4 \in Q$. If $c(u_1v_4) = j$, without loss of generality, we suppose that $j \neq 4$. The number of edges of $G' - u_1$ whose color is not j is at least $\binom{2k-4}{2} - 2$ and $\binom{2k-4}{2} - 2 > \text{ext}(2k - 4, (k - 2)K_2) = \binom{2k-5}{2}$. Then there is a TMC $H = (k - 2)K_2$ in $G' - u_1$ which contains no color j . We can obtain a TMC $kK_2 = H \cup u_1v_4 \cup v_1v_2$ in K_n .

Case II. $k = 3$.

Without loss of generality, we suppose that G' and G'' is a K_3 and a P_3 , respectively. We suppose that K_n contains no TMC $3K_2$. Then, $c(u_1v_4) \in \{2, 5\} \cap \{2, 4\}$, i.e., $c(u_1v_4) = 2$, $c(u_3v_3) \in \{2, 4\} \cap \{1, 4\}$, i.e., $c(u_3v_3) = 4$, $c(u_2v_1) \in \{2, 5\} \cap \{3, 5\}$, i.e., $c(u_2v_1) = 5$. Now we obtain a TMC $3K_2 = u_1v_4 \cup u_3v_3 \cup u_2v_1$. See Figure 3. ■

Now we turn back to the proof of Theorem 3.3. Let $D(G), A(G), C(G)$ as the canonical decomposition of G and $c(D(G)) = q, |A(G)| = s, |V(G)| = n$. Since the size of the maximum matchings of G is $k - 1$, by Theorem 2.4 (e), $k - 1 = \frac{1}{2}(n - q + s)$, i.e., $q = n - 2k + 2 + s$. Let the components of $D(G)$ be D_1, D_2, \dots, D_q . By Theorem 2.4 (a), the components of the subgraph induced by $D(G)$ are factor-critical, hence we suppose that $|V(D_i)| = 2l_i + 1$ for $1 \leq i \leq q$, without loss of generality, $l_1 \geq l_2 \geq \dots \geq l_q \geq 0$. Let the components of $C(G)$ be $C_1, C_2, \dots, C_{q'}$ with $|V(C_i)| = 2t_i$ for $1 \leq i \leq q'$.

Since $s + q = s + n - 2k + 2 + s \leq n$, then $0 \leq s \leq k - 1$. Moreover,

$$\begin{aligned}
n = s + \sum_{i=1}^q (2l_i + 1) + |C(G)| &\geq s + (2l_1 + 1) + \sum_{i=2}^q (2l_i + 1) \\
&\geq s + (2l_1 + 1) + (q - 1) \\
&\geq s + (2l_1 + 1) + (n - 2k + 2 + s - 1),
\end{aligned}$$

hence $2l_1 + 1 \leq 2k - 2s - 1$. We distinguish four cases to finish the proof of Theorem 3.3.

Case 1. $s = k - 1$.

In this case, since $s + q = (k - 1) + n - 2k + 2 + (k - 1) = n$, then $C(G) = \emptyset$ and $l_1 = l_2 = \dots = l_q = 0$. The components of the subgraph induced by $D(G)$ are isolated vertices. We distinguish two subcases to finish the proof of the case.

Subcase 1.1. There is at most one vertex u in $D(G)$ such that $d_G(u) < k - 1$.

We suppose $v \in D(G)$ and $u \neq v$. Let $G(n - k - 1, k - 1)$ be the bipartite graph obtained from G by deleting the vertices u, v and the edges spanned by $A(G)$. It is obvious that $uv \in E(K_n)$ and $uv \notin E(G)$, without loss of generality, we suppose $c(uv) = 1$. Then the number of edges in $G(n - k - 1, k - 1)$ whose color is not 1 is at least $(n - k - 1)(k - 1) - 1$. Since $n - k - 1 \geq 2$, then $(n - k - 1)(k - 1) - 1 > \text{ext}(n - k - 1, k - 1, (k - 1)K_2) = (n - k - 1)(k - 2)$. By Lemma 2.2, there exists a TMC $H = (k - 1)K_2$ in $G(n - k - 1, k - 1)$ which contains no color 1, thus we obtain a TMC $kK_2 = H \cup uv$ in K_n .

Subcase 1.2. There exist at least two vertices u, v in $D(G)$ such that $d_G(u) < k - 1$ and $d_G(v) < k - 1$.

We suppose that $c(uv) = 1$. Let $G'(n - k - 1, k - 1)$ be the bipartite graph obtained from G by deleting the vertices u, v and the edges spanned by $A(G)$ and the edge whose color is 1. Thus there is no TMC $(k - 1)K_2$ in $G'(n - k - 1, k - 1)$. Hence, by Lemma 2.2,

$$\begin{aligned} |E(G)| &\leq 1 + \text{ext}(n - k - 1, k - 1, (k - 1)K_2) + 2(k - 2) + \binom{k - 1}{2} \\ &\leq 1 + (k - 2)(n - k - 1) + 2(k - 2) + \binom{k - 1}{2} \\ &= \binom{k - 2}{2} + (k - 2)(n - k + 2) + 1 \\ &< \text{ext}(n, (k - 1)K_2) + 2, \end{aligned}$$

which contradicts $|E(G)| \geq \text{ext}(n, (k - 1)K_2) + 2$.

Case 2. $0 \leq s \leq k - 2$ and $2l_1 + 1 \leq 2k - 2s - 3$.

In this case, if $2k - 2s - 3 = 1$, then $l_1 = l_2 = \dots = l_q = 0$, $s = k - 2$ and $|C(G)| = 2$, hence

$$\begin{aligned} |E(G)| &\leq \binom{s}{2} + s(n - s) + \binom{2}{2} \\ &= \binom{k - 2}{2} + (k - 2)(n - k + 2) + 1 \\ &< \text{ext}(n, (k - 1)K_2) + 2, \end{aligned}$$

which contradicts $|E(G)| \geq \text{ext}(n, (k - 1)K_2) + 2$.

If $2k - 2s - 3 \geq 3$, then $0 \leq s \leq k - 3$ and

$$\begin{aligned} \sum_{i=2}^q (2l_i + 1) + \sum_{i=1}^{q'} (2t_i) &= n - s - (2l_1 + 1) \\ &\geq n - s - (2k - 2s - 3) = (q - 1) + 2. \end{aligned}$$

Thus, if $|C(G)| \geq 2$, then

$$\begin{aligned} |E(G)| &\leq \binom{s}{2} + s(n - s) + \sum_{i=1}^q \binom{2l_i + 1}{2} + \sum_{i=1}^{q'} \binom{2t_i}{2} \\ &\leq \binom{s}{2} + s(n - s) + \binom{2l_1 + 1 + \sum_{i=2}^q 2l_i}{2} + \sum_{i=1}^{q'} \binom{2t_i}{2} \\ &\leq \binom{s}{2} + s(n - s) + \binom{2l_1 + 1 + \sum_{i=2}^q 2l_i + (\sum_{i=1}^{q'} 2t_i - 2)}{2} + \binom{2}{2} \\ &= \binom{s}{2} + s(n - s) + \binom{n - s - (q - 1) - 2}{2} + \binom{2}{2} \\ &= \binom{s}{2} + s(n - s) + \binom{2k - 2s - 3}{2} + \binom{2}{2} := f_1(s). \end{aligned}$$

Hence,

$$\begin{aligned} f_1(0) &= \binom{2k - 3}{2} + 1 < \text{ext}(n, (k - 1)K_2) + 2, \\ f_1(k - 3) &= \binom{k - 2}{2} + (k - 2)(n - k + 2) - (n - k) + 2 \\ &< \binom{k - 2}{2} + (k - 2)(n - k + 2) < \text{ext}(n, (k - 1)K_2) + 2. \end{aligned}$$

Since $0 \leq s \leq k - 3$, $|E(G)| \leq \max\{f_1(0), f_1(k - 3)\} < \text{ext}(n, (k - 1)K_2) + 2$, which contradicts $|E(G)| \geq \text{ext}(n, (k - 1)K_2) + 2$.

If $|C(G)| = 0$, then $2l_2 + 1 \geq 3$ and

$$\begin{aligned} |E(G)| &\leq \binom{s}{2} + s(n - s) + \sum_{i=1}^q \binom{2l_i + 1}{2} + \sum_{i=1}^{q'} \binom{2t_i}{2} \\ &\leq \binom{s}{2} + s(n - s) + \binom{2l_1 + 1 + \sum_{i=3}^q 2l_i + \sum_{i=1}^{q'} 2t_i}{2} + \binom{2l_2 + 1}{2} \\ &\leq \binom{s}{2} + s(n - s) + \binom{2l_1 + 1 + \sum_{i=3}^q 2l_i + \sum_{i=1}^{q'} 2t_i + (2l_2 - 2)}{2} + \binom{3}{2} \\ &= \binom{s}{2} + s(n - s) + \binom{n - s - (q - 1) - 2}{2} + \binom{3}{2} \\ &= \binom{s}{2} + s(n - s) + \binom{2k - 2s - 3}{2} + \binom{3}{2} := f_2(s). \end{aligned}$$

Thus,

$$\begin{aligned}
f_2(0) &= \binom{2k-3}{2} + 3, \\
f_2(1) &= \binom{2k-3}{2} + n - 4k + 11, \\
f_2(k-3) &= \binom{k-2}{2} + (k-2)(n-k+2) - (n-k) + 4 \\
&\leq \binom{k-2}{2} + (k-2)(n-k+2) + 1 < \text{ext}(n, (k-1)K_2) + 2.
\end{aligned}$$

If $s = 0$ and $|E(G)| = \binom{2k-3}{2} + 3$, then $G \cong SG_1$. By Lemma 3.4, we can obtain a TMC kK_2 in K_n . If $s = 0$, $n \geq 2k + 1$ and $|E(G)| = \binom{2k-3}{2} + 2$, then $G \cong SG_2$. By Lemma 3.5, we can obtain a TMC kK_2 in K_n . So, if $n \geq 2k + 1$, then $|E(G)| \leq \binom{2k-3}{2} + 1 < \text{ext}(n, (k-1)K_2) + 2$, which contradicts $|E(G)| = \text{ext}(n, (k-1)K_2) + 2$. If $n = 2k$ and $k \geq 7$, then $|E(G)| \leq \binom{2k-3}{2} + 2 = \text{ext}(n, (k-1)K_2) + 2$, which contradicts $|E(G)| = \text{ext}(n, (k-1)K_2) + 3$. If $n = 2k$ and $3 \leq k \leq 6$, then $|E(G)| \leq \binom{2k-3}{2} + 2 \leq \binom{k-2}{2} + (k-2)(k+2) = \text{ext}(n, (k-1)K_2)$, which contradicts $|E(G)| = \text{ext}(n, (k-1)K_2) + 2$.

If $1 \leq s \leq k-3$, then $k \geq 4$ and $|E(G)| \leq \max\{f_2(1), f_2(k-3)\}$. So, if $f_2(k-3) \geq f_2(1)$, then $|E(G)| \leq f_2(k-3) < \text{ext}(n, (k-1)K_2) + 2$, a contradiction. If $f_2(1) > f_2(k-3)$, then

$$\binom{2k-3}{2} + n - 4k + 11 > \binom{k-2}{2} + (k-2)(n-k+2) - (n-k) + 4.$$

Hence $2k \leq n < \frac{1}{2}(5k-7)$, $k > 7$ and

$$\begin{aligned}
|E(G)| &\leq f_2(1) = \binom{2k-3}{2} + n - 4k + 11 \\
&< \binom{2k-3}{2} + \frac{1}{2}(15-3k) \\
&< \text{ext}(n, (k-1)K_2) + 2,
\end{aligned}$$

a contradiction.

Case 3. $0 \leq s \leq k-2$, $2l_1 + 1 = 2k - 2s - 1$ and $n \geq 2k + 1$.

In this case, $s + (2l_1 + 1) + (q-1) = n$, hence $C(G) = \emptyset$, $l_2 = l_3 = \dots = l_q = 0$ and each D_i for $2 \leq i \leq q$ is an isolated vertex.

Let $G(q, s)$ be the bipartite graph obtained from G by deleting the edges spanned by $A(G)$ and by contracting the component D_1 to a single vertex p . Thus by Theorem 2.4 (c) and (d), we can obtain a maximum matching M of size $k-1$ such that M contains a maximum matching M_1 of $G(q, s)$ which does not match vertex p and a near-perfect matching M_2 of D_1 . Since $q = n - 2k + 2 + s \geq s + 3$, there exist two vertices $u, v \in D(G) - D_1$ and $u, v \notin \langle M \rangle$. It is obvious that $uv \in E(K_n)$ and $uv \notin E(G)$. We




Figure 4: If $yz_1 \in E_G(y, D_1)$, we can obtain a TMC $kK_2 = M'_1 \cup M'_2 \cup uv$ in K_n .

suppose that $c(uv) = 1$, hence there exists an edge $e = yz \in M$ with $c(e) = 1$. Now we distinguish two subcases to complete the proof of the case.

Subcase 3.1. $e \in M_1$.

In this subcase, $s \geq 1$ and $yz \in E_G(A(G), D(G))$, without loss of generality, we suppose that $y \in A(G)$. If there exists an edge $yz_1 \in E_G(y, D_1)$ with $z_1 \in D_1$, then we can obtain another maximum matching M'_1 of $G(q, s)$ with $M'_1 = M_1 \cup yz_1 - yz$ and a near-perfect matching M'_2 of D_1 which does not match z_1 . Thus we obtain a TMC $kK_2 = M'_1 \cup M'_2 \cup uv$ in K_n . See Figure 4.

Thus we suppose that $E_G(y, D_1) = \emptyset$. There is no matching of size s in $G'(q-3, s) = G(q, s) - p - u - v - e$. By Lemma 2.2, $|E_G(G')| \leq (s-1)(q-3) = (s-1)(n-2k+s-1)$. Now

$$\begin{aligned}
|E(G)| &\leq \binom{s}{2} + \binom{2k-2s-1}{2} + 1 + |E_G(G')| \\
&\quad + |E_G(D_1, A(G))| + |E_G(\{u, v\}, A(G))| \\
&\leq \binom{s}{2} + \binom{2k-2s-1}{2} + 1 + (s-1)(n-2k+s-1) \\
&\quad + (2k-2s-1)(s-1) + 2s := f_3(s)
\end{aligned}$$

Hence,

$$\begin{aligned}
f_3(1) &= \binom{2k-3}{2} + 3, \\
f_3(2) &= \binom{2k-3}{2} + n - 4k + 11, \\
f_3(k-2) &= \binom{k-2}{2} + (k-2)(n-k+2) - (n-k) + 4 \\
&\leq \binom{k-2}{2} + (k-2)(n-k+2) < \text{ext}(n, (k-1)K_2) + 2.
\end{aligned}$$

If $s = 1$, then $|E(G)| \leq \binom{2k-3}{2} + 3$. If $|E(G)| = \binom{2k-3}{2} + 3$, then $(G - e + uv) \cong SG_1$. By the proof of Lemma 3.4, we can obtain a TMC kK_2 in K_n . If $|E(G)| = \binom{2k-3}{2} + 2$, then

$(G - e + uv) \cong SG_2$. By the proof of Lemma 3.5, we can obtain a TMC kK_2 in K_n . If $|E(G)| \leq \binom{2k-3}{2} + 1 \leq \text{ext}(n, (k-1)K_2) + 1$, this contradicts $|E(G)| = \text{ext}(n, (k-1)K_2) + 2$.

If $2 \leq s \leq k-2$, then $k \geq 4$ and $|E(G)| \leq \max\{f_3(2), f_3(k-2)\}$. So, if $f_3(k-2) \geq f_3(2)$, then $|E(G)| \leq f_3(k-2) < \text{ext}(n, (k-1)K_2) + 2$, a contradiction. If $f_3(2) > f_3(k-2)$, then

$$\binom{2k-3}{2} + n - 4k + 11 > \binom{k-2}{2} + (k-2)(n-k+2) - (n-k) + 4.$$

Hence, $2k \leq n < \frac{1}{2}(5k-7)$, $k > 7$ and

$$\begin{aligned} |E(G)| &\leq f_3(2) = \binom{2k-3}{2} + n - 4k + 11 \\ &< \binom{2k-3}{2} + \frac{1}{2}(15-3k) \\ &< \text{ext}(n, (k-1)K_2) + 2, \end{aligned}$$

a contradiction.

Subcase 3.2. $e \in M_2$.

In this subcase, $y \in D_1$ and $z \in D_1$. By Theorem 2.4 (a), D_1 is factor-critical, there exists a near-perfect matching M'_2 which does not match y , So M'_2 does not contain $e = yz$. Now we obtain a TMC $kK_2 = M'_2 \cup M_1 \cup uv$ in K_n .

Case 4. $0 \leq s \leq k-2$, $2l_1 + 1 = 2k - 2s - 1$ and $n = 2k$.

In this case, $q = s + 2$ and $s + (2l_1 + 1) + (q - 1) = 2k$, hence $C(G) = \emptyset$, $l_2 = l_3 = \dots = l_q = 0$ and each D_i for $2 \leq i \leq q$ is an isolated vertex. Now we distinguish two subcases to complete the proof of the case.

Subcase 4.1. $1 \leq s \leq k-2$.

If $E_G(D_1, A(G)) = \emptyset$, then

$$|E(G)| \leq \binom{2k-2s-1}{2} + \binom{s}{2} + s(s+1) := f_4(s).$$

Thus,

$$\begin{aligned} f_4(1) &= \binom{2k-3}{2} + 2, \\ f_4(k-2) &= \binom{k-2}{2} + (k-2)(k+2) + 3 - 3(k-2) \end{aligned}$$

Since $k \geq 3$, then $f_4(1) \geq f_4(k-2)$ and $|E(G)| \leq \max\{f_4(1), f_4(k-2)\} = f_4(1) = \binom{2k-3}{2} + 2$. If $k \geq 7$, this contradicts $|E(G)| = \text{ext}(2k, (k-1)K_2) + 3 = \binom{2k-3}{2} + 3$. If $3 \leq k \leq 6$, then

$$\begin{aligned} |E(G)| &\leq \binom{2k-3}{2} + 2 \\ &\leq \binom{k-2}{2} + (k-2)(k+2) = \text{ext}(2k, (k-1)K_2), \end{aligned}$$




Figure 5: The special graph SG_3 and $|E(SG_3)| = \binom{2k-3}{2} + 3$.

which contradicts $|E(G)| = ext(2k, (k-1)K_2) + 2$.

So we suppose that $E_G(D_1, A(G)) \neq \emptyset$. Let $G(s+2, s)$ be the bipartite graph obtained from G by deleting the edges spanned by $A(G)$ and by contracting the component D_1 to a single vertex p . Thus by Theorem 2.4 (d), we can obtain a maximum matching M of size $k-1$ such that M contains a near-perfect matching M_1 of D_1 which does not match w with $w \in D_1$ and a matching M_2 of size s which matches all vertices of $A(G)$ with vertices in $\{w\} \cup (D(G) - D_1)$. Since $E_G(D_1, A(G)) \neq \emptyset$, we can suppose that $w \in \langle M_2 \rangle$. There exist exactly two vertices $u, v \in D(G) - D_1$ and $u, v \notin \langle M \rangle$. It is obvious that $uv \in E(K_n)$ and $uv \notin E(G)$. We suppose that $c(uv) = 1$, hence there exists an edge $e = yz \in M$ with $c(e) = 1$. Now we distinguish two subcases to complete the proof of the subcase 4.1.

Subcase 4.1.1. $e = yz \in M_1$.

If $s = 1$, then $|D_1| = 2k-3$ and we suppose $A(G) = \{x\}$. Thus the size of M_1 is $k-2$ and there is no $H = (k-2)K_2$ in $D'_1 = D_1 - w - yz$, for otherwise, we can obtain a TMC $kK_2 = H \cup xw \cup uv$ in K_{2k} . If $E_G(x, \{y, z\}) \neq \emptyset$, say $xy \in E(G)$, then we can obtain a perfect matching M'_1 of $D_1 - y$ and a TMC $kK_2 = M'_1 \cup uv \cup xy$ in K_{2k} . So, $E_G(x, \{y, z\}) = \emptyset$ and

$$\begin{aligned}
|E(G)| &= 1 + |E_G(D'_1)| + |E_G(w, D'_1)| + |E_G(x, D_1)| + |E_G(x, \{u, v\})| \\
&\leq 1 + ext(2k-4, (k-2)K_2) + (2k-4) + (2k-5) + 2 \\
&= \binom{2k-5}{2} + 4k-6 \\
&= \binom{2k-3}{2} + 3.
\end{aligned}$$

Denote SG_3 be the special graph G shown in Figure 5, whence $E(SG_3) = E(K_{2k-3}^-) \cup xu \cup xv \cup yw \cup yz$. Without loss of generality, we suppose that $c(wy) = 4$. If $|E(G)| = \binom{2k-3}{2} + 3$, it is easy to check that $G \cong SG_3$.

If $k \geq 7$, then by the beginning hypothesis $|E(G)| = ext(2k, (k-1)K_2) + 3 = \binom{2k-3}{2} + 3$, whence $G \cong SG_3$. Now $\binom{2k-4}{2} - 1 > ext(2k-4, (k-2)K_2)$, we can obtain a TMC $H = (k-2)K_2$ in $K_{2k-3}^- - w$, whence a TMC $kK_2 = H \cup yw \cup uv$ in K_{2k} .




Figure 6: There is no $(k - s - 1)K_2$ in $D'_1 = D_1 - w - yz$. If $x'y \in E(G)$, there is no $(s - 1)K_2$ in bipartite graph $G'(s - 1, s - 1) = G - \{D_1 \cup u \cup v \cup x'\}$.

If $3 \leq k \leq 6$, then

$$\binom{2k - 3}{2} + 3 \leq \binom{k - 2}{2} + (k - 2)(k + 2) + 1 = \text{ext}(2k, (k - 1)K_2) + 1,$$

which contradicts $|E(G)| = \text{ext}(2k, (k - 1)K_2) + 2$. If $2 \leq s \leq k - 2$, then $k \geq 4$. We suppose that $x \in A(G)$ and $xw \in M_2$. By the the same token, $E_G(x, \{y, z\}) = \emptyset$ and there is no $(k - s - 1)K_2$ in $D'_1 = D_1 - w - yz$.

If $E_G(A(G) - x, \{y, z\}) \neq \emptyset$, say $x'y \in E(G)$, then there is no $H = (s - 1)K_2$ in bipartite graph $G'(s - 1, s - 1) = G - \{D_1 \cup u \cup v \cup x'\}$, for otherwise, we can obtain a perfect matching M'_1 in $D_1 - y$ and a TMC $kK_2 = M'_1 \cup H \cup uv \cup x'y$. See Figure 6.

Thus,

$$\begin{aligned} |E_G(A(G), D(G))| &= |E_G(A(G), D_1 - y - z)| + |E(A(G), \{y, z\})| \\ &\quad + |E_G(A(G), \{u, v\})| + |E_G(G'(s - 1, s - 1))| \\ &\quad + |E_G(x', D(G) - D_1 - u - v)| \\ &\leq (2k - 2s - 3)s + 2(s - 1) + 2s \\ &\quad + \text{ext}(s - 1, s - 1, (s - 1)K_2) + (s - 1) \\ &= (2k - 2s - 3)s + 2s + (s - 1)(s + 1). \end{aligned}$$

If $E_G(A(G) - x, \{y, z\}) = \emptyset$, then

$$\begin{aligned} |E_G(A(G), D(G))| &= |E_G(A(G), D_1 - y - z)| + |E_G(A(G), D(G) - D_1)| \\ &\leq (2k - 2s - 3)s + s(s + 1). \end{aligned}$$

So,

$$\begin{aligned} &|E_G(A(G), D(G))| \\ &\leq \max\{(2k - 2s - 3)s + 2s + (s - 1)(s + 1), (2k - 2s - 3)s + s(s + 1)\} \\ &= (2k - 2s - 3)s + 2s + (s - 1)(s + 1). \end{aligned}$$

Now, we have

$$\begin{aligned}
|E(G)| &= \binom{s}{2} + 1 + |E_G(D'_1)| + |E_G(w, D'_1)| + |E_G(A(G), D(G))| \\
&\leq \binom{s}{2} + 1 + \binom{2k-2s-3}{2} + (2k-2s-2) + \\
&\quad (2k-2s-3)s + 2s + (s-1)(s+1) := f_5(s).
\end{aligned}$$

Thus,

$$\begin{aligned}
f_5(2) &= \binom{2k-3}{2} - 2k + 11, \\
f_5(k-2) &= \binom{k-2}{2} + (k-2)(k+2) - k + 4 \\
&< \text{ext}(2k, (k-1)K_2) + 2.
\end{aligned}$$

If $4 \leq k \leq 6$, then $f_5(k-2) \geq f_5(2)$ and $|E(G)| \leq \max\{f_5(2), f_5(k-2)\} = f_5(k-2) < \text{ext}(2k, (k-1)K_2) + 2$, which contradicts $|E(G)| = \text{ext}(2k, (k-1)K_2) + 2$.

If $k \geq 7$, then $f_5(2) \geq f_5(k-2)$ and $|E(G)| \leq \max\{f_5(2), f_5(k-2)\} = f_5(2) = \binom{2k-3}{2} - 2k + 11 < \binom{2k-3}{2} = \text{ext}(2k, (k-1)K_2)$, which contradicts $|E(G)| = \text{ext}(2k, (k-1)K_2) + 3$.

Subcase 4.1.2. $e = yz \in M_2$.

Without loss of generality, we suppose that $y \in A(G)$.

If $s = 1$, then $A(G) = \{y\}$, $yz = yw$ and $c(yw) = c(uv) = 1$. Then $E_G(y, D_1 - w) = \emptyset$, for otherwise, say $yw' \in E_G(y, D_1 - w)$ with $w' \in (D_1 - w)$, we can obtain a TMC $H = (k-2)K_2$ in $D_1 - w'$ and a TMC $kK_2 = H \cup yw' \cup uv$ in K_{2k} . So,

$$|E(G)| = |E_G(D_1)| + |E_G(y, \{w, u, v\})| \leq \binom{2k-3}{2} + 3.$$

If $3 \leq k \leq 6$, then

$$\binom{2k-3}{2} + 3 \leq \binom{k-2}{2} + (k-2)(k+2) + 1 = \text{ext}(2k, (k-1)K_2) + 1,$$

which contradicts $|E(G)| = \text{ext}(2k, (k-1)K_2) + 2$.

If $k \geq 7$, since $|E(G)| = \binom{2k-3}{2} + 3$, it is easy to check that $(G - e + uv) \cong SG_1$. By the proof of Lemma 3.4, we can obtain a TMC kK_2 in K_{2k} .

If $2 \leq s \leq k-2$, first we look at the bipartite graph $G(s+2, s)$. We suppose that M'_2 is any maximum matching of size s in $G(s+2, s)$ with $p \in \langle M'_2 \rangle$ and $u_1, v_1 \notin \langle M'_2 \rangle$. By Subcase 4.1.1, we can suppose that there exists an edge $e_1 \in M'_2$ such that $c(e_1) = c(u_1v_1)$. If $d_{G(s+2,s)}(p) = s$ and there is at most one vertex u_2 in $D(G) - D_1$ such that $d_{G(s+2,s)}(u) \leq s-1$, we suppose $v_2 \in D(G) - D_1$ and $u_2 \neq v_2$. Let $G(s, s)$ be the bipartite graph obtained from $G(s+2, s)$ by deleting the vertices u_2, v_2 . It is obvious that $u_2v_2 \in E(K_n)$ and $u_2v_2 \notin E(G)$. Then the number of edges in $G(s, s)$ whose color

is not $c(u_2v_2)$ is at least $s^2 - 1$. Since $s \geq 2$, then $s^2 - 1 \geq \text{ext}(s, s, sK_2) = s(s-1) + 1$. By Lemma 2.2, there exists a TMC $H = sK_2$ in $G(s, s)$ which contains no color $c(u_2v_2)$, thus we obtain a TMC $(s+1)K_2 = H \cup u_2v_2$. By Theorem 2.4, we can obtain a TMC kK_2 in K_{2k} .

So, if $d_{G(s+2,s)}(p) = s$, then we suppose there exist at least two vertices u_3, v_3 in $D(G) - D_1$ such that $d_{G(s+2,s)}(u_3) \leq s-1$ and $d_{G(s+2,s)}(v_3) \leq s-1$. Let $G'(s, s)$ be the bipartite graph obtained from $G(s+2, s)$ by deleting the vertices u_3, v_3 and the edge whose color is $c(u_3v_3)$. Thus there is no TMC sK_2 in $G'(s, s)$. By Lemma 2.2, $E(G(s+2, s)) \leq 1 + 2(s-1) + s(s-1)$ and

$$|E_G(A(G), D(G))| \leq 1 + 2(s-1) + s((2k-2s-1) + (s-2)) = 1 + 2(s-1) + s(2k-s-3).$$

Now we suppose that $d_{G(s+2,s)}(p) \leq s-1$. Since $E(A(G), D_1) \neq \emptyset$, if there exists an edge $w''x' \in E(A(G), D_1)$ with $x' \in A(G)$, $w'' \in D_1$ and $w''x' \neq wx$. Thus there is no TMC $H = (s-1)K_2$ in $G(s+2, s) - \{p \cup u \cup v \cup x'\} - yz$, for otherwise, we can obtain a TMC $(s+1)K_2 = H \cup uv \cup w''x'$, a TMC $(k-s-1)K_2$ in $D_1 - w''$ and a TMC kK_2 in K_{2k} . We have

$$\begin{aligned} |E_G(A(G), D(G))| &\leq |E_G(A(G), D_1)| + (s-1)(s-2) + 1 \\ &\quad + |E_G(x', D(G) - D_1 - u - v)| + |E_G(A(G), \{u, v\})| \\ &\leq (2k-2s-1)(s-1) + (s-1)(s-2) + 1 + (s-1) + 2s \\ &= (2k-2s-1)(s-1) + s^2 + 2. \end{aligned}$$

If $E(A(G), D_1) = \{xw\}$, then

$$|E_G(A(G), D(G))| \leq 1 + s(s+1).$$

Thus,

$$\begin{aligned} &|E_G(A(G), D(G))| \\ &\leq \max\{1 + 2(s-1) + s(2k-s-3), (2k-2s-1)(s-1) + s^2 + 2, 1 + s(s+1)\} \\ &= 1 + 2(s-1) + s(2k-s-3). \end{aligned}$$

So,

$$|E(G)| \leq \binom{s}{2} + \binom{2k-2s-1}{2} + 1 + 2(s-1) + s(2k-s-3) := f_6(s).$$

We have

$$\begin{aligned} f_6(2) &= \binom{2k-3}{2} + 3, \\ f_6(3) &= \binom{2k-3}{2} - 2k + 12, \\ f_6(k-2) &= \binom{k-2}{2} + (k-2)(k+2) - k + 4 \\ &< \text{ext}(2k, (k-1)K_2) + 2. \end{aligned}$$




Figure 7: G is isomorphic to one of the above three graphs.

If $s = 2$ and $|E(G)| = f_6(2) = \binom{2k-3}{2} + 3$, then it is easy to check that G has a structure shown in Figure 7. By the proof Lemma 3.4, we can obtain a TMC kK_2 in K_{2k} .

If $3 \leq s \leq k - 2$, then $k \geq 5$. If $5 \leq k \leq 6$, then $f_6(k - 2) = f_6(3)$ and $|E(G)| \leq f_6(k - 2) < \text{ext}(2k, (k - 1)K_2) + 2$, which contradicts $|E(G)| = \text{ext}(2k, (k - 1)K_2) + 2$. If $k \geq 7$, then $f_6(3) > f_6(k - 2)$ and $|E(G)| \leq f_6(3) = \binom{2k-3}{2} - 2k + 12 < \binom{2k-3}{2} = \text{ext}(2k, (k - 1)K_2)$, which contradicts $|E(G)| = \text{ext}(2k, (k - 1)K_2) + 3$.

Subcase 4.2. $s = 0$.

In this subcase, $|V(D_1)| = 2k - 1$ and $q = 2$. We suppose that $z_1 \in D_1$ and $D_2 = \{z_2\}$. Let M be a perfect matching of $D_1 - z_1$. Then there exists an edge $e \in M$ such that $c(e) = c(z_1 z_2)$. So, there is no TMC $(k - 1)K_2$ in $D_1 - z_1 - e$. Let D'_1 be $D_1 - z_1 - e$ and $D(D'_1)$, $A(D'_1)$ and $C(D'_1)$ as the canonical decomposition of D'_1 . We look at the graph $G_1 = G - e + z_1 z_2$. Let $A'(G_1) = A(D'_1) \cup z_1$ and $D'(G_1) = D(D'_1) \cup z_2$ and $C'(G_1) = C(D'_1)$. Let $|A'(G_1)| = s'$, $q' = c(D'(G_1)) = c(D(D'_1)) + 1 = (2k - 2) - 2(k - 2) + s - 1 + 1 = s + 2$. Obviously, $1 \leq s' \leq k - 1$. Employing similar technique as in the proofs of Cases 1, 2 and Subcase 4.1, we can obtain contradictions. The details are omitted. Up to now, the proof is complete. \blacksquare

Acknowledgment: The authors are very grateful to the referees for their helpful comments and suggestions.

References

- [1] N. Alon, On a conjecture of Erdős, Simonovits and Sós concerning anti-Ramsey theorems, *J. Graph Theory* **7**(1983), 91-94.
- [2] M. Axenovich and A. Kündgen, On a generalized anti-Ramsey problem, *Combinatorica* **21**(2001), 335-349.

- [3] J.A. Bondy and U.S.R. Murty, “Graph Theory with Applications”, Macmillan, London; Elsevier, New York, 1976.
- [4] P. Erdős and T. Gallai, On maximal paths and circuits of graphs, *Acta Math. Acad. Sc. Hungar.* **10**(1959), 337-356.
- [5] P. Erdős, M. Simonovits and V.T. Sós, Anti-Ramsey theorems, in: A. Hajnal, R. Rado and V.T. Sós (Eds), Infinite and Finite Sets, Vol.II, Colloq. Math. Soc. János Bolyai **10**(1975), 633-643.
- [6] T. Jiang, Edge-colorings with no large polychromatic stars, *Graphs and Combin.* **18**(2002), 303-308.
- [7] T. Jiang and D.B. West, Edge-colorings of complete graphs that avoid polychromatic trees, *Discrete Math.* **274**(2004), 137-145.
- [8] L. Lovász and M.D. Plummer, “Matching Theory”, North-Holland-Amsterdam, New York, Oxford, Tokyo, 1986.
- [9] J.J. Montellano-Ballesteros and V. Neumann-Lara, An anti-Ramsey theorem on cycles, *Graphs and Combin.* **21**(2005), 343C354.
- [10] I. Schiermeyer, Rainbow numbers for matchings and complete graphs, *Discrete Math.* **286**(2004), 157-162.
- [11] M. Simonovits and V.T. Sós, On restricted colourings of K_n , *Combinatorica* **4**(1984), 101-110.