

FULLY COMMUTATIVE ELEMENTS AND KAZHDAN-LUSZTIG CELLS IN THE FINITE AND AFFINE COXETER GROUPS, II

JIAN-YI SHI

ABSTRACT. Let W be an irreducible finite or affine Coxeter group and let W_c be the set of fully commutative elements in W . We prove that the set W_c is closed under the Kazhdan-Lusztig's preorder $\underset{LR}{\geq}$ if and only if W_c is a union of two-sided cells of W .

Introduction.

Let $W = (W, S)$ be a Coxeter group with S the distinguished generator set. For any $J = \{s_1, \dots, s_r\} \subseteq S$, denote by w_J or $w_{s_1 s_2 \dots s_r}$ the longest element in the subgroup W_J of W generated by J . The fully commutative elements of W were defined by Stembridge: $w \in W$ is *fully commutative*, if any two reduced expressions of w can be transformed from each other by only applying the relations $st = ts$ with $s, t \in S$ and $o(st) = 2$ ($o(st)$ being the order of st), or equivalently, w has no reduced expression of the form $w = xw_{st}y$ with $o(st) > 2$ for some $s \neq t$ in S (see [17, Proposition 2.1]). The fully commutative elements were studied extensively by a number of people (see [2], [4], [6], [7], [16], [17]). Let W_c be the set of all the fully commutative elements in W .

In the present paper, we only consider (and always assume) the case where W is an irreducible finite or affine Coxeter group unless otherwise specified. The paper is a continuation of my previous paper [16]; the latter proved that the set W_c is a union of

Supported by Nankai Univ., the 973 Project of MST of China, the NSF of China, the SF of the Univ. Doctoral Program of ME of China, the Shanghai Priority Academic Discipline, and the CST of Shanghai

two-sided cells (in the sense of Kazhdan-Lusztig, see [8]) if and only if W has a non-branching Coxeter graph and is not \tilde{F}_4 . The aim of this paper is to give a necessary and sufficient condition for the set W_c being closed under the Kazhdan–Lusztig preorder $\underset{LR}{\geq}$ (see Theorem 2.1). We use the result of [16] mentioned above and the following key observation: If W has a non-branching Coxeter graph and is not \tilde{F}_4 , then for any $w \notin W_c$, there exists some $y \in M(w)$ (see 1.5 for the notation) such that $\mathcal{L}(y)$ is not fully commutative (see 1.1). Then we get our result by comparing the generalized τ -invariants on the elements in the set W_c and in its complement $W \setminus W_c$ (see [12, Section 4]).

In [7, Section 3.1], Green and Losonczy proved that an irreducible finite Coxeter group W contains no subgraph of type D_4 in its Coxeter graph if and only if the set W_c is closed under $\underset{LR}{\geq}$ and is a union of two-sided cells. They gave a conceptual (resp., a computer) proof for $W = B_m, A_n, m \geq 2, n \geq 1$ (resp., $W = F_4, H_3, H_4$) and referred the proof for the other cases to the papers [3], [5]. Then in [6, Theorem 3.4], Green proved that W is a union of two-sided cells closed under $\underset{LR}{\geq}$ for $W = \tilde{A}_n, n \geq 1$. The results [6, Theorem 3.4] and [7, Section 3.1] on $\tilde{A}_n, A_n, n \geq 1$, may also be obtained from my earlier results [11, Theorem 17.4], [13, Theorem 3.1] and [14, Section 2.9] by [17, Theorem 2.1].

In the proof of our main result (i.e., Theorem 2.1), we use the right cell graphs, rather than a computer, in dealing with the cases of $W = \tilde{G}_2, F_4, H_3, H_4$ (see Appendix and the proof of Lemma 2.2).

The contents of the paper are organized as follows. We collect some notations, terminology and known results concerning Kazhdan–Lusztig cells of a Coxeter group W in Section 1. Then we prove our main result in Section 2. In Appendix we list some right cell graphs in $W \setminus W_c$ for $W = \tilde{G}_2, F_4, H_4, H_3$, which are used in the proof of Lemma 2.2.

§1. Some results on Coxeter groups.

Let (W, S) be a Coxeter system. In the Introduction we defined the set W_c of all the fully commutative elements of W . In this section, we collect some notations, terminology and known results for later use.

1.1. Let \leq be the Bruhat–Chevalley order and $\ell(w)$ the length function on W . Call a subset J of S *fully commutative* if the element w_J is so.

For $w, x, y \in W$, we use the notation $w = x \cdot y$ to mean $w = xy$ and $\ell(w) = \ell(x) + \ell(y)$. If $w = x \cdot y \in W_c$ then $x, y \in W_c$. In particular, if $w \in W_c$ has an expression $w = x \cdot w_J \cdot y$ with $x, y \in W$ and $J \subseteq S$, then J is fully commutative.

1.2. Let \leq_L (resp., \leq_R , \leq_{LR}) be the preorder on W defined as in [8], and let \sim_L (resp., \sim_R , \sim_{LR}) be the equivalence relation on W determined by \leq_L (resp., \leq_R , \leq_{LR}). The corresponding equivalence classes are called *left* (resp., *right*, *two-sided*) *cells* of W . The preorder \leq_L (resp., \leq_R , \leq_{LR}) on W induces a partial order on the set of left (resp., right, two-sided) cells of W .

1.3. For any $w \in W$, let $\mathcal{L}(w) = \{s \in S \mid sw < w\}$ and $\mathcal{R}(w) = \{s \in S \mid ws < w\}$.

Assume $m = o(st) > 2$ for some $s, t \in S$. A sequence of elements

$$\underbrace{sy, tsy, stsy, \dots}_{m-1 \text{ terms}}$$

is called a *left $\{s, t\}$ -string* if $y \in W$ satisfies $\mathcal{L}(y) \cap \{s, t\} = \emptyset$.

We say that z is obtained from w by a *left $\{s, t\}$ -star operation*, if z, w are two neighboring terms in a left $\{s, t\}$ -string. Clearly, a resulting element z of a left $\{s, t\}$ -star operation on w , when it exists, need not be unique unless w is a terminal term of the left $\{s, t\}$ -string containing it.

The following result follows directly from the definition of the relation \sim_L on W .

Lemma. *If $x, y \in W$ can be obtained from each other by successively applying left star operations, then $x \sim_L y$.*

1.4. By the notation $x \text{---} y$ in W , we mean that either $x < y$ or $y < x$ holds and that $\max\{\deg P_{x,y}, \deg P_{y,x}\} = \frac{1}{2}(|\ell(x) - \ell(y)| - 1)$, where $P_{x,y}$ is the celebrated Kazhdan–Lusztig polynomial associated to $x, y \in W$ (see [8, Theorem 1.1]).

(a) The relation $x \underset{L}{\leq} y$ (resp., $x \underset{R}{\leq} y$) implies $\mathcal{R}(x) \supseteq \mathcal{R}(y)$ (resp., $\mathcal{L}(x) \supseteq \mathcal{L}(y)$). In particular, the relation $x \underset{L}{\sim} y$ (resp., $x \underset{R}{\sim} y$) implies $\mathcal{R}(x) = \mathcal{R}(y)$ (resp., $\mathcal{L}(x) = \mathcal{L}(y)$) (see [8, Proposition 2.4]). Hence it makes sense to write $\mathcal{L}(\Gamma)$ (resp., $\mathcal{R}(\Gamma)$) for any right (resp., left) cell Γ of W , where $\mathcal{L}(\Gamma) = \mathcal{L}(z)$ (resp., $\mathcal{R}(\Gamma) = \mathcal{R}(z)$) for any $z \in \Gamma$.

(b) If $x, y \in W$ with $x \text{---} y$ are in some left $\{s, t\}$ -strings (not necessarily in the same left string; see 1.3) for some $s, t \in S$ with $st \neq ts$, then there exist some $x', y' \in W$ which are obtained from x, y respectively by a left $\{s, t\}$ -star operation and satisfy $x' \text{---} y'$ (see [9, Section 10.4]).

(c) $x \underset{LR}{\sim} x^{-1}$ for any $x \in W$ (see [10, Corollary 1.9 (a) and Theorem 1.10] and [1, Corollary 3.2]).

1.5. For any $w \in W$, let $M(w)$ be the set of all the elements y satisfying: there exists a sequence of elements $z_0 = w, z_1, \dots, z_t = y$ in W with $t \geq 0$ such that z_i is obtained from z_{i-1} by a left star operation for every $1 \leq i \leq t$. We see by Lemma 1.3 that all the elements in $M(w)$ are in the same left cell of W .

§2. The condition for W_c being closed under the preorder $\underset{LR}{\geq}$.

In this section, assume that W is an irreducible finite or affine Coxeter group. In [16, Theorem 3.4 and Sections 3.5–3.7], we showed that the set W_c is a union of two-sided cells of W if and only if W has a non-branching Coxeter graph and is not \tilde{F}_4 . We understand that this result was already known in the case where W is any irreducible finite Coxeter group (see [7]).

A subset K of W is *closed under the preorder* $\underset{LR}{\geq}$ if the conditions $x \in K, y \in W$ and $y \underset{LR}{\geq} x$ together imply $y \in K$.

In the present section, we want to give a necessary and sufficient condition for the set W_c to be closed under $\underset{LR}{\geq}$.

Theorem 2.1. *Let W be an irreducible finite or affine Coxeter group. Then W_c is closed under $\underset{LR}{\geq}$ if and only if W_c is a union of two-sided cells of W .*

To prove Theorem 2.1, we need prove some lemmas.

Lemma 2.2. *If W is an irreducible finite or affine Coxeter group such that W_c is a union of two-sided cells of W , then for any $w \in W \setminus W_c$, there exists some $y \in M(w)$ (see 1.5) such that $\mathcal{L}(y)$ is not fully commutative (see 1.1).*

Proof. By [16, Theorem 3.4 and 3.5–3.7], we know that W_c is a union of two-sided cells of W if and only if W has a non-branching Coxeter graph and is not \tilde{F}_4 , i.e., W is one of the following groups: $A_n, \tilde{A}_n, I_2(m), \tilde{C}_l, B_l, F_4, H_3, H_4, \tilde{G}_2$, where $n \geq 1, m \geq 5$ and $l \geq 2$. The result follows by [11, Theorems 17.4, 17.6 and Propositions 9.3.7, 16.2.4] for the groups \tilde{A}_n and A_n , and by [16, Corollary 3.3] for the groups \tilde{C}_l . By the fact that B_l is a standard parabolic subgroup of \tilde{C}_l , we can show the result for the groups B_l by the same argument as that for [16, Corollary 3.3]. Then the result for the groups F_4, H_3, H_4 and \tilde{G}_2 can be checked directly from their right cell graphs (see Appendix). Finally, the result for the groups $I_2(m)$ is obvious. \square

Remark 2.3. It is necessary for the assumption that W_c is a union of two-sided cells of W in Lemma 2.2. There is a counter-example when such a condition is removed. Let $W = \tilde{F}_4$ and $S = \{s_0, s_1, s_2, s_3, s_4\}$ be with $o(s_0s_1) = o(s_1s_2) = o(s_3s_4) = 3$ and $o(s_2s_3) = 4$. Then the element $w = s_4s_2s_3s_2s_0s_1s_0$ is not fully commutative. However, $\mathcal{L}(y)$ is fully commutative for any element y in $M(w)$ (see [12, Section 5.4]).

By Lemma 2.2, we can prove the following

Lemma 2.4. *When it is a union of two-sided cells of W , the set W_c is closed under the*

$preorder \underset{LR}{\geq}$.

Proof. Suppose not. Then there exist some $x \in W_c$ and some $w \in W \setminus W_c$ with $x \underset{L}{\leq} w$. We may assume $x \text{---} w$ and $\mathcal{L}(x) \not\subseteq \mathcal{L}(w)$ without loss of generality. So $\mathcal{R}(x) \supseteq \mathcal{R}(w)$ by 1.4 (a). Hence $\mathcal{L}(x^{-1}) \supseteq \mathcal{L}(w^{-1})$. By Lemma 2.2, there exists an element y in $M(w^{-1})$ with $\mathcal{L}(y)$ not fully commutative. Then there exists a sequence of elements $w_0 = w^{-1}, w_1, \dots, w_r = y$ in $M(w^{-1})$ such that w_i is obtained from w_{i-1} by a left $\{s_i, t_i\}$ -star operation for every $1 \leq i \leq r$ and some $s_i, t_i \in S$ with $s_i t_i \neq t_i s_i$. We may assume r minimal with this property. Hence the $\mathcal{L}(w_i)$'s, $0 \leq i < r$, are all fully commutative. Since w_1 is obtained from w^{-1} by a left $\{s_1, t_1\}$ -star operation, we have $|\{s_1, t_1\} \cap \mathcal{L}(w^{-1})| = 1$. Since $\mathcal{L}(x^{-1})$ is fully commutative and $\mathcal{L}(x^{-1}) \supseteq \mathcal{L}(w^{-1})$, we have $|\{s_1, t_1\} \cap \mathcal{L}(x^{-1})| = 1$ also. So we can apply a left $\{s_1, t_1\}$ -star operation on x^{-1} to obtain some element x_1 in $M(x^{-1})$ with $x_1 \text{---} w_1$ by 1.4 (b). Since $\mathcal{R}(x_1) = \mathcal{R}(x^{-1}) = \mathcal{L}(x) \not\subseteq \mathcal{L}(w) = \mathcal{R}(w^{-1}) = \mathcal{R}(w_1)$, we have $x_1 \underset{R}{\leq} w_1$ and hence $\mathcal{L}(x_1) \supseteq \mathcal{L}(w_1)$ by 1.4 (a). When $r > 1$, we can apply a left $\{s_2, t_2\}$ -star operation on x_1 to obtain some element x_2 with $x_2 \text{---} w_2$ by the same reason as that for getting x_1 from x^{-1} . Continuing this process, we get a sequence of elements $x_0 = x^{-1}, x_1, \dots, x_r$ in $M(x^{-1})$ such that x_i is obtained from x_{i-1} by a left $\{s_i, t_i\}$ -star operation and $x_i \text{---} w_i$ for $1 \leq i \leq r$. By the assumption that W_c is a union of two-sided cells of W and by the facts that $x_r \underset{L}{\sim} x^{-1} \underset{LR}{\sim} x$ (by 1.4 (c)) and $x \in W_c$, we have $x_r \in W_c$ and hence the set $\mathcal{L}(x_r)$ is fully commutative. Since $\mathcal{L}(w_r)$ is not fully commutative, we have $\mathcal{L}(w_r) \not\subseteq \mathcal{L}(x_r)$. Since $x_r \text{---} w_r$, this implies $w_r \underset{L}{\leq} x_r$ and hence $x \underset{L}{\leq} w \underset{LR}{\sim} w^{-1} \underset{LR}{\sim} w_r \underset{L}{\leq} x_r \underset{L}{\sim} x^{-1} \underset{LR}{\sim} x$ by 1.4 (c). We get $x \underset{LR}{\sim} w$, contradicting the assumption that W_c is a union of two-sided cells of W . So our result follows. \square

2.5. Proof of Theorem 2.1. The implication “ \Leftarrow ” is just Lemma 2.4. For the implication “ \Rightarrow ”, we need only show that $x \not\underset{LR}{\sim} y$ for any $x \in W_c$ and any $y \in W \setminus W_c$.

Suppose not. Then there exist some $x \in W_c$ and some $y \in W \setminus W_c$ with $x \underset{LR}{\sim} y$ (and hence $y \underset{LR}{\geq} x$). But this would imply $y \in W_c$ by the assumption that W_c is closed under $\underset{LR}{\geq}$, a contradiction. So Theorem 2.1 follows. \square

Appendix.

A *right cell graph associated to an element $x \in W$* (written $\mathfrak{M}_R(x)$) is by definition a graph whose vertex set $V(x)$ consists of all the right cells Γ of W with $\Gamma \cap M(x) \neq \emptyset$ (each right cell is represented by a box). Two vertices Γ, Γ' of $\mathfrak{M}_R(x)$ are joined by an edge, if there are some $y \in M(x) \cap \Gamma$ and $z \in M(x) \cap \Gamma'$ such that y, z are two neighboring terms of a left string. Each vertex Γ of $\mathfrak{M}_R(x)$ is labelled by the set $\mathcal{L}(\Gamma)$ (see 1.4 (a)).


It is easily seen that the set of the subsets of S occurring as the labels of the vertices in $\mathfrak{M}_R(x)$ is equal to the set $\{I \subseteq S \mid I = \mathcal{L}(y) \text{ for some } y \in M(x)\}$.

Two right cell graphs $\mathfrak{M}_R(x)$ and $\mathfrak{M}_R(y)$ are *isomorphic* if there exists a bijection $\phi : V(x) \rightarrow V(y)$ such that $\mathcal{L}(\Gamma) = \mathcal{L}(\phi(\Gamma))$ for any $\Gamma \in V(x)$ and such that any pair $\Gamma, \Gamma' \in V(x)$ are joined by an edge if and only if $\phi(\Gamma), \phi(\Gamma')$ are so.

Note that the definition of a right cell graph imitates that of a left cell graph, the latter was given in my previous paper [15, Subsection 2.11].

We work out all the right cell graphs in $W \setminus W_c$ (resp., a representative set of the isomorphism classes of those graphs) for the groups $W = \tilde{G}_2, F_4$ (resp., H_4, H_3) according to the results in [9], [18], [1].


(1) $W = \tilde{G}_2$ with $S = \{s_0, s_1, s_2\}$ satisfying $o(s_0s_2) = 3$ and $o(s_1s_2) = 6$:






Here and later the boldfaced numbers in a box Γ represent the elements in $\mathcal{L}(\Gamma)$. The box

of $\mathfrak{M}_R(x)$ with inside numbers underlined represents the right cell Γ_x containing x . For example, the box $\underline{02}$ in $\mathfrak{M}_R(s_0s_2s_1s_2s_0w_{12})$ represents the right cell $\Gamma = \Gamma_{s_0s_2s_1s_2s_0w_{12}}$ with $\mathcal{L}(\Gamma) = \{s_0, s_2\}$; while two boxes $\underline{01}$ in $\mathfrak{M}_R(w_{12})$ represent respectively two right cells $\Gamma, \Gamma' \in V(w_{12})$ with $\mathcal{L}(\Gamma) = \mathcal{L}(\Gamma') = \{s_0, s_1\}$. The notation $w_{ij\dots}$ stands for the element $w_{s_i s_j \dots}$ (see the first paragraph in Introduction)


(2) $W = F_4$ with $S = \{s_1, s_2, s_3, s_4\}$ satisfying $o(s_1s_2) = o(s_3s_4) = 3$ and $o(s_2s_3) = 4$.



(3) $W = H_4$ with $S = \{s_1, s_2, s_3, s_4\}$ satisfying $o(s_1s_2) = o(s_2s_3) = 3$ and $o(s_3s_4) = 5$.



(4) $W = H_3$ with $S = \{s_1, s_2, s_3\}$ satisfying $o(s_1 s_2) = 3$ and $o(s_2 s_3) = 5$.



Acknowledgement. I would like to take this opportunity to express my deep gratitude to Professor Y. C. Chen who invited me to visit Nankai University and to the Center for Combinatorics, Nankai University for hospitality during the writing of this paper.

REFERENCES

1. D. Alvis, *The left cells of the Coxeter group of type H_4* , J. Algebra **107** (1987), 160–168.
2. S. C. Billey and G. S. Warrington, *Kazhdan–Lusztig polynomials for 321-hexagon-avoiding permutations*, J. Algebraic Combin. **13** (2001), 111–136.
3. C. K. Fan, *A Hecke algebra quotient and properties of commutative elements of a Weyl group*, Ph.D. thesis, M.I.T., 1995.
4. C. K. Fan and J. R. Stembridge, *Nilpotent orbits and commutative elements*, J. Algebra **196** (1997), 490–498.
5. J.J.Graham, *Modular representations of Hecke algebras and related algebras*, Ph.D. thesis, Univ. of Sydney, 1995.
6. R.M.Green, *On 321-avoiding permutations in affine Weyl groups*, J. Alg. Combin. **15** (2002), 241–252.
7. R. M. Green and J. Losonczy, *Fully commutative Kazhdan–Lusztig cells*, Ann. Inst. Fourier (Grenoble) **51** (2001), 1025–1045.
8. D. Kazhdan and G. Lusztig, *Representations of Coxeter groups and Hecke algebras*, Invent. Math. **53** (1979), 165–184.
9. G. Lusztig, *Cells in affine Weyl groups*, in “Algebraic Groups and Related Topics”(R. Hotta, ed.), Advanced Studies in Pure Math., Kinokuniya and North Holland,(1985), 255–287.
10. G. Lusztig, *Cells in affine Weyl groups, II*, J. Algebra **109** (1987), 536–548.
11. Jian-yi Shi, *The Kazhdan–Lusztig cells in certain affine Weyl groups*, vol. 1179, Springer–Verlag, Lecture Notes in Mathematics, 1986.
12. Jian-yi Shi, *Left cells in affine Weyl groups*, Tôhoku Math. J. **46** (1994), 105–124.
13. Jian-yi Shi, *Some results relating two presentations of certain affine Weyl groups*, J. Algebra **163**(1) (1994), 235–257.
14. Jian-yi Shi, *The partial order on two-sided cells of certain affine Weyl groups*, J. Algebra. **179**(2) (1996), 607–621.
15. Jian-yi Shi, *Left cells in the affine Weyl group of type \tilde{F}_4* , J. Algebra **200** (1998), 173–206.
16. Jian-yi Shi, *Fully commutative elements and Kazhdan–Lusztig cells in the finite and affine Coxeter groups*, Proc. of AMS **131** (2003), 3371–3378.
17. J. R. Stembridge, *On the fully commutative elements of Coxeter groups*, J. Algebraic Combin. **5** (1996), 353–385.
18. K. Takahashi, *The left cells and their W -graphs of Weyl group of type F_4* , Tokyo J. Math. **13** (1990), 327–340.

Center for Combinatorics,
 The Key Laboratory of Pure Mathematics and
 Combinatorics of Ministry of Education,
 Nankai University,
 Tianjin, 300071,
 P. R. China

Department of Mathematics,
 East China Normal University,
 Shanghai, 200062,
 P. R. China