

A planar network proof for Hankel total positivity of type B Narayana polynomials

Ethan Y.H. Li¹, Grace M.X. Li², Arthur L.B. Yang³ and Candice X.T. Zhang⁴

Center for Combinatorics, LPMC
Nankai University, Tianjin 300071, P. R. China

Email: ¹yinhao_li@mail.nankai.edu.cn, ²limengxing@mail.nankai.edu.cn,
³yang@nankai.edu.cn, ⁴zhang_xutong@mail.nankai.edu.cn

Abstract. The Hankel matrix of type B Narayana polynomials was proved to be totally positive by Wang and Zhu, and independently by Sokal. Pan and Zeng raised the problem of giving a planar network proof of this result. In this paper, we present such a proof by constructing a planar network allowing negative weights, applying the Lindström-Gessel-Viennot lemma and establishing an involution on the set of nonintersecting families of directed paths.

AMS Mathematics Subject Classification 2020: 05A15, 05A19, 05A20

Keywords: Narayana polynomials of type B , Hankel total positivity, planar network, coefficient matrix, the Lindström-Gessel-Viennot lemma

1 Introduction

For any $n \geq 0$, let

$$W_n(q) = \sum_{k=0}^n \binom{n}{k}^2 q^k$$

denote the n -th Narayana polynomial of type B . Wang and Zhu [25], and Sokal [22] independently proved that the Hankel matrix

$$H = (W_{i+j}(q))_{i,j \geq 0} \tag{1.1}$$

is q -totally positive, namely, any minor of H is a polynomial in q with nonnegative coefficients. The main objective of this paper is to give a combinatorial proof of the q -total positivity of H , which solves a problem of Pan and Zeng [18].

The q -total positivity of the Hankel matrix H arose in the study of the q -log-convexity of the polynomial sequence $(W_n(q))_{n \geq 0}$. For the convenience of introducing related definitions and results, we make use of the notion of q -nonnegativity and the symbol \geq_q . A polynomial $f(q)$ with real coefficients is called q -nonnegative, written $f(q) \geq_q 0$, if all

its coefficients are nonnegative. Accordingly, for two polynomials $f(q)$ and $g(q)$ we write $f(q) \geq_q g(q)$ if $f(q) - g(q) \geq_q 0$. Recall that a sequence $\alpha = (a_n(q))_{n \geq 0}$ of polynomials in q is said to be q -log-convex if for any $n \geq 1$ there holds $a_{n+1}(q)a_{n-1}(q) \geq_q a_n^2(q)$. Furthermore, if $a_{m+1}(q)a_{n-1}(q) \geq_q a_m(q)a_n(q)$ holds for any $m \geq n \geq 1$, then we call α a strongly q -log-convex sequence. Conversely, we say that α is a q -log-concave sequence if for any $n \geq 1$ we have $a_n^2(q) \geq_q a_{n+1}(q)a_{n-1}(q)$, and it is a strongly q -log-concave sequence if $a_m(q)a_n(q) \geq_q a_{m+1}(q)a_{n-1}(q)$ holds for any $m \geq n \geq 1$. The concept of q -log-concavity was introduced by Stanley, and the notion of strong q -log-concavity was due to Sagan [19]. Many polynomial sequences have been proved to be q -log-concave, or even strongly q -log-concave, see Butler [2], Krattenthaler [10], Leroux [12], Sagan [19, 20], and Chen, Wang and Yang [5]. However, q -log-convex sequences received very little attention until the work of Liu and Wang [16], who first introduced the notion of q -log-convexity. Liu and Wang established the q -log-convexity of many combinatorial polynomials, such as the Eulerian polynomials. For further progress on q -log-convexity, see [4, 27] for instance.

The q -log-convexity of $(W_n(q))_{n \geq 0}$ was conjectured by Liu and Wang [16], and was proved later by Chen, Tang, Wang and Yang [3] by using the theory of symmetric functions. Zhu [26] further established the strong q -log-convexity of $(W_n(q))_{n \geq 0}$ by identifying this polynomial sequence as the first column of the triangular array $B = (b_{n,k}(q))_{n,k \geq 0}$, which is generated by

$$\begin{aligned} b_{n,0}(q) &= (1+q) \cdot b_{n-1,0}(q) + 2q \cdot b_{n-1,1}(q); \\ b_{n,k}(q) &= b_{n-1,k-1}(q) + (1+q) \cdot b_{n-1,k}(q) + q \cdot b_{n-1,k+1}(q) \quad (k \geq 1, n \geq 1) \end{aligned} \tag{1.2}$$

with $b_{0,0}(q) = 1$ and $b_{n,k}(q) = 0$ for $k > n$. The triangular array B belongs to a wide class of matrices, called q -recursive matrices in [25], or Catalan-Stieltjes matrices in [18, 13], which we will recall below. Let $\gamma = (r_k(q))_{k \geq 0}$, $\sigma = (s_k(q))_{k \geq 0}$ and $\tau = (t_k(q))_{k \geq 1}$ be three sequences of polynomials in q . The Catalan-Stieltjes matrix with respect to γ, σ, τ , denoted by $C^{\gamma, \sigma, \tau} = (c_{n,k}(q))_{n,k \geq 0}$, is generated by the following recursive relations:

$$\begin{aligned} c_{n,0}(q) &= s_0(q)c_{n-1,0}(q) + t_1(q)c_{n-1,1}(q); \\ c_{n,k}(q) &= r_{k-1}(q)c_{n-1,k-1}(q) + s_k(q)c_{n-1,k}(q) + t_{k+1}(q)c_{n-1,k+1}(q) \quad (k \geq 1, n \geq 1), \end{aligned}$$

where $c_{0,0}(q) = 1$ and $c_{n,k}(q) = 0$ unless $n \geq k \geq 0$. Actually, Zhu [26] gave a general criterion for the strong q -log-convexity of $(c_{n,0}(q))_{n \geq 0}$ of $C^{\gamma, \sigma, \tau}$. Further, Wang and Zhu [25] proved that the Hankel matrix $(c_{i+j,0}(q))_{i,j \geq 0}$ is q -totally positive provided that the

matrix

$$\begin{pmatrix} 1 & & & & \\ s_0(q) & r_0(q) & & & \\ t_1(q) & s_1(q) & r_1(q) & & \\ & t_2(q) & s_2(q) & r_2(q) & \\ & & \ddots & \ddots & \ddots \end{pmatrix},$$

called the coefficient matrix of $C^{\gamma,\sigma,\tau}$, is q -totally positive. Let L^B denote the corresponding coefficient matrix of B . Wang and Zhu first proved the q -total positivity of L^B and hence obtained the q -total positivity of the Hankel matrix $H = (W_{i+j}(q))_{i,j \geq 0}$. The latter result was also independently proved by Sokal [22] based on the continued fraction expression of the generating function $\sum_{n \geq 0} W_n(q)x^n$.

We would like to note that the q -total positivity of the Hankel matrix $(c_{i+j,0}(q))_{i,j \geq 0}$ is also closely related to that of $C^{\gamma,\sigma,\tau}$, for details see [14] and [25]. Chen, Liang and Wang [6] raised the problem of giving a combinatorial interpretation for the q -total positivity of $C^{\gamma,\sigma,\tau}$. An ideal tool to combinatorially proving the positivity of a matrix is the famous Lindström-Gessel-Viennot lemma, see [15, 7, 8]. A natural strategy is to construct a planar network with nonnegative weights for the target matrix, and then to apply the Lindström-Gessel-Viennot lemma to interpret each minor of this matrix as the generating function of nonintersecting families of directed paths, which are obviously nonnegative. In the spirit of this method, Pan and Zeng [18] provided a general planar network construction for the Catalan-Stieltjes matrices and their associated Hankel matrices, which enables them to give combinatorial proofs of the q -total positivity for many such matrices, such as those related to the Eulerian polynomials, Schröder polynomials, and Narayana polynomials of type A . However, their approach did not work for Narayana polynomials of type B , and they proposed it as an open problem to find a planar network proof of the q -total positivity of $H = (W_{i+j}(q))_{i,j \geq 0}$. It seems impossible to find a planar network with only nonnegative weights for H .

In this paper, inspired by our recent work [13], we construct for H a suitable planar network allowing negative weights and solve Pan and Zeng's problem. In our construction, the planar network for H can be naturally divided into serial segments which are essentially subnetworks of the planar network for the coefficient matrix L^B of B . By applying the Lindström-Gessel-Viennot lemma and establishing a sign-reversing involution on the nonintersecting families of each segment, we combinatorially prove the q -total positivity of L^B and H .

This paper is organized as follows. In Section 2, we will introduce the Lindström-Gessel-Viennot lemma. In Section 3, we will present our planar network construction for

the coefficient matrix L^B , as well as a combinatorial proof of its q -total positivity. In Section 4 we will make use of the results in Section 3 to obtain a planar network for H and a combinatorial proof of its q -total positivity. We conclude this paper in Section 5 with a conjecture on the immanant positivity for H .

2 The Lindström-Gessel-Viennot lemma

The Lindström-Gessel-Viennot lemma was originally proved by Lindström [15] and further developed by Gessel and Viennot [7, 8]. It has a broad range of applications, see [9, 11, 17, 24] for instance. In this section, we will give an overview of the Lindström-Gessel-Viennot lemma, which plays a key role in our combinatorial proof of the q -total positivity of the Hankel matrix of type B Narayana polynomials.

To state the Lindström-Gessel-Viennot lemma, we need some notations. Let D be a directed graph, or digraph for short, with vertex set $V(D)$ and arc set $A(D)$. A digraph D is said to be acyclic if it contains no directed cycles. Throughout this paper we may assume that D is locally finite, namely, for any two vertices $u, v \in V(D)$ the number of directed paths from u to v is finite. We say two directed paths intersect if they have a vertex in common. A sequence (p_1, \dots, p_n) of directed paths is called a nonintersecting family if p_i and p_j do not intersect for any $i \neq j$. Let $\mathbf{U} = (u_1, \dots, u_n)$ and $\mathbf{V} = (v_1, \dots, v_n)$ be two sequences of vertices in D , and let $\mathcal{N}_D(\mathbf{U}, \mathbf{V})$ denote the set of nonintersecting families (p_1, \dots, p_n) such that p_i is a directed path from u_i to v_i for each $1 \leq i \leq n$. If for any permutation σ of $\{1, 2, \dots, n\}$, the set $\mathcal{N}_D(\mathbf{U}, \sigma(\mathbf{V})) = \mathcal{N}_D((u_1, \dots, u_n), (v_{\sigma(1)}, \dots, v_{\sigma(n)}))$ is empty unless σ is the identity permutation, then \mathbf{U} and \mathbf{V} are said to be compatible. A weight function wt of D is a map from $A(D)$ to R , where R is a commutative ring with identity. The weight of a directed path in D is the product of the weights of all its arcs, and the weight of a nonintersecting family is defined to be the product of the weights of all its components. Given two vertices u and v of D , let $GF_{D, \text{wt}}(u, v)$ denote the sum of the weights of all directed paths from u to v . For notational convenience we will use $GF_D(u, v)$ instead of $GF_{D, \text{wt}}(u, v)$ when the weight function is easy to recognize. For two sequences \mathbf{U} and \mathbf{V} of vertices in D , let $GF(\mathcal{N}_D(\mathbf{U}, \mathbf{V}))$ denote the sum of the weights of all elements in $\mathcal{N}_D(\mathbf{U}, \mathbf{V})$. For a matrix M , we denote by $\det[M]$ the determinant of M . The celebrated Lindström-Gessel-Viennot lemma is stated as follows.

Lemma 2.1 ([8, Corollary 2]) *Let D be a locally finite and acyclic digraph with a weight function, and let $\mathbf{U} = (u_1, \dots, u_n)$, $\mathbf{V} = (v_1, \dots, v_n)$ be two compatible sequences*

of vertices in D . Then

$$\det \left[(GF_D(u_i, v_j))_{1 \leq i, j \leq n} \right] = GF(\mathcal{N}_D(\mathbf{U}, \mathbf{V})).$$

In this paper, we mainly apply the above lemma to a special class of digraphs, called planar networks. Recall that a digraph D is said to be planar if it can be embedded in the plane with edges meeting only at endpoints. We call $\mathcal{D} = (D, \text{wt}_D)$ a planar network if D is a locally finite, acyclic, and planar digraph, and wt_D is a weight function of D . Given an $n \times n$ matrix M , then \mathcal{D} is called a planar network for M if there exist two sequences (u_1, \dots, u_n) and (v_1, \dots, v_n) of vertices in D such that

$$M = (GF_D(u_i, v_j))_{1 \leq i, j \leq n}.$$

In the remaining part of this paper, we usually specify the vertices and say that $\mathcal{D} = (D, \text{wt}_D, (u_1, \dots, u_n), (v_1, \dots, v_n))$ is a planar network for M .

3 The coefficient matrix L^B

In this section we will establish the planar network for the coefficient matrix L^B and prove its q -total positivity. By (1.2), we have

$$L^B = (l_{i,j})_{i,j \geq 0} = \begin{pmatrix} 1 & & & & & & \\ 1+q & 1 & & & & & \\ 2q & 1+q & 1 & & & & \\ & q & 1+q & 1 & & & \\ & & q & 1+q & 1 & & \\ & & & q & 1+q & 1 & \\ & & & & \ddots & \ddots & \ddots \end{pmatrix}.$$

Now we give the construction of the planar network for L^B . Let D^{L^B} be the infinite planar digraph with vertex set

$$V(D^{L^B}) = \{P_i \mid i \geq 0\} \cup \{Q_i \mid i \geq 0\} \cup \{P'_i \mid i \geq 0\}$$

and arc set

$$\begin{aligned} A(D^{L^B}) = & \{P_i \rightarrow Q_i \mid i \geq 0\} \cup \{P_i \rightarrow Q_{i-1} \mid i \geq 1\} \\ & \cup \{Q_i \rightarrow P'_i \mid i \geq 0\} \cup \{Q_i \rightarrow P'_{i-1} \mid i \geq 2\} \\ & \cup \{P_1 \rightarrow P'_0, Q_1 \xrightarrow{\mathcal{L}} P'_0, Q_1 \xrightarrow{\mathfrak{A}} P'_0\}, \end{aligned}$$

where the coordinates of vertices are given by $P_i = (0, -i)$, $Q_i = (1, -i)$ and $P'_i = (2, -i)$, and $Q_1 \xrightarrow{\text{L}} P'_0$, $Q_1 \xrightarrow{\text{R}} P'_0$ are multiple arcs from Q_1 to P'_0 with one drawn on the left and the other on the right, respectively, as shown in Figure 3.1. The weight function $\text{wt}_{D^{LB}}$ is defined by

$$\text{wt}_{D^{LB}}(P_1 \rightarrow P'_0) = -1, \quad \text{wt}_{D^{LB}}(P_i \rightarrow Q_{i-1}) = q \quad \text{for } i \geq 1,$$

and $\text{wt}_{D^{LB}}(a) = 1$ for the other arcs a in $A(D^{LB})$. Then we have the following result.

Lemma 3.1 *Let D^{LB} and $\text{wt}_{D^{LB}}$ be defined as above. Then*

$$\mathcal{L}^B = (D^{LB}, \text{wt}_{D^{LB}}, (P_0, P_1, \dots), (P'_0, P'_1, \dots))$$

is a planar network for L^B , or equivalently,

$$L^B = (GF_{D^{LB}}(P_i, P'_j))_{i,j \geq 0}.$$

Proof. By the definitions in Section 2 and the above construction, it is straightforward to verify that

$$l_{i,j} = GF_{D^{LB}}(P_i, P'_j)$$

for $i, j \geq 0$. Then the proof follows. ■

Figure 3.1 provides an illustration of the planar network \mathcal{L}^B , where we only label the weights not equal to 1.




Figure 3.1: The planar network \mathcal{L}^B

The remaining part of this section is devoted to giving a combinatorial proof of the q -total positivity of L^B by using the planar network \mathcal{L}^B . To this end, we define the following three properties of nonintersecting families $\mathbf{p} = (p_1, \dots, p_k)$:

$$(\mathcal{P}_1) \quad p_1 = P_1 \rightarrow P'_0;$$

$$(\mathcal{P}_2) \quad p_1 = P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0;$$

$$(\mathcal{P}_3) \quad \text{there exists } l \ (l \geq 2) \text{ such that } p_m = P_m \rightarrow Q_{m-1} \rightarrow P'_{m-1} \text{ for } 1 \leq m \leq l-1 \text{ and } p_l = P_l \rightarrow Q_l \rightarrow P'_{l-1}.$$

Given a positive integer k and two sequences

$$\begin{aligned} I &= (i_1, \dots, i_k), \text{ where } 0 \leq i_1 < \dots < i_k, \\ J &= (j_1, \dots, j_k), \text{ where } 0 \leq j_1 < \dots < j_k, \end{aligned} \tag{3.1}$$

let

$$\mathbf{P}_I = (P_{i_1}, \dots, P_{i_k}), \quad \mathbf{P}'_J = (P'_{j_1}, \dots, P'_{j_k}), \tag{3.2}$$

and let

$$S_{I,J} = \{\mathbf{p} \in \mathcal{N}_{DL^B}(\mathbf{P}_I, \mathbf{P}'_J) \mid \mathbf{p} \text{ satisfies none of } (\mathcal{P}_1), (\mathcal{P}_2), (\mathcal{P}_3)\}. \tag{3.3}$$

It is clear that each nonintersecting family in $S_{I,J}$ has a q -nonnegative weight. By virtue of this, the following result provides a combinatorial interpretation for the q -total positivity of L^B .

Theorem 3.2 *Let $I, J, \mathbf{P}_I, \mathbf{P}'_J, S_{I,J}$ be as given in (3.1), (3.2) and (3.3) respectively, and let $L^B_{I,J}$ denote the submatrix of L^B whose rows are indexed by I and columns indexed by J . Then we have*

$$\det [L^B_{I,J}] = GF(S_{I,J}), \tag{3.4}$$

where $GF(S_{I,J})$ denotes the sum of weights of all elements in $S_{I,J}$. In particular, $\det [L^B_{I,J}]$ is q -nonnegative.

Proof. By Lemma 3.1 and Lemma 2.1, we have

$$\det [L^B_{I,J}] = GF(\mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J)),$$

where we use $\mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J)$ to stand for $\mathcal{N}_{DL^B}(\mathbf{P}_I, \mathbf{P}'_J)$ for convenience. In the following we may assume that $\mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J) \neq \emptyset$, otherwise, $S_{I,J} = \emptyset$, $\det [L^B_{I,J}] = 0$, and (3.4) holds trivially. For $i = 1, 2, 3$, let

$$\mathcal{N}_i = \{\mathbf{p} \in \mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J) \mid \mathbf{p} \text{ satisfies } (\mathcal{P}_i)\}.$$

Clearly, $\mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J)$ is the disjoint union of $\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_3$ and $S_{I,J}$. Now it suffices to give an involution ϕ on $\mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J)$ such that any nonintersecting family $\mathbf{p} \in \mathcal{N}_1 \cup \mathcal{N}_2 \cup \mathcal{N}_3$ and its image $\phi(\mathbf{p})$ have opposite weights, and the restriction of ϕ to $S_{I,J}$ is the identity map. Thus, we only need to define the action of ϕ on $\mathcal{N}_1 \cup \mathcal{N}_2 \cup \mathcal{N}_3$.

Let us first consider the case $k = 1$, for which we have $\mathcal{N}_3 = \emptyset$. If $i_1 = 1, j_1 = 0$, then we have

$$\mathcal{N}_1 = \{P_1 \rightarrow P'_0\}, \quad \mathcal{N}_2 = \{P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0\}.$$

Define ϕ to be the map which sends $P_1 \rightarrow P'_0$ and $P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0$ to each other. Note that the weight of $P_1 \rightarrow P'_0$ is -1 , while the weight of $P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0$ is 1 . This establishes the desired involution. If $i_1 \neq 1$ or $j_1 \neq 0$, then $\mathcal{N}_1 = \mathcal{N}_2 = \mathcal{N}_3 = \emptyset$ and hence $\mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J) = S_{I,J}$; for this subcase, we simply take ϕ to be the identity map.

We proceed to define ϕ for $k \geq 2$. In this case we divide \mathcal{N}_1 into the following two subsets:

$$\mathcal{N}_{1,1} = \{\mathbf{p} \in \mathcal{N}_1 \mid p_2 = P_2 \rightarrow Q_1 \rightarrow P'_1\}, \quad \mathcal{N}_{1,2} = \mathcal{N}_1 \setminus \mathcal{N}_{1,1}.$$

In the following, we will define a sign-reversing involution ϕ on $\mathcal{N}_1 \cup \mathcal{N}_2 \cup \mathcal{N}_3$ such that

$$\phi(\mathcal{N}_{1,1}) = \mathcal{N}_3, \quad \phi(\mathcal{N}_{1,2}) = \mathcal{N}_2.$$

There are several subcases to consider.

- (i) If $i_1 = 0, i_1 \geq 2$, or $i_1 = j_1 = 1$, then $\mathcal{N}_{1,1} = \mathcal{N}_{1,2} = \mathcal{N}_2 = \mathcal{N}_3 = \emptyset$. For these three situations, take ϕ to be the identity map.
- (ii) If $i_1 = 1, j_1 = 0$, and moreover $i_2 \geq 3$ or $j_2 \geq 2$, then $\mathcal{N}_{1,1} = \mathcal{N}_3 = \emptyset$ and $\mathcal{N}_{1,2} = \mathcal{N}_1$. For these situations, we take ϕ to be the map which sends

$$(P_1 \rightarrow P'_0, p_2, \dots, p_k) \in \mathcal{N}_{1,2} \quad \text{and} \quad (P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0, p_2, \dots, p_k) \in \mathcal{N}_2$$

to each other. It is clear that $(P_1 \rightarrow P'_0, p_2, \dots, p_k)$ and $(P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0, p_2, \dots, p_k)$ have opposite weights.

- (iii) If $i_1 = 1, j_1 = 0$ and $i_2 = 2, j_2 = 1$, then

$$\begin{aligned} \mathcal{N}_{1,1} &= \{\mathbf{p} \in \mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J) \mid p_1 = P_1 \rightarrow P'_0, p_2 = P_2 \rightarrow Q_1 \rightarrow P'_1\}, \\ \mathcal{N}_{1,2} &= \{\mathbf{p} \in \mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J) \mid p_1 = P_1 \rightarrow P'_0, p_2 = P_2 \rightarrow Q_2 \rightarrow P'_1\}, \\ \mathcal{N}_2 &= \{\mathbf{p} \in \mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J) \mid p_1 = P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0, p_2 = P_2 \rightarrow Q_2 \rightarrow P'_1\}, \end{aligned}$$

$$\mathcal{N}_3 = \left\{ \mathbf{p} \in \mathcal{N}(\mathbf{P}_I, \mathbf{P}'_J) \left| \begin{array}{l} p_1 = P_1 \rightarrow Q_0 \rightarrow P'_0, \\ \exists l \geq 2 \text{ such that } p_l = P_l \rightarrow Q_l \rightarrow P'_{l-1}, \\ p_m = P_m \rightarrow Q_{m-1} \rightarrow P'_{m-1}, \forall 2 \leq m \leq l-1 \end{array} \right. \right\}.$$

Now we are going to define the map ϕ on $\mathcal{N}_1 \cup \mathcal{N}_2 \cup \mathcal{N}_3$.

If $\mathbf{p} \in \mathcal{N}_{1,1}$, say $\mathbf{p} = (P_1 \rightarrow P'_0, p_2, \dots, p_k)$, then we take l to be the largest number such that $p_m = P_m \rightarrow Q_{m-1} \rightarrow P'_{m-1}$ for $2 \leq m \leq l$, and let

$$\phi(\mathbf{p}) = (P_1 \rightarrow Q_0 \rightarrow P'_0, p_2, \dots, p_{l-1}, P_l \rightarrow Q_l \rightarrow P'_{l-1}, p_{l+1}, \dots, p_k).$$

Thus, $\phi(\mathbf{p}) \in \mathcal{N}_3$.

If $\mathbf{p} \in \mathcal{N}_3$, namely, there exists $l \geq 2$ such that $\mathbf{p} = (P_1 \rightarrow Q_0 \rightarrow P'_0, p_2, \dots, p_k)$ with $p_l = P_l \rightarrow Q_l \rightarrow P'_{l-1}$ and $p_m = P_m \rightarrow Q_{m-1} \rightarrow P'_{m-1}$ for any $2 \leq m \leq l-1$, then let

$$\phi(\mathbf{p}) = (P_1 \rightarrow P'_0, p_2, \dots, p_{l-1}, P_l \rightarrow Q_{l-1} \rightarrow P'_{l-1}, p_{l+1}, \dots, p_k).$$

Here the map ϕ is well-defined since the number l exists then it must be unique by the definition of (\mathcal{P}_3) . It is also clear that $\phi(\mathbf{p}) \in \mathcal{N}_{1,1}$.

If $\mathbf{p} \in \mathcal{N}_{1,2}$, say $\mathbf{p} = (P_1 \rightarrow P'_0, p_2, \dots, p_k)$, then let

$$\phi(\mathbf{p}) = (P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0, p_2, \dots, p_k).$$

Hence, we have $\phi(\mathbf{p}) \in \mathcal{N}_2$.

If $\mathbf{p} \in \mathcal{N}_2$, say $\mathbf{p} = (P_1 \rightarrow Q_1 \xrightarrow{\xi} P'_0, p_2, \dots, p_k)$, then let

$$\phi(\mathbf{p}) = (P_1 \rightarrow P'_0, p_2, \dots, p_k).$$

It is obvious that $\phi(\mathbf{p}) \in \mathcal{N}_{1,2}$.

Figure 3.2 gives an illustration of ϕ for this subcase.

With the above definition of ϕ , it is straightforward to verify that ϕ is a sign-reversing involution on $\mathcal{N}_1 \cup \mathcal{N}_2 \cup \mathcal{N}_3$, as desired. ■

4 The Hankel matrix H

The aim of this section is to give a combinatorial proof of the q -total positivity of the Hankel matrix H of Narayana polynomials of type B . It suffices to combinatorially prove




Figure 3.2: An illustration of ϕ for subcase (iii)

the q -total positivity of each leading principal submatrix of H . To this end, let us first establish the planar networks for its leading principal submatrices.

Let $L_n^B = (l_{i,j}(q))_{0 \leq i,j \leq n+1}$, $B_n = (b_{i,j}(q))_{0 \leq i,j \leq n}$ and $H_n = (h_{i,j}(q))_{0 \leq i,j \leq n}$. By (1.2) it is evident that

$$B_{n+1} = \bar{B}_n L_n^B, \text{ where } \bar{B}_n = \begin{pmatrix} 1 & O \\ O & B_n \end{pmatrix}. \quad (4.1)$$

Aigner [1] proved that

$$H_n = B_n T_n B_n^t, \quad (4.2)$$

where

$$T_n = \begin{pmatrix} 1 & & & & \\ & 2q & & & \\ & & 2q^2 & & \\ & & & \ddots & \\ & & & & 2q^n \end{pmatrix}_{(n+1) \times (n+1)},$$

and B_n^t denotes the transpose of B_n .

These two formulas allow us to recursively construct the planar network for H_n . Precisely, we mainly make use of the following lemma, which provides a way to build a network for a product of matrices. Recall that in a digraph, a vertex v is called a source (resp. sink) if there is no arcs point in (resp. out of) it. The following result could be considered as a corollary of the transfer-matrix method (see [23, Theorem 4.7.1] for instance), but for self-containedness we will give a detailed proof.

Lemma 4.1 *Given k square matrices M_1, M_2, \dots, M_k of order n , for each $1 \leq i \leq k$ assume that $\mathcal{M}_i = (D^{M_i}, \text{wt}_{D^{M_i}}, (u_{i,1}, \dots, u_{i,n}), (v_{i,1}, \dots, v_{i,n}))$ is a planar network for M_i*

with $u_{i,j}$ being a source and $v_{i,j}$ being a sink for all $1 \leq j \leq n$. Let D^M be the digraph obtained by placing $D^{M_1}, D^{M_2}, \dots, D^{M_k}$ in succession and identifying $v_{i,j}$ with $u_{i+1,j}$ for each $1 \leq i \leq k-1$ and $1 \leq j \leq n$, and let wt_{D^M} be the weight function inherited from $\text{wt}_{D^{M_1}}, \dots, \text{wt}_{D^{M_k}}$ in an obvious way. Then

$$\mathcal{M} = (D^M, \text{wt}_{D^M}, (u_{1,1}, \dots, u_{1,n}), (v_{k,1}, \dots, v_{k,n}))$$

is a planar network for the product $M_1 \cdots M_k$.

Proof. The proof is by induction on k . Let us first prove the base case $k = 2$. The construction tells that for any $1 \leq i, j \leq n$, each directed path from $u_{1,i}$ to $v_{2,j}$ must pass through exactly one vertex $v_{1,l}$ ($= u_{2,l}$) for some $1 \leq l \leq n$, and hence

$$GF_{D^M}(u_{1,i}, v_{2,j}) = \sum_{l=1}^n GF_{D^{M_1}}(u_{1,i}, v_{1,l}) GF_{D^{M_2}}(u_{2,l}, v_{2,j}),$$

as desired. Assume the assertion for k ($k \geq 2$). By applying the preceding proof to $M_1 \cdots M_k$ and M_{k+1} , we find that the assertion also holds for $k+1$. This completes the proof. \blacksquare

Now we present the construction for the planar network \mathcal{H}_n for H_n , which is essentially based on \mathcal{L}^B . At first, we give the planar network for L_n^B , which is actually obtained by cutting off the part of \mathcal{L}^B below $y = -n - 1$. Precisely, let $D^{L_n^B}$ be the subgraph of D^{L^B} induced by the vertices $P_0, \dots, P_{n+1}, Q_0, \dots, Q_{n+1}, P'_0, \dots, P'_{n+1}$, and let $\text{wt}_{D^{L_n^B}}$ be the restriction of $\text{wt}_{D^{L^B}}$ to $D^{L_n^B}$. Then $(D^{L_n^B}, \text{wt}_{D^{L_n^B}}, (P_0, \dots, P_{n+1}), (P'_0, \dots, P'_{n+1}))$ is a planar network for L_n^B . Unfortunately, this labeling is not convenient for introducing the recursive construction of \mathcal{H}_n . In the rest of this paper, we will label the vertex P_i by $P_{n+1-i}^{(n)}$, Q_i by $Q_{n+1-i}^{(n)}$, P'_i by $P_{n+1-i}^{(n+1)}$ for $0 \leq i \leq n+1$ in the digraph $D^{L_n^B}$. Moreover, we may shift the digraphs $D^{L_n^B}$ in the plane such that $P_j^{(i)} = (2i, j)$ and $Q_j^{(i)} = (2i+1, j)$ for all $i, j \geq 0$. Then

$$\mathcal{L}_n^B = (D^{L_n^B}, \text{wt}_{D^{L_n^B}}, (P_{n+1}^{(n)}, \dots, P_0^{(n)}), (P_{n+1}^{(n+1)}, \dots, P_0^{(n+1)}))$$

is a planar network for L_n^B , and

$$L_n^B = \left(GF_{D^{L_n^B}}(P_{n+1-i}^{(n)}, P_{n+1-j}^{(n+1)}) \right)_{0 \leq i, j \leq n+1}.$$

Figure 4.1 shows the planar network \mathcal{L}_2^B .

Using the planar networks $\mathcal{L}_0^B, \mathcal{L}_1^B, \dots, \mathcal{L}_n^B$, we can construct the planar network \mathcal{B}_{n+1} for B_{n+1} .




Figure 4.1: Planar network \mathcal{L}_2^B

- For $n = 0$, we take \mathcal{B}_1 to be the planar network \mathcal{L}_0^B since $B_1 = L_0^B$.
- Assuming that \mathcal{B}_n has been constructed for some $n \geq 1$, we continue to construct \mathcal{B}_{n+1} . Let $D^{\bar{B}_n}$ be the digraph with $V(D^{\bar{B}_n}) = V(D^{B_n}) \cup \{P_{n+1}^{(0)}, P_{n+1}^{(1)}, \dots, P_{n+1}^{(n)}\}$ and $A(D^{\bar{B}_n}) = A(D^{B_n}) \cup \{P_{n+1}^{(i)} \rightarrow P_{n+1}^{(i+1)} \mid 0 \leq i \leq n-1\}$, and let $\text{wt}_{D^{\bar{B}_n}}(a)$ be equal to $\text{wt}_{D^{B_n}}(a)$ for $a \in A(D^{B_n})$ and equal to 1 for the other arcs. Then

$$\bar{B}_n = (D^{\bar{B}_n}, \text{wt}_{D^{\bar{B}_n}}, (P_{n+1}^{(0)}, P_n^{(0)}, \dots, P_0^{(0)}), (P_{n+1}^{(n)}, P_n^{(n)}, \dots, P_0^{(n)}))$$

is a planar network for \bar{B}_n . By (4.1) and Lemma 4.1, we obtain that

$$\mathcal{B}_{n+1} = (D^{B_{n+1}}, \text{wt}_{D^{B_{n+1}}}, (P_{n+1}^{(0)}, P_n^{(0)}, \dots, P_0^{(0)}), (P_{n+1}^{(n+1)}, P_n^{(n+1)}, \dots, P_0^{(n+1)}))$$

is a planar network for B_{n+1} , where $D^{B_{n+1}}$ and $\text{wt}_{D^{B_{n+1}}}$ are defined in the way as described in Lemma 4.1. See Figure 4.2 for an illustration of D^{B_3} .

Based on (4.2) and Lemma 4.1, we proceed to build the planar network for H_n from \mathcal{B}_n . Firstly, we construct a planar network for B_n^t . We take $D^{B_n^t}$ to be the digraph obtained by reflecting D^{B_n} about the vertical line $x = 2n + 1/2$ and reversing the direction of all arcs. We also label the image of $P_j^{(i)}$ (resp. $Q_j^{(i)}$) by $\bar{P}_j^{(i)}$ (resp. $\bar{Q}_j^{(i)}$). We also let $\text{wt}_{D^{B_n^t}}$ be the function which assigns to each arc of $D^{B_n^t}$ the weight of its preimage. Then it is easy to verify that

$$\mathcal{B}_n^t = (D^{B_n^t}, \text{wt}_{D^{B_n^t}}, (\bar{P}_n^{(n)}, \dots, \bar{P}_0^{(n)}), (\bar{P}_n^{(0)}, \dots, \bar{P}_0^{(0)}))$$

is a planar network for B_n^t . Next, we define D^{T_n} to be the digraph whose vertex set is $\{P_i^{(n)} \mid 0 \leq i \leq n\} \cup \{\bar{P}_j^{(n)} \mid 0 \leq j \leq n\}$ and arc set is $\{P_i^{(n)} \rightarrow \bar{P}_i^{(n)} \mid 0 \leq i \leq n\}$, and let




Figure 4.2: Digraph D^{B_3}

$\text{wt}_{D^{T_n}}(P_i^{(n)} \rightarrow \bar{P}_i^{(n)}) = (T_n)_{i,i}$ for $0 \leq i \leq n$. Then

$$\mathcal{T}_n = (D^{T_n}, \text{wt}_{D^{T_n}}, (P_n^{(n)}, \dots, P_0^{(n)}), (\bar{P}_n^{(n)}, \dots, \bar{P}_0^{(n)}))$$

is a planar network for T_n . Finally, we combine \mathcal{B}_n , \mathcal{T}_n and $\bar{\mathcal{B}}_n^t$ to get the following planar network for H_n :

$$\mathcal{H}_n = (D^{H_n}, \text{wt}_{D^{H_n}}, (P_n^{(0)}, \dots, P_0^{(0)}), (\bar{P}_n^{(0)}, \dots, \bar{P}_0^{(0)})), \quad (4.3)$$

where D^{H_n} and $\text{wt}_{D^{H_n}}$ are defined in the way as described in Lemma 4.1. Figure 4.3 shows the digraph D^{H_3} .




Figure 4.3: Digraph D^{H_3}

We are now in a position to give a combinatorial proof of the q -total positivity of H_n for any nonnegative integer n . Given a positive integer k and two sequences $I = (i_1, \dots, i_k)$, $J = (j_1, \dots, j_k)$ of indices such that $0 \leq i_1 < \dots < i_k \leq n$ and $0 \leq j_1 < \dots < j_k \leq n$, let

$$\mathbf{P}_I = (P_{n-i_1}^{(0)}, \dots, P_{n-i_k}^{(0)}), \quad \bar{\mathbf{P}}_J = (\bar{P}_{n-j_1}^{(0)}, \dots, \bar{P}_{n-j_k}^{(0)}). \quad (4.4)$$

Let $H_{I,J}$ denote the submatrix of H_n whose rows are indexed by I and columns indexed by J . By Lemma 2.1 and (4.3), we have

$$\det [H_{I,J}] = GF(\mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J)). \quad (4.5)$$

We further need to find a subset of $\mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J)$, say $\mathcal{S}_{I,J}^{H_n}$, which will play the same role as $S_{I,J}$ in Theorem 3.2.

Observe that by the recursive construction of D^{H_n} , it can be naturally divided into $2n + 1$ parts: $D_1^{H_n}, \dots, D_n^{H_n}, D^{T_n}, \bar{D}_n^{H_n}, \dots, \bar{D}_1^{H_n}$, where for each $1 \leq i \leq n$ the graph $D_i^{H_n}$ is the subgraph of D^{H_n} induced by the vertices $P_n^{(i-1)}, \dots, P_0^{(i-1)}, Q_i^{(i-1)}, \dots, Q_0^{(i-1)}, P_n^{(i)}, \dots, P_0^{(i)}$, and $\bar{D}_i^{H_n}$ is the subgraph of D^{H_n} induced by the vertices $\bar{P}_n^{(i)}, \dots, \bar{P}_0^{(i)}, \bar{Q}_i^{(i-1)}, \dots, \bar{Q}_0^{(i-1)}, \bar{P}_n^{(i-1)}, \dots, \bar{P}_0^{(i-1)}$. Graphically, D^{H_n} is divided into $2n + 1$ parts by $2n$ lines parallel to the y -axis. Thus, each member $\mathbf{p} = (p_1, \dots, p_k) \in \mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J)$ is also divided into $2n + 1$ nonintersecting families $\mathbf{p}_1, \dots, \mathbf{p}_n, \mathbf{p}_T, \bar{\mathbf{p}}_n, \dots, \bar{\mathbf{p}}_1$ by these lines, where \mathbf{p}_i (resp. $\bar{\mathbf{p}}_i$) is the restriction of \mathbf{p} to $D_i^{H_n}$ (resp. $\bar{D}_i^{H_n}$) for each $1 \leq i \leq n$, and \mathbf{p}_T is the restriction of \mathbf{p} to D^{T_n} . For this reason, we may adopt the notation

$$\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_n, \mathbf{p}_T, \bar{\mathbf{p}}_n, \dots, \bar{\mathbf{p}}_1)$$

to represent a nonintersecting family of $\mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J)$.

Note that $D_i^{H_n}$ ($1 \leq i \leq n$) can be regarded as the digraph obtained by adding $n - i$ parallel arcs (namely, $P_n^{(i-1)} \rightarrow P_n^{(i)}, \dots, P_{i+1}^{(i-1)} \rightarrow P_{i+1}^{(i)}$) to $D^{L_{i-1}^B}$. By simply mimicking the definitions of (\mathcal{P}_1) , (\mathcal{P}_2) and (\mathcal{P}_3) as given immediately before (3.2), we may define the following properties on nonintersecting families \mathbf{p}_i in $D_i^{H_n}$:

$(\mathcal{P}_1^{(i)})$ There exists $1 \leq j \leq k$ such that the j -th component of \mathbf{p}_i is the directed path $P_{i-1}^{(i-1)} \rightarrow P_i^{(i)}$;

$(\mathcal{P}_2^{(i)})$ There exists $1 \leq j \leq k$ such that the j -th component of \mathbf{p}_i is the directed path $P_{i-1}^{(i-1)} \rightarrow Q_{i-1}^{(i-1)} \xrightarrow{\Sigma} P_i^{(i)}$;

$(\mathcal{P}_3^{(i)})$ There exist $1 \leq j \leq k$ and $l \geq 2$ such that the $(j + l - 1)$ -th component of \mathbf{p}_i is $P_{i-l}^{(i-1)} \rightarrow Q_{i-l}^{(i-1)} \rightarrow P_{i-l+1}^{(i)}$ and the m -th component is $P_{i-1-(m-j)}^{(i-1)} \rightarrow Q_{i-(m-j)}^{(i-1)} \rightarrow P_{i-(m-j)}^{(i)}$ for each $j \leq m \leq j + l - 2$.

For $\bar{\mathbf{p}}_i$ in $\bar{D}_i^{H_n}$, if its preimage with respect to the reflection satisfies $(\mathcal{P}_1^{(i)})$, $(\mathcal{P}_2^{(i)})$, or $(\mathcal{P}_3^{(i)})$, we say that $\bar{\mathbf{p}}_i$ satisfies Property $(\bar{\mathcal{P}}_1^{(i)})$, $(\bar{\mathcal{P}}_2^{(i)})$, or $(\bar{\mathcal{P}}_3^{(i)})$, respectively. Then we take

$$\mathcal{S}_{I,J}^{H_n} = \left\{ \mathbf{p} \in \mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J) \left| \begin{array}{l} \mathbf{p}_i \text{ satisfies none of } (\mathcal{P}_1^{(i)}), (\mathcal{P}_2^{(i)}), (\mathcal{P}_3^{(i)}) \text{ and} \\ \bar{\mathbf{p}}_i \text{ satisfies none of } (\bar{\mathcal{P}}_1^{(i)}), (\bar{\mathcal{P}}_2^{(i)}), (\bar{\mathcal{P}}_3^{(i)}) \\ \text{for each } 1 \leq i \leq n \end{array} \right. \right\}. \quad (4.6)$$

It is clear that each $\mathbf{p} \in \mathcal{S}_{I,J}^{H_n}$ has a q -nonnegative weight.

We would like to point out that the involution ϕ defined in the proof of Theorem 3.2 can also be mimicked to define a sign-reversing involution ϕ_i on nonintersecting families \mathbf{p}_i in $D_i^{H_n}$. Suppose that $\mathbf{p}_i = (p_{i,1}, \dots, p_{i,k})$ and $p_{i,1}, \dots, p_{i,m}$ are those parallel arcs out of $D^{L_{i-1}^B}$. Then $(p_{i,m+1}, \dots, p_{i,k})$ is a nonintersecting family in $D^{L_{i-1}^B}$. If $\phi((p_{i,m+1}, \dots, p_{i,k})) = (p'_{i,m+1}, \dots, p'_{i,k})$, then define

$$\phi_i(\mathbf{p}_i) = (p_{i,1}, \dots, p_{i,m}, p'_{i,m+1}, \dots, p'_{i,k}). \quad (4.7)$$

Similarly, we can define a sign-reversing involution $\bar{\phi}_i$ on nonintersecting families $\bar{\mathbf{p}}_i$ in $\bar{D}_i^{H_n}$. Note that if \mathbf{p}_i satisfies Property $(\mathcal{P}_1^{(i)})$, $(\mathcal{P}_2^{(i)})$, or $(\mathcal{P}_3^{(i)})$, then $(p'_{i,m+1}, \dots, p'_{i,k})$ satisfies Property (\mathcal{P}_1) , (\mathcal{P}_2) , or (\mathcal{P}_3) (with a change of labeling), respectively, and hence $\phi((p_{i,m+1}, \dots, p_{i,k})) \neq (p_{i,m+1}, \dots, p_{i,k})$ and $\phi_i(\mathbf{p}_i) \neq \mathbf{p}_i$. An analogous result holds for $\bar{\mathbf{p}}_i$ and $\bar{\phi}_i$.

The main result of this section is as follows, which provides a combinatorial proof of the q -total positivity of H .

Theorem 4.2 *Given a nonnegative integer n and two sequences $I = (i_1, \dots, i_k)$, $J = (j_1, \dots, j_k)$ of indices such that $0 \leq i_1 < \dots < i_k \leq n$ and $0 \leq j_1 < \dots < j_k \leq n$, let $H_{I,J}$ denote the submatrix of H_n whose rows are indexed by I and columns indexed by J , let $\mathbf{P}_I, \bar{\mathbf{P}}_J$ be as given by (4.4), and let $\mathcal{S}_{I,J}^{H_n}$ be as given by (4.6). Then we have*

$$\det [H_{I,J}] = GF(\mathcal{S}_{I,J}^{H_n}), \quad (4.8)$$

where $GF(\mathcal{S}_{I,J}^{H_n})$ denotes the sum of weights of all elements in $\mathcal{S}_{I,J}^{H_n}$. In particular, H is q -totally positive.

Proof. By (4.5), it suffices to give a sign-reversing involution, say ϕ^{H_n} , on $\mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J)$ with $\mathcal{S}_{I,J}^{H_n}$ being the set of all fixed points. We proceed to define ϕ^{H_n} by using the aforementioned involutions ϕ_i and $\bar{\phi}_i$, see (4.7). Given $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_n, \mathbf{p}_T, \bar{\mathbf{p}}_n, \dots, \bar{\mathbf{p}}_1) \in \mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J)$, if $\mathbf{p} \in \mathcal{S}_{I,J}^{H_n}$, then let $\phi^{H_n}(\mathbf{p}) = \mathbf{p}$.

Next, we consider the case $\mathbf{p} \in \mathcal{N}_{D^{H_n}}(\mathbf{P}_I, \bar{\mathbf{P}}_J) \setminus \mathcal{S}_{I,J}^{H_n}$. If there exists some i such that \mathbf{p}_i satisfies Property $(\mathcal{P}_1^{(i)})$, $(\mathcal{P}_2^{(i)})$ or $(\mathcal{P}_3^{(i)})$, or equivalently, $\phi_i(\mathbf{p}_i) \neq \mathbf{p}_i$, then let l be the smallest such index and

$$\phi^{H_n}((\mathbf{p}_1, \dots, \mathbf{p}_n, \mathbf{p}_T, \bar{\mathbf{p}}_n, \dots, \bar{\mathbf{p}}_1)) = (\mathbf{p}_1, \dots, \phi_l(\mathbf{p}_l), \dots, \mathbf{p}_n, \mathbf{p}_T, \bar{\mathbf{p}}_n, \dots, \bar{\mathbf{p}}_1).$$

Otherwise, if such an index does not exist, then there must exist some i such that $\bar{\mathbf{p}}_i$ satisfies Property $(\bar{\mathcal{P}}_1^{(i)})$, $(\bar{\mathcal{P}}_2^{(i)})$ or $(\bar{\mathcal{P}}_3^{(i)})$, or equivalently, $\bar{\phi}_i(\bar{\mathbf{p}}_i) \neq \bar{\mathbf{p}}_i$. In this subcase, we let l be the largest such index and define

$$\phi^{H_n}((\mathbf{p}_1, \dots, \mathbf{p}_n, \mathbf{p}_T, \bar{\mathbf{p}}_n, \dots, \bar{\mathbf{p}}_1)) = (\mathbf{p}_1, \dots, \mathbf{p}_n, \mathbf{p}_T, \bar{\mathbf{p}}_n, \dots, \bar{\phi}_l(\bar{\mathbf{p}}_l), \dots, \bar{\mathbf{p}}_1).$$

By the construction of the involutions ϕ_i and $\bar{\phi}_i$ for $1 \leq i \leq n$, it is easy to verify that ϕ^{H_n} is a sign-reversing involution on nonintersecting families in D^{H_n} . Hence ϕ^{H_n} induces the q -total positivity of H_n in the same way that ϕ induces the q -total positivity of L_n^B . Further, the q -total positivity of H_n implies the q -total positivity of H since each minor of H is a minor of H_n for some n . The proof is complete. \blacksquare

By applying the same kind of reasoning of the proof of Theorem 4.2, we can give a combinatorial proof of the q -total positivity of the triangular array $B = (b_{n,k}(q))_{n,k \geq 0}$ as defined by (1.2). We leave the details to the reader. It is interesting to note that similar reasoning can be used to establish the following result.

Theorem 4.3 *Let $C = (c_{n,k}(q))_{n,k \geq 0}$ be the Catalan-Stieltjes matrix generated by one of the following two recurrences:*

(1) *We have*

$$\begin{aligned} c_{n,0}(q) &= [(f - e) + eq] \cdot c_{n-1,0}(q) + fq \cdot c_{n-1,1}(q); \\ c_{n,k}(q) &= c_{n-1,k-1}(q) + (1 + q) \cdot c_{n-1,k}(q) + q \cdot c_{n-1,k+1}(q), \quad (k \geq 1, n \geq 1) \end{aligned}$$

for some $f \geq e \geq 0$.

(2) *We have*

$$\begin{aligned} c_{n,0}(q) &= [(f - 1) + eq] \cdot c_{n-1,0}(q) + eq \cdot c_{n-1,1}(q); \\ c_{n,k}(q) &= c_{n-1,k-1}(q) + (1 + eq) \cdot c_{n-1,k}(q) + eq \cdot c_{n-1,k+1}(q), \quad (k \geq 1, n \geq 1) \end{aligned}$$

for some $e, f \geq 1$.

Then the Hankel matrix $(c_{i+j,0})_{i,j \geq 0}$ is q -totally positive.

Proof. Note that, for either case, we can construct a planar network for the leading principal submatrix $(c_{i+j,0})_{0 \leq i,j \leq n}$ by using the same underlying graph D^{H_n} of the planar network for H_n , but with a different weight function. Since the weight function of \mathcal{H}_n is naturally inherited from that of \mathcal{L}^B , it is sufficient to assign a new weight function to D^{L^B} .

For the first case, we let

$$\begin{aligned} \text{wt}_{D^{L^B}}(P_1 \rightarrow Q_0) &= eq, & \text{wt}_{D^{L^B}}(P_i \rightarrow Q_{i-1}) &= q \text{ for } i \geq 2, \\ \text{wt}_{D^{L^B}}(P_1 \rightarrow P'_0) &= -e, & \text{wt}_{D^{L^B}}(Q_1 \xrightarrow{\xi} P'_0) &= e, \\ \text{wt}_{D^{L^B}}(Q_1 \xrightarrow{\alpha} P'_0) &= f - e, \end{aligned}$$

and $\text{wt}_{D^{LB}}(a) = 1$ for the other arcs a in D^{LB} .

For the second case, we let

$$\begin{aligned} \text{wt}_{D^{LB}}(P_i \rightarrow Q_{i-1}) &= eq \text{ for } i \geq 1, & \text{wt}_{D^{LB}}(P_1 \rightarrow P'_0) &= -1, \\ \text{wt}_{D^{LB}}(Q_1 \xrightarrow{\xi} P'_0) &= 1, & \text{wt}_{D^{LB}}(Q_1 \xrightarrow{\eta} P'_0) &= f - 1, \end{aligned}$$

and $\text{wt}_{D^{LB}}(a) = 1$ for the other arcs a in D^{LB} .

With these new weights, it is straightforward to verify that the involution ϕ^{H_n} constructed in the proof of Theorem 4.2 is still a sign-reversing involution on nonintersecting families of D^{H_n} . As a result, we obtain the q -total positivity of $(c_{i+j,0})_{i,j \geq 0}$. \blacksquare

5 A conjecture on immanant positivity

Let $M = (m_{i,j})$ be a square matrix of order n , \mathfrak{S}_n be the symmetric group of order n , λ be a partition of n , and χ^λ be the irreducible character of \mathfrak{S}_n associated with λ . Recall that the immanant of M with respect to λ is defined by

$$\text{Imm}_\lambda M = \sum_{\sigma \in \mathfrak{S}_n} \chi^\lambda(\sigma) m_{1,\sigma(1)} \cdots m_{n,\sigma(n)}.$$

When $\lambda = (1^n)$, the immanant $\text{Imm}_\lambda M$ specializes to $\det[M]$. In [13] we proved the immanant positivity for a large family of Catalan-Stieltjes matrices and their associated Hankel matrices. This motivates us to study the immanant positivity for the Hankel matrix H defined as in (1.1). We have the following conjecture.

Conjecture 5.1 *Let $k \geq 1$ and $I = (i_1, \dots, i_k)$, $J = (j_1, \dots, j_k)$ be two sequences of indices with $0 \leq i_1 < \dots < i_k$ and $0 \leq j_1 < \dots < j_k$. Let $H_{I,J}$ be the submatrix of H whose rows are indexed by I and columns are indexed by J . Then*

$$\text{Imm}_\lambda H_{I,J} \geq_q 0$$

for any partition λ of k .

We have verified the immanant positivity of all square submatrices of H_n for $n \leq 6$ by Sage [21]. Note that, our method in [13] does not apply to H directly, since there exist some arcs weighted by -1 in our planar network for H .

Acknowledgments. We are very grateful to the referee for valuable comments. This work is supported in part by the Fundamental Research Funds for the Central Universities and the National Science Foundation of China (Nos. 11522110, 11971249).

References

- [1] M. Aigner, Catalan and other numbers: a recurrent theme, in: H. Crapo, D. Senato (Eds.), Algebraic combinatorics and computer science, 347–390, Springer Italia, Milan, 2001.
- [2] L.M. Butler, The q -log-concavity of q -binomial coefficients, J. Combin. Theory Ser. A **54** (1990), no. 1, 54–63.
- [3] W.Y.C. Chen, R.L. Tang, L.X.W. Wang and A.L.B. Yang, The q -log-convexity of the Narayana polynomials of type B , Adv. in Appl. Math. **44** (2010), no. 2, 85–110.
- [4] W.Y.C. Chen, L.X.W. Wang and A.L.B. Yang, Schur positivity and the q -log-convexity of the Narayana polynomials, J. Algebraic Combin. **32** (2010), no. 3, 303–338.
- [5] W.Y.C. Chen, L.X.W. Wang and A.L.B. Yang, Recurrence relations for strongly q -log-convex polynomials, Canad. Math. Bull. **54** (2011), no. 2, 217–229.
- [6] X. Chen, H. Liang and Y. Wang, Total positivity of recursive matrices, Linear Algebra Appl. **471** (2015), 383–393.
- [7] I.M. Gessel and X.G. Viennot, Binomial determinants, paths, and hook length formulae, Adv. in Math. **58** (1985), no. 3, 300–321.
- [8] I.M. Gessel and X.G. Viennot, Determinants, paths, and plane partitions, preprint.
- [9] A.M. Hamel and I.P. Goulden, Planar decompositions of tableaux and Schur function determinants, European J. Combin. **16** (1995), no. 5, 461–477.
- [10] C. Krattenthaler, On the q -log-concavity of Gaussian binomial coefficients, Monatsh. Math. **107** (1989), no. 4, 333–339.
- [11] C. Krattenthaler, Oscillating tableaux and nonintersecting lattice paths, J. Statist. Plann. Inference **54** (1996), no. 1, 75–85.
- [12] P. Leroux, Reduced matrices and q -log-concavity properties of q -Stirling numbers, J. Combin. Theory Ser. A **54** (1990), no. 1, 64–84.
- [13] E.Y.H. Li, G.M.X. Li, A.L.B. Yang and C.X.T. Zhang, Immanant Positivity for Catalan-Stieltjes Matrices, arXiv:2106.12816.

- [14] H. Liang, L. Mu and Y. Wang, Catalan-like numbers and Stieltjes moment sequences, *Discrete Math.* **339** (2016), no. 2, 484–488.
- [15] B. Lindström, On the vector representations of induced matroids, *Bull. London Math. Soc.* **5** (1973), 85–90.
- [16] L.L. Liu and Y. Wang, On the log-convexity of combinatorial sequences, *Adv. in Appl. Math.* **39** (2007), no. 4, 453–476.
- [17] M.E. Mays and J. Wojciechowski, A determinant property of Catalan numbers, *Discrete Math.* **211** (2000), no. 1-3, 125–133.
- [18] Q. Pan and J. Zeng, On total positivity of Catalan-Stieltjes matrices, *Electron. J. Combin.* **23** (2016), no. 4, Paper 4.33, 18 pp.
- [19] B.E. Sagan, Inductive proofs of q -log concavity, *Discrete Math.* **99** (1992), no. 1-3, 289–306.
- [20] B.E. Sagan, Log concave sequences of symmetric functions and analogs of the Jacobi-Trudi determinants, *Trans. Amer. Math. Soc.* **329** (1992), no. 2, 795–811.
- [21] SageMath, the Sage Mathematics Software System (Version 9.1), The Sage Developers, 2020, <https://www.sagemath.org>.
- [22] A.D. Sokal, Total positivity: A concept at the interface between algebra, analysis and combinatorics, talk given at Institut Camille Jordan, Université Lyon 1, April 20 2015.
- [23] R.P. Stanley, *Enumerative combinatorics. Volume 1. Second edition.* Cambridge Studies in Advanced Mathematics, 49. Cambridge University Press, Cambridge, 2012.
- [24] J.R. Stembridge, Nonintersecting paths, Pfaffians, and plane partitions, *Adv. Math.* **83** (1990), no. 1, 96–131.
- [25] Y. Wang, B.-X. Zhu, Log-convex and Stieltjes moment sequences, *Adv. in Appl. Math.* **81** (2016), 115–127.
- [26] B.-X. Zhu, Log-convexity and strong q -log-convexity for some triangular arrays, *Adv. in Appl. Math.* **50** (2013), no. 4, 595–606.
- [27] B.-X. Zhu, Some positivities in certain triangular arrays, *Proc. Amer. Math. Soc.* **142** (2014), no. 9, 2943–2952.