



ELSEVIER

Available online at www.sciencedirect.com

European Journal of Combinatorics xx (xxxx) xxx–xxx

European Journal
of Combinatoricswww.elsevier.com/locate/ej

On Postnikov's hook length formula for binary trees

William Y.C. Chen, Laura L.M. Yang

Center for Combinatorics, LPMC-TJKLC, Nankai University, Tianjin 300071, PR China

Received 15 November 2005; accepted 22 November 2007

Abstract

We present a combinatorial proof of Postnikov's hook length formula for binary trees.
© 2007 Elsevier Ltd. All rights reserved.

Let $[n] = \{1, 2, \dots, n\}$. It is well known that the number of labeled trees on $[n]$ equals n^{n-2} , and the number of rooted trees on $[n]$ equals n^{n-1} [5,8]. Recently, Postnikov [6] derived an identity on binary trees and asked for a combinatorial proof [6]. We adopt the terminology of Postnikov [6]. Given a binary tree T and a vertex v of T , we use $h(v)$ to denote the hook length of v , namely, the number of descendants of v (including v itself). Postnikov's hook length formula for binary trees reads as follows [6].

Theorem 1. For $n \geq 1$, we have

$$(n+1)^{n-1} = \sum_T \frac{n!}{2^n} \prod_{v \in T} \left(1 + \frac{1}{h(v)}\right), \quad (1)$$

where the sum ranges over all binary trees T with n vertices.

Our combinatorial proof is based on the following reformulation of (1) in terms of rooted trees:

$$(n+1)^n = \sum_T \frac{(n+1)!}{2^n} \prod_{v \in T} \left(1 + \frac{1}{h(v)}\right). \quad (2)$$

Proof of (2). Let F_n denote the sum on the right hand side of (2). For any unlabeled binary tree T with n vertices, the hook length of the root is always n . Let us consider a binary tree T such

E-mail addresses: chen@nankai.edu.cn (W.Y.C. Chen), yanglm@hotmail.com (L.L.M. Yang).

0195-6698/\$ - see front matter © 2007 Elsevier Ltd. All rights reserved.
doi:10.1016/j.ejc.2007.11.025

Please cite this article in press as: W.Y.C. Chen, L.L.M. Yang, On Postnikov's hook length formula for binary trees, European Journal of Combinatorics (2007), doi:10.1016/j.ejc.2007.11.025

1 that the left subtree T_1 has k vertices and the right subtree T_2 has $n - k - 1$ vertices. From the
 2 relation

$$3 \quad \frac{(n+1)!}{2^n} \left(1 + \frac{1}{n}\right) = \frac{n+1}{2n} \binom{n+1}{k+1} \frac{(k+1)!}{2^k} \frac{(n-k)!}{2^{n-k-1}},$$

4 it can be deduced that

$$5 \quad F_n = \frac{n+1}{2n} \sum_{k=0}^{n-1} \binom{n+1}{k+1} \sum_{T_1} \frac{(k+1)!}{2^k} \prod_{v \in T_1} \left(1 + \frac{1}{h(v)}\right) \sum_{T_2} \frac{(n-k)!}{2^{n-k-1}} \prod_{v \in T_2} \left(1 + \frac{1}{h(v)}\right),$$

6 where T_1 and T_2 range over all binary trees on k and $n - k - 1$ vertices, respectively. Hence F_n
 7 satisfies the following recurrence relation:

$$8 \quad F_n = \frac{n+1}{2n} \sum_{k=0}^{n-1} \binom{n+1}{k+1} F_k F_{n-k-1}. \tag{3}$$

9 It is known that the number $T_n = n^{n-2}$ of labeled trees with n vertices satisfies the recurrence
 10 relation

$$11 \quad 2nT_{n+1} = \sum_{k=0}^{n-1} \binom{n+1}{k+1} (k+1)T_{k+1}(n-k)T_{n-k}. \tag{4}$$

12 Let $R_n = nT_n$ denote the number of rooted trees on n vertices. Then the above recurrence (4)
 13 can be recast as

$$14 \quad R_{n+1} = \frac{n+1}{2n} \sum_{k=0}^{n-1} \binom{n+1}{k+1} R_{k+1}R_{n-k}. \tag{5}$$

15 A combinatorial interpretation of (4) is given by Moon [5]: The left hand side of (4) equals the
 16 number of labeled trees on $[n+1]$ with a distinguished edge and a direction on this distinguished
 17 edge. Let T be such a tree, we may decompose it into an ordered pair of rooted trees by cutting
 18 off the distinguished edge.

19 Combining the recurrence (3) of F_n with the recurrence (5) of R_n , we arrive at the conclusion
 20 **Q1** that $F_n = R_{n+1} = (n+1)^n$. Thus we obtain (2). ■

21 We note that Seo [7] also found a combinatorial proof of the identity (1). Further studies
 22 related to Postnikov's hook length formula (1) have been carried out by Du and Liu [1], Gessel
 23 and Seo [2], Hivert, Novelli and Thibon [3], and Liu [4].

24 **Acknowledgments**

25 The authors wish to thank Richard Stanley for helpful comments. This work was supported
 26 by the 973 Project, the PCSIRT Project of the Ministry of Education, the Ministry of Science and
 27 Technology, and the National Science Foundation of China.

28 **References**

29 [1] R.R.X. Du, F. Liu, (k, m) -Catalan numbers and hook length polynomials for plane trees, European J. Combin. 28
 30 (2007) 1312–1321.
 31 [2] I.M. Gessel, S. Seo, A refinement of Cayley's formula for trees, Electron. J. Combin. 11 (2) (2006) #R27.

- Q2 [3] F. Hivert, J.-C. Novelli, J.-Y. Thibon, Trees, functional equations, and combinatorial Hopf algebras, *European J. Combin.* (in press). 1
- [4] F. Liu, Hook length polynomials for plane forests of a certain type. [arXiv:math.CO/0511055](https://arxiv.org/abs/math.CO/0511055). 2
- [5] J.W. Moon, Counting labelled trees, in: *Canadian Mathematical Monographs*, 1970. 3
- [6] A. Postnikov, Permutohedra, associahedra, and beyond, in: *Conference in Honor of Richard Stanley's Sixtieth Birthday*, June 26, 2004, MIT. [arXiv:math.Co/0507163](https://arxiv.org/abs/math.CO/0507163). 4
- [7] S. Seo, A combinatorial proof of Postnikov's identity and a generalized enumeration of labeled trees, *Electron. J. Combin.* 11 (2) (2005) #N3. 5
- [8] R.P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, Cambridge, UK, 1999. 6
- 7
- 8
- 9

UNCORRECTED PROOF