

# Hypergraph Turán numbers of vertex disjoint cycles\*

Ran Gu, Xueliang Li, Yongtang Shi

Center for Combinatorics and LPMC-TJKLC

Nankai University, Tianjin 300071, China

guran323@163.com, lxl@nankai.edu.cn, shi@nankai.edu.cn

## Abstract

The Turán number of a  $k$ -uniform hypergraph  $H$ , denoted by  $ex_k(n; H)$ , is the maximum number of edges in any  $k$ -uniform hypergraph  $F$  on  $n$  vertices which does not contain  $H$  as a subgraph. Let  $\mathcal{C}_\ell^{(k)}$  denote the family of all  $k$ -uniform minimal cycles of length  $\ell$ ,  $\mathcal{S}(\ell_1, \dots, \ell_r)$  denote the family of hypergraphs consisting of unions of  $r$  vertex disjoint minimal cycles of length  $\ell_1, \dots, \ell_r$ , respectively, and  $\mathbb{C}_\ell^{(k)}$  denote a  $k$ -uniform linear cycle of length  $\ell$ . We determine precisely  $ex_k(n; \mathcal{S}(\ell_1, \dots, \ell_r))$  and  $ex_k(n; \mathbb{C}_{\ell_1}^{(k)}, \dots, \mathbb{C}_{\ell_r}^{(k)})$  for sufficiently large  $n$ . The results extend recent results of Füredi and Jiang who determined the Turán numbers for single  $k$ -uniform minimal cycles and linear cycles.

**Keywords:** Turán number; cycles; extremal hypergraphs

**AMS Subject Classification (2010):** 05D05, 05C35, 05C65

## 1 Introduction

In this paper, we employ standard definitions and notation from hypergraph theory (see e.g., [1]). A *hypergraph* is a pair  $H = (V, E)$  consisting of a set  $V$  of vertices and a set  $E \subseteq \mathcal{P}(V)$  of edges. If every edge contains exactly  $k$  vertices, then  $H$  is a  *$k$ -uniform hypergraph*. For two hypergraphs  $G$  and  $H$ , we write  $G \subseteq H$  if there is an injective homomorphism from  $G$  into  $H$ . We use  $G \cup H$  to denote the disjoint union

---

\*Supported by NSFC and the “973” program.

of (hyper)graphs  $G$  and  $H$ . By disjoint, we will always mean vertex disjoint. A *Berge path* of length  $\ell$  is a family of distinct sets  $\{F_1, \dots, F_\ell\}$  and  $\ell + 1$  distinct vertices  $v_1, \dots, v_{\ell+1}$  such that for each  $1 \leq i \leq \ell$ ,  $F_i$  contains  $v_i$  and  $v_{i+1}$ . Let  $\mathcal{B}_\ell^{(k)}$  denote the family of  $k$ -uniform Berge paths of length  $\ell$ . A *linear path* of length  $\ell$  is a family of sets  $\{F_1, \dots, F_\ell\}$  such that  $|F_i \cap F_{i+1}| = 1$  for each  $i$  and  $F_i \cap F_j = \emptyset$  whenever  $|i - j| > 1$ . Let  $\mathbb{P}_\ell^{(k)}$  denote the  $k$ -uniform linear path of length  $\ell$ . It is unique up to isomorphisms. A  $k$ -uniform *Berge cycle* of length  $\ell$  is a cyclic list of distinct  $k$ -sets  $A_1, \dots, A_\ell$  and  $\ell$  distinct vertices  $v_1, \dots, v_\ell$  such that for each  $1 \leq i \leq \ell$ ,  $A_i$  contains  $v_i$  and  $v_{i+1}$  (where  $v_{\ell+1} = v_1$ ). A  $k$ -uniform *minimal cycle* of length  $\ell$  is a cyclic list of  $k$ -sets  $A_1, \dots, A_\ell$  such that consecutive sets intersect in at least one element and nonconsecutive sets are disjoint. Denote the family of all  $k$ -uniform minimal cycles of length  $\ell$  by  $\mathcal{C}_\ell^{(k)}$ . A  $k$ -uniform *linear cycle* of length  $\ell$ , denoted by  $\mathbb{C}_\ell^{(k)}$ , is a cyclic list of  $k$ -sets  $A_1, \dots, A_\ell$  such that consecutive sets intersect in exactly one element and nonconsecutive sets are disjoint.

The *Turán number*, or extremal number, of a  $k$ -uniform hypergraph  $H$ , denoted by  $ex_k(n; H)$ , is the maximum number of edges in any  $k$ -uniform hypergraph  $F$  on  $n$  vertices which does not contain  $H$  as a subgraph. This is a natural generalization of the classical Turán number for 2-uniform graphs; we restrict ourselves to the case of  $k$ -uniform hypergraphs. Let  $ex_k(n; F_1, F_2, \dots, F_r)$  denote the  $k$ -uniform hypergraph Turán Number of a list of  $k$ -uniform hypergraphs  $F_1, F_2, \dots, F_r$ , i.e.,  $ex_k(n; F_1, F_2, \dots, F_r) = ex_k(n; F_1 \cup F_2 \cup \dots \cup F_r)$ .

For the class of  $k$ -uniform Berge paths of length  $\ell$ , Györi et al [5] determined  $ex_k(n; \mathcal{B}_\ell^{(k)})$  exactly for infinitely many  $n$ . In [2], Füredi et al. established the following results.

**Theorem 1** [2] *Let  $k, t$  be positive integers, where  $k \geq 3$ . For sufficiently large  $n$ , we have*

$$ex_k \left( n; \mathbb{P}_{2t+1}^{(k)} \right) = \binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{n-t}{k-1}.$$

*The only extremal family consists of all the  $k$ -sets in  $[n]$  that meet some fixed set  $S$  of  $t$  vertices. Also,*

$$ex_k \left( n; \mathbb{P}_{2t+2}^{(k)} \right) = \binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{n-t}{k-1} + \binom{n-t-2}{k-2}.$$

*The only extremal family consists of all the  $k$ -sets in  $[n]$  that meet some fixed set  $S$  of  $t$  vertices plus all the  $k$ -sets in  $[n] \setminus S$  that contain some two fixed elements.*

For more results we refer to [2, 6].

For the minimal and linear cycles, Füredi and Jiang [3], determined the extremal numbers when the forbidden hypergraph is a single minimal cycle or a single linear cycle. This confirms, in a stronger form, a conjecture of Mubayi and Verstraëte [6] for  $k \geq 5$  and adds to the limited list of hypergraphs whose Turán numbers have been known either exactly or asymptotically. Their main results are as follows:

**Theorem 2** [3] *Let  $t$  be a positive integer,  $k \geq 4$ . For sufficiently large  $n$ , we have  $ex_k(n; \mathcal{C}_{2t+1}^{(k)}) = \binom{n}{k} - \binom{n-t}{k}$ , and  $ex_k(n; \mathcal{C}_{2t+2}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + 1$ . For  $\mathcal{C}_{2t+1}^{(k)}$ , the only extremal family consists of all the  $k$ -sets in  $[n]$  that meet some fixed  $k$ -set  $S$ . For  $\mathcal{C}_{2t+2}^{(k)}$ , the only extremal family consists of all the  $k$ -sets in  $[n]$  that intersect some fixed  $t$ -set  $S$  plus one additional  $k$ -set outside  $S$ .*

**Theorem 3** [3] *Let  $t$  be a positive integer,  $k \geq 5$ . For sufficiently large  $n$ , we have  $ex_k(n; \mathcal{C}_{2t+1}^{(k)}) = \binom{n}{k} - \binom{n-t}{k}$ , and  $ex_k(n; \mathcal{C}_{2t+2}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + \binom{n-t-2}{k-2}$ . For  $\mathcal{C}_{2t+1}^{(k)}$ , the only extremal family consists of all the  $k$ -sets in  $[n]$  that meet some fixed  $k$ -set  $S$ . For  $\mathcal{C}_{2t+2}^{(k)}$ , the only extremal family consists of all the  $k$ -sets in  $[n]$  that intersect some fixed  $t$ -set  $S$  plus all the  $k$ -sets in  $[n] \setminus S$  that contain some two fixed elements.*

From the definition of  $k$ -uniform minimal cycles, two  $k$ -uniform minimal cycles of the same length may not be isomorphic. Hence we define the following family of hypergraphs, where every member consists of  $r$  vertex disjoint cycles:

$$\mathcal{S}(\ell_1, \dots, \ell_r) = \{C_1 \cup \dots \cup C_r : C_i \in \mathcal{C}_{\ell_i}^{(k)} \text{ for } i \in [r]\}$$

Apart from the results above, we will need the following two results:

**Theorem 4** [4] *Let  $H$  be a  $k$ -uniform hypergraph on  $n$  vertices with no two edges intersecting in exactly one vertex, where  $k \geq 3$ . Then  $|E(H)| \leq \binom{n}{k-2}$ .*

We build on earlier work of Füredi and Jiang [3], in this paper, we determine precisely the exact Turán numbers when forbidden hypergraphs are  $r$  vertex disjoint minimal cycles or  $r$  vertex disjoint linear cycles. Our main results are as follows:

**Theorem 5** *Let integers  $k \geq 4$ ,  $r \geq 1$ ,  $\ell_1, \dots, \ell_r \geq 3$ ,  $t = \sum_{i=1}^r \lfloor \frac{\ell_i+1}{2} \rfloor - 1$ , and  $I = 1$ , if all the  $\ell_1, \dots, \ell_r$  are even,  $I = 0$  otherwise. For sufficiently large  $n$ ,*

$$ex_k(n; \mathcal{S}(\ell_1, \dots, \ell_r)) = \binom{n}{k} - \binom{n-t}{k} + I.$$

**Theorem 6** Let integers  $k \geq 5$ ,  $r \geq 1$ ,  $\ell_1, \dots, \ell_r \geq 3$ ,  $t = \sum_{i=1}^r \lfloor \frac{\ell_i+1}{2} \rfloor - 1$ , and  $J = \binom{n-t-2}{k-2}$ , if all the  $\ell_1, \dots, \ell_r$  are even,  $J = 0$  otherwise. For sufficiently large  $n$ ,

$$ex_k(n; \mathcal{C}_{\ell_1}^{(k)}, \dots, \mathcal{C}_{\ell_r}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + J.$$

Sometimes, we allow the hypergraph to contain less than  $r$  minimal or linear cycles, consider the Turán number in such cases, we have the following two corollaries. We use notation  $r \cdot F$  to denote  $r$  vertex disjoint copies of hypergraph  $F$ . Let  $\ell_1 = \dots = \ell_r = \ell$ , we can immediately get the following two corollaries from Theorems 5 and 6.

**Corollary 1** Let integers  $k \geq 4$ ,  $r \geq 1$ ,  $\ell \geq 3$ ,  $t = r \lfloor \frac{\ell+1}{2} \rfloor - 1$ , and  $I = 1$ , if  $\ell$  is even,  $I = 0$ , if  $\ell$  is odd. For sufficiently large  $n$ ,

$$ex_k(n; r \cdot \mathcal{C}_{\ell}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + I.$$

**Corollary 2** Let integers  $k \geq 5$ ,  $r \geq 1$ ,  $\ell \geq 3$ ,  $t = r \lfloor \frac{\ell+1}{2} \rfloor - 1$ , and  $J = \binom{n-t-2}{k-2}$ , if  $\ell$  is even,  $J = 0$ , if  $\ell$  is odd. For sufficiently large  $n$ ,

$$ex_k(n; r \cdot \mathcal{C}_{\ell}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + J.$$

## 2 Proof of Theorem 5

For convenience, we define  $f(n, k, \{\ell_1, \dots, \ell_r\}) = \binom{n}{k} - \binom{n-t}{k} + I$ . Note that the hypergraph on  $n$  vertices that has every edge incident to some fixed  $t$ -set  $S$ , along with one additional edge disjoint from  $S$  when all of  $\ell_1, \dots, \ell_r$  are even, has exactly  $f(n, k, \{\ell_1, \dots, \ell_r\})$  edges and does not contain a copy of any member of  $\mathcal{S}(\ell_1, \dots, \ell_r)$ .

Thus, to prove Theorem 5, it suffices to prove that  $ex_k(n; \mathcal{S}(\ell_1, \dots, \ell_r)) \leq \binom{n}{k} - \binom{n-t}{k} + I$ , i.e., any hypergraph on  $n$  vertices with more than  $f(n, k, \{\ell_1, \dots, \ell_r\})$  edges must contain a member of  $\mathcal{S}(\ell_1, \dots, \ell_r)$ . We use induction on  $r$ . From Theorem

2, the case  $r = 1$  has been proved. Assume that  $r \geq 2$ , and Theorem 5 holds for smaller  $r$ .

Let  $H$  be a hypergraph on  $n$  vertices with  $m$  edges and  $m > f(n, k, \{\ell_1, \dots, \ell_r\})$ .

Since  $f(n, k, \{\ell_1, \dots, \ell_r\}) > f(n, k, \ell_1)$  for sufficiently large  $n$ , there exists at least one  $k$ -uniform minimal  $\ell_1$ -cycle in  $H$ . Take one of them, denote its vertex set by  $C$ , so  $\ell_1 \leq |C| \leq (k-1)\ell_1$ . We have that  $|E(H \setminus C)| \leq f(n - |C|, k, \{\ell_2, \dots, \ell_r\})$ , since otherwise, by induction hypothesis, we can find vertex disjoint copies of  $\mathcal{C}_{\ell_2}^{(k)} \cup \dots \cup \mathcal{C}_{\ell_r}^{(k)}$  in  $H$ ; plus the minimal  $\ell_1$ -cycle on  $C$ , there is a copy of a member of  $\mathcal{S}(\ell_1, \dots, \ell_r)$  in  $H$  already.

Let  $m_C$  denote the number of edges in  $H$  incident to vertices in  $C$ . Then,

$$m_C \geq m - f(n - |C|, k, \{\ell_2, \dots, \ell_r\}) \quad (1)$$

$$\geq f(n, k, \{\ell_1, \dots, \ell_r\}) - f(n - \ell_1, k, \{\ell_2, \dots, \ell_r\}) \quad (2)$$

$$= \frac{\lfloor \frac{\ell_1+1}{2} \rfloor}{(k-1)!} n^{k-1} + O(n^{k-2}). \quad (3)$$

We call an edge in  $H$  is a *terminal edge* if it contains exactly one vertex in  $C$ . Let  $T$  denote the set of all terminal edges in  $H$ . For every  $(k-1)$ -set  $R$  in  $V(H) \setminus C$ , define

$$T_R = \{E \in T : R \subseteq E\}.$$

According to the size of each set  $T_R$ , we partite all the  $(k-1)$ -sets in  $V(H) \setminus C$  into two sets, such that:

$$X = \{R \subseteq V(H) \setminus C \text{ and } |R| = k-1 : |T_R| \leq \lfloor \frac{\ell_1+1}{2} \rfloor - 1\}$$

$$Y = \{R \subseteq V(H) \setminus C \text{ and } |R| = k-1 : |T_R| \geq \lfloor \frac{\ell_1+1}{2} \rfloor\}.$$

It is not difficult to give an upper bound of  $m_C$  with the terms  $|X|$  and  $|Y|$  as follows:

$$\begin{aligned} m_C &\leq \binom{|C|}{2} \binom{n-2}{k-2} + |X| \left( \lfloor \frac{\ell_1+1}{2} \rfloor - 1 \right) + |Y| \cdot |C| \\ &\leq \binom{|C|}{2} \binom{n-2}{k-2} + \binom{n}{k-1} \left( \lfloor \frac{\ell_1+1}{2} \rfloor - 1 \right) + |Y| \cdot \ell_1 (k-1). \end{aligned}$$

Combine with (3), we have that

$$|Y| \geq \frac{n^{k-1}}{(k-1)\ell_1(k-1)!} + O(n^{k-2}). \quad (4)$$

For any  $(k-1)$ -set  $R \in Y$ , there are at least  $\lfloor \frac{\ell_1+1}{2} \rfloor$  vertices in  $C$  that can form terminal edges with  $R$ . We choose exactly  $\lfloor \frac{\ell_1+1}{2} \rfloor$  of them, call the vertex set of these

$\lfloor \frac{\ell_1+1}{2} \rfloor$  vertices *terminal set* relative to  $R$ . Since the number of  $\lfloor \frac{\ell_1+1}{2} \rfloor$ -sets in  $C$  is at most  $\binom{|C|}{\lfloor \frac{\ell_1+1}{2} \rfloor}$ , we can get that some elements in  $Y$  may have the same terminal set. And it is easy to derive that, the number of  $(k-1)$ -sets in  $Y$  with the same terminal set, is at least

$$\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \binom{|C|}{\lfloor \frac{\ell_1+1}{2} \rfloor}^{-1} + O(n^{k-2}) \geq \frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor}^{-1} + O(n^{k-2}).$$

Choose one terminal set  $U$  in  $C$ , such that there are at least  $\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor}^{-1} + O(n^{k-2})$   $(k-1)$ -sets in  $V(H) \setminus C$ , every such  $(k-1)$ -set can form a terminal edge with every vertex in  $U$ . Let  $R_U$  be the set of all the common  $(k-1)$ -sets associate with  $U$  in  $V(H) \setminus C$ , we have that

$$|R_U| \geq \frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor}^{-1} + O(n^{k-2}). \quad (5)$$

Let  $m_U$  denote the number of edges incident to vertices in  $U$ , then,

$$m_U \leq \binom{\lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} + m',$$

where  $m'$  is the number of edges which contain at least two vertices in  $U$ . With some calculations, we have that

$$\begin{aligned} & f(n, k, \{\ell_1, \dots, \ell_r\}) - f\left(n - \left\lfloor \frac{\ell_1+1}{2} \right\rfloor, k, \{\ell_2, \dots, \ell_r\}\right) - m_U \\ &= \binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} - m_U \\ &\geq \left[ \binom{n-1}{k-1} - \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} \right] + \left[ \binom{n-2}{k-1} - \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} \right] \\ &\quad + \dots + \left[ \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor + 1}{k-1} - \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} \right] - m'. \end{aligned}$$

It is not difficult to deduce that the last expression is no less than zero (consider the combinatorial meaning of that expression), hence, we can derive that

$$\begin{aligned} E(H \setminus U) &= m - m_U > f(n, k, \{\ell_1, \dots, \ell_r\}) - m_U \\ &\geq f\left(n - \left\lfloor \frac{\ell_1+1}{2} \right\rfloor, k, \{\ell_2, \dots, \ell_r\}\right). \end{aligned}$$

Thus by the induction hypothesis, there exists a member of  $\mathcal{S}(\ell_2, \dots, \ell_r)$  with vertex set  $W$  in  $V(H) \setminus U$ , also we have that

$$|W| \leq (k-1) \sum_{i=2}^r \ell_i. \quad (6)$$

Now we focus on finding a  $k$ -uniform minimal  $\ell_1$ -cycle disjoint from  $W$ .

Considering the  $(k-1)$ -uniform hypergraph  $H_0$  with vertex set  $V(H) \setminus U$  and edge set  $R_U$ , we will prove the following claim:

**Claim 1** *There are  $\lfloor \frac{\ell_1}{2} \rfloor$  pairs of  $(k-1)$ -edges in  $H_0$ , say  $\{a_i, b_i\}$ ,  $i = 1, \dots, \lfloor \frac{\ell_1}{2} \rfloor$ , such that for every  $i$ ,  $a_i$  and  $b_i$  have exactly one common vertex, and for any  $j \neq i$ ,  $\{a_i, b_i\}$  and  $\{a_j, b_j\}$  are vertex disjoint, moreover, all these  $(k-1)$ -edges disjoint from  $W$ .*

*Proof.* The number of  $(k-1)$ -edges incident with vertices in  $W$  is at most  $|W| \cdot \binom{n-1}{k-2}$ . With the aid of (5) and (6), in  $R_U$ , the number of  $(k-1)$ -edges disjoint from  $W$  is at least

$$\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left( \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2}) - (k-1) \sum_{i=2}^r \ell_i \binom{n-1}{k-2} > \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-2}.$$

By Theorem 4, we can find a pair of  $(k-1)$ -edges  $\{a_1, b_1\}$  with exactly one common vertex. Let  $p = \lfloor \frac{\ell_1}{2} \rfloor (2k-3)$ , since  $\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left( \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2}) - (k-1) \sum_{i=2}^r \ell_i \binom{n-1}{k-2} - p \binom{n-1}{k-2} > \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-2}$ , we can repeat the argument above to find  $\{a_2, b_2\}, \dots, \{a_{\lfloor \frac{\ell_1}{2} \rfloor}, b_{\lfloor \frac{\ell_1}{2} \rfloor}\}$  satisfying the properties described in Claim 1.  $\square$

Let  $U = \{u_1, \dots, u_{\lfloor \frac{\ell_1+1}{2} \rfloor}\}$ . To form the required minimal  $\ell_1$ -cycle, we need to consider such two cases:

**Case 1.**  $\ell_1$  is even.

Find  $\frac{\ell_1}{2}$  pairs of  $(k-1)$ -edges in  $H_0$  as described in Claim 1, still denote them by  $\{a_i, b_i\}$ ,  $i = 1, \dots, \frac{\ell_1}{2}$ . Construct a  $k$ -uniform minimal  $\ell_1$ -cycle in  $H$  with edges:

$$a_1 \cup \{u_1\}, b_1 \cup \{u_2\}, a_2 \cup \{u_2\}, \dots, b_{\frac{\ell_1}{2}-1} \cup \{u_{\frac{\ell_1}{2}}\}, a_{\frac{\ell_1}{2}} \cup \{u_{\frac{\ell_1}{2}}\}, b_{\frac{\ell_1}{2}} \cup \{u_1\}.$$

**Case 2.**  $\ell_1$  is odd.

Find  $\frac{\ell_1-3}{2}$  pairs of  $(k-1)$ -edges in  $H_0$  as described in Claim 1. Similar to the proof of Claim 1. Let  $Q$  be the vertex set of  $W$  and all these  $\frac{\ell_1-3}{2}$  pairs of  $(k-1)$ -edges, hence,  $|Q| = \frac{\ell_1-3}{2}(2k-3) + |W|$ . By Theorem 1,  $ex_{k-1} \left( n - \lfloor \frac{\ell_1+1}{2} \rfloor; \mathbb{P}_3^{(k-1)} \right) = \frac{1}{(k-2)!} n^{k-2} + O(n^{k-3})$ , for sufficiently large  $n$ . In  $H_0$ , the number of  $(k-1)$ -edges disjoint from  $Q$  is at least  $\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left( \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2}) - |Q| \binom{n-1}{k-2} > \frac{1}{(k-2)!} n^{k-2} + O(n^{k-3})$ . That implies in  $H_0$ , we can find a  $\mathbb{P}_3^{(k-1)}$  in remaining  $(k-1)$ -edges disjoint from  $Q$ . Let  $x, y, z$  be the three consecutive  $(k-1)$ -edges in that  $\mathbb{P}_3^{(k-1)}$ , then, in  $H$ , we can form a  $k$ -uniform minimal  $\ell_1$ -cycle with edges:

$$a_1 \cup \{u_1\}, b_1 \cup \{u_2\}, a_2 \cup \{u_2\}, \dots, a_{\frac{\ell_1-3}{2}} \cup \{u_{\frac{\ell_1-3}{2}}\},$$

$$b_{\frac{\ell_1-3}{2}} \cup \{u_{\frac{\ell_1-1}{2}}\}, x \cup \{u_{\frac{\ell_1-1}{2}}\}, y \cup \{u_{\frac{\ell_1+1}{2}}\}, z \cup \{u_1\}.$$

Moreover, it is easy to see that this  $k$ -uniform minimal  $\ell_1$ -cycle is not only minimal, but also linear, whenever  $\ell_1$  is even or odd. Thus, we have constructed  $r$  disjoint  $k$ -uniform minimal cycles. So the hypergraph which contains no member of  $\mathcal{S}(\ell_1, \dots, \ell_r)$  can not have more than  $f(n, k, \{\ell_1, \dots, \ell_r\})$  edges. Thus completes the proof.  $\blacksquare$

### 3 Proof of Theorem 6

The argument in the proof of Theorem 6 is similar to the proof of Theorem 5.

Let  $g(n, k, \{\ell_1, \dots, \ell_r\}) = \binom{n}{k} - \binom{n-t}{k} + J$ . Firstly, we point out that the hypergraph on  $n$  vertices that has every edge incident to some fixed  $t$ -set  $S$ , along with all the  $k$ -edges disjoint from  $S$  containing some two fixed elements not in  $S$  when all of  $\ell_1, \dots, \ell_r$  are even, has exactly  $g(n, k, \{\ell_1, \dots, \ell_r\})$  edges and dose not contain a copy of any member of  $\mathbb{C}_{\ell_1}^{(k)} \cup \dots \cup \mathbb{C}_{\ell_r}^{(k)}$ .

Hence it suffices to prove that  $ex_k(n; \mathbb{C}_{\ell_1}^{(k)}, \dots, \mathbb{C}_{\ell_r}^{(k)}) \leq g(n, k, \{\ell_1, \dots, \ell_r\})$ . We proceed by induction on  $r$  again since the case  $r = 1$  is provided by Theorem 3. Let  $H$  be a hypergraph on  $n$  vertices with  $m > g(n, k, \{\ell_1, \dots, \ell_r\})$  edges. If one of  $\ell_1, \dots, \ell_r$  is even, rearrange the sequence to make sure  $\ell_1$  is even.

As in the proof of Theorem 5, since  $g(n, k, \{\ell_1, \dots, \ell_r\}) > g(n, k, \ell_1)$  for sufficiently large  $n$ , there exists at least one  $k$ -uniform linear  $\ell_1$ -cycle in  $H$ . Take one of them, denote its vertex set by  $C$ . Similarly, we have that  $|E(H \setminus C)| \leq g(n - |C|, k, \{\ell_2, \dots, \ell_r\})$ . Still let  $m_C$  denote the number of edges in  $H$  incident to vertices in  $C$ , with some calculations, we can get that:

$$m_C \geq \frac{\lfloor \frac{\ell_1+1}{2} \rfloor}{(k-1)!} n^{k-1} + O(n^{k-2}).$$

Again we define terminal edges,  $T_R, X, Y$  as before, we can find the  $\lfloor \frac{\ell_1+1}{2} \rfloor$ -set  $U$ , too. Then by induction hypothesis, we can find a copy of  $\mathbb{C}_{\ell_2}^{(k)} \cup \dots \cup \mathbb{C}_{\ell_r}^{(k)}$  on vertex set  $W$  in  $V(H) \setminus U$ . With the same method used in the proof of Theorem 5, we can select a terminal set of size  $\lfloor \frac{\ell_1+1}{2} \rfloor$  in  $C$ , then, similarly, we can construct a  $k$ -uniform linear  $\ell_1$ -cycle in  $H$  since the  $k$ -uniform minimal  $\ell_1$ -cycle we described in the proof of Theorem 5 is also linear. And this  $k$ -uniform linear  $\ell_1$ -cycle avoid the vertices in  $W$ , hence we know that the hypergraph which contains no  $\mathbb{C}_{\ell_1}^{(k)} \cup \dots \cup \mathbb{C}_{\ell_r}^{(k)}$  can not have more than  $g(n, k, \{\ell_1, \dots, \ell_r\})$  edges. The proof is thus complete.  $\blacksquare$



## References

- [1] C. Berge, *Hypergraphs*, North-Holland Mathematical Library 45, North-Holland, Amsterdam, 1989.
- [2] Z. Füredi, T. Jiang, R. Seiver, Exact Solution of the hypergraph Turán problem for  $k$ -uniform linear paths, *Combinatorica*, to appear.
- [3] Z. Füredi, T. Jiang, Hypergraph Turán numbers of linear cycles, arXiv:1302.2387. [math.CO] 2013.
- [4] P. Keevash, D. Mubayi, R. Wilson, Set systems with no singleton intersection, *SIAM J. Discrete Math.* **20** (2006), 1031–1041.
- [5] E. Györi, G. Katona, N. Lemons, Hypergraph extensions of the Erdős-Gallai theorem, *Electronic Notes in Disc. Math.* **36** (2010), 655–662.
- [6] D. Mubayi, J. Verstraëte: Minimal paths and cycles in set systems, *European J. Combin.* **28**(2007), 1681–1693.