

A PROOF OF XIN–ZHANG’S TRIDIAGONAL DETERMINANT CONJECTURE

JIAQIANG HU  and CHEN ZHANG  

(Received 11 April 2026 UTC; accepted 30 June 2026 UTC)

Abstract

We confirm a recent conjecture of Xin and Zhang [‘A variation of the Morris constant term’, Preprint, 2024, arXiv:2409.14356v2], which establishes a simple product formula for the characteristic polynomial of an $(n-1) \times (n-1)$ tridiagonal matrix. This characteristic polynomial arises from a recurrence relation that enumerates $n \times n$ nonnegative integer matrices with all row and column sums equal to t , also called the Ehrhart polynomial of the n th Birkhoff polytope.

2020 *Mathematics subject classification*: primary 15A15; secondary 05A10, 05A19.

Keywords and phrases: determinant, tridiagonal matrix, characteristic polynomial, combinatorial identity, Pascal’s triangle.

1. Introduction

Our main objective is to prove a conjecture of Xin and Zhang [13], which gives a product formula for the characteristic polynomial of a certain tridiagonal matrix.

CONJECTURE 1.1 [13, Conjecture 4.8]. Let n be a positive integer and let C be the $(n-1) \times (n-1)$ matrix defined by

$$C_{ij} = \begin{cases} t - (n - 2i + 2)(n - 1 - i) - 1 & \text{if } i = j, \\ (i - 1)(n - i) & \text{if } i = j + 1, \\ -(n - 1 - i)(n - i) & \text{if } i = j - 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then, $\det C$ is equal to

$$\begin{cases} (t - n + 1) \prod_{i=0}^{r-2} (t - n + 1 - r(r-1) + i(i+1))^2 & \text{if } n = 2r, \\ (t - n + 1)(t - n + 1 - (r-1)^2) \prod_{i=1}^{r-2} (t - n + 1 - (r-1)^2 + i^2)^2 & \text{if } n = 2r - 1. \end{cases}$$

This work was supported by the Postdoctoral Fellowship Program and China Postdoctoral Science Foundation (No. BX20250066).

© The Author(s), 2026. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

This conjecture arises in the study of the Ehrhart polynomial $H_n(t)$ of the n th Birkhoff polytope, where $H_n(t)$ counts the number of $n \times n$ nonnegative integer matrices with all row and column sums equal to t . The computation of $H_n(t)$ has been a subject of extensive research (see, for example, [1, 5, 6, 8–11, 14]). So far, $H_n(t)$ is known only for $n \leq 9$, with $H_9(t)$ computed by Beck and Pixton [3] using residue techniques.

From [3, Corollary 4],

$$H_n(t) = \sum_{\mathbf{m}} \binom{n}{\mathbf{m}} \text{CT}_{\mathbf{x}} \mathcal{H}^{\mathbf{m}}(t),$$

where the sum is over all nonnegative integer sequences $\mathbf{m} = (m_1, m_2, \dots, m_n)$ such that $\sum_{i=1}^n m_i = n$ and $\sum_{i=1}^r m_i > r$ for $1 \leq r < n$. Here, $\text{CT}_{\mathbf{x}} f$ denotes the constant term of the Laurent series f in $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and

$$\mathcal{H}^{\mathbf{m}}(t) := \prod_{i=1}^n \frac{x_i^{(m_i-1)t}}{\prod_{j=1, j \neq i}^n (1 - x_j/x_i)^{m_i}}.$$

In particular, for $\mathbf{m} = (2, 1^{n-2}, 0)$, we define

$$\begin{aligned} h_n(t) &:= (-1)^{(n-1)(n-2)/2} \text{CT}_{\mathbf{x}} \mathcal{H}^{(2, 1^{n-2}, 0)}(t) \\ &= \text{CT}_{\mathbf{x}} \frac{(x_n/x_1)^{-t} \prod_{i=2}^{n-1} (x_i/x_1)^{2i-n} \prod_{i=1}^{n-1} (1 - x_n/x_i)}{\prod_{i=2}^n (1 - x_i/x_1)^3 \prod_{2 \leq i < j \leq n} (1 - x_j/x_i)^2}. \end{aligned}$$

Noting the similarity with the Morris constant term identity studied in [2, 12], Xin and Zhang [13] derived a recurrence for $h_n(t)$ of the form

$$h_n(t) = -\frac{1}{c_0(t+n-1)} \sum_{i=1}^{n-1} c_i(t+n-1)h_n(t-i), \quad (1.1)$$

in which $c_0(t) = \det C$. Therefore, if Conjecture 1.1 holds, then (1.1) indeed yields a recurrence when

$$t > \left\lfloor \frac{(n-1)^2}{4} \right\rfloor = \begin{cases} r(r-1) & \text{if } n = 2r, \\ (r-1)^2 & \text{if } n = 2r-1. \end{cases}$$

The ideas in [13] also apply to $\text{CT}_{\mathbf{x}} \mathcal{H}^{(k, 1^{n-k}, 0^{k-1})}(t)$ for $3 \leq k \leq n-1$. Moreover, the determinant $\det C$ arises in a similar but more complicated recurrence. These aspects will be elaborated on in an upcoming paper [15].

Further developments motivate us to consider the equivalent determinant of

$$tI - \widetilde{C}^{(n-1)} = C + (n-1)I,$$

where I is the identity matrix and $\widetilde{C}^{(n)}$ is an $n \times n$ tridiagonal matrix with entries

$$(\widetilde{C}^{(n)})_{i,j} = \begin{cases} (n+1-i)(n+1-2i) & \text{if } i=j, \\ (n+1-i)(1-i) & \text{if } i=j+1, \\ (n+1-i)(n-i) & \text{if } i=j-1, \\ 0 & \text{otherwise.} \end{cases} \quad (1.2)$$

It is hard to determine the characteristic polynomial of $\widetilde{C}^{(n)}$: none of the minors of $tI - \widetilde{C}^{(n)}$ admits a simple factorisation. This prevents us from proving the conjecture by traditional techniques such as diagonalisation, LU decompositions, continued fractions and so on. Ultimately, we discovered that $\widetilde{C}^{(n)}$ is similar to a lower triangular matrix. It is noteworthy that the similar matrix is a variation of Pascal’s triangle. We state this result as the following theorem.

THEOREM 1.2. *For a positive integer n , let $\widetilde{C}^{(n)}$ be the matrix defined in (1.2). There exists an $n \times n$ nonsingular upper triangular matrix U with entries*

$$U_{i,j} = \binom{n-i}{n-j}, \quad (1.3)$$

such that $U\widetilde{C}^{(n)}U^{-1}$ is a lower triangular matrix whose entries are given by

$$(U\widetilde{C}^{(n)}U^{-1})_{i,j} = \begin{cases} i(n-i) & \text{if } i=j, \\ (n+1-i)(1-i) & \text{if } i=j+1, \\ 0 & \text{otherwise.} \end{cases}$$

Consequently, Conjecture 1.1 holds and we have the nice formula

$$\det C = \prod_{i=1}^{n-1} (t - n + 1 - i(n-1-i)).$$

For instance,

$$\widetilde{C}^{(4)} = \begin{pmatrix} 12 & 12 & 0 & 0 \\ -3 & 3 & 6 & 0 \\ 0 & -4 & -2 & 2 \\ 0 & 0 & -3 & -3 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & 3 & 3 & 1 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$U\widetilde{C}^{(4)}U^{-1} = \begin{pmatrix} 3 & 0 & 0 & 0 \\ -3 & 4 & 0 & 0 \\ 0 & -4 & 3 & 0 \\ 0 & 0 & -3 & 0 \end{pmatrix}.$$

In Section 2, we prove Theorem 1.2 using an explicit formula for U^{-1} , together with several combinatorial identities. Section 3 contains concluding remarks.

2. The proof of Theorem 1.2

Our proof of Theorem 1.2 is based on the following well-known combinatorial identities (the proofs are omitted):

$$\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}, \quad (2.1)$$

$$n \binom{n-1}{k-1} = k \binom{n}{k} = (n-k+1) \binom{n}{k-1}, \quad (2.2)$$

$$\binom{n}{k} \binom{k}{j} = \binom{n}{j} \binom{n-j}{k-j}, \quad (2.3)$$

where $n \geq k \geq j$ are nonnegative integers.

PROOF OF THEOREM 1.2. By [4, Theorem 1], the entries of U^{-1} are given by

$$(U^{-1})_{i,j} = (-1)^{i+j} U_{i,j} = (-1)^{i+j} \binom{n-i}{n-j}.$$

Consider the (i, j) -entry of $U\widetilde{C}^{(n)}U^{-1}$:

$$\begin{aligned} (U\widetilde{C}^{(n)}U^{-1})_{i,j} &= \sum_{1 \leq l, m \leq n} U_{i,l}(\widetilde{C}^{(n)})_{l,m}(U^{-1})_{m,j} \\ &= \sum_{l=1}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l}(U^{-1})_{l,j} + \sum_{l=2}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l-1}(U^{-1})_{l-1,j} \\ &\quad + \sum_{l=1}^{n-1} U_{i,l}(\widetilde{C}^{(n)})_{l,l+1}(U^{-1})_{l+1,j}, \end{aligned}$$

where the second equality holds because $(\widetilde{C}^{(n)})_{l,m} = 0$ when $|l-m| > 1$. Observe that $(\widetilde{C}^{(n)})_{l,l} = (\widetilde{C}^{(n)})_{l,l-1} + (\widetilde{C}^{(n)})_{l,l+1}$, and the formulae for $(\widetilde{C}^{(n)})_{l,l-1}$ and $(\widetilde{C}^{(n)})_{l,l+1}$ vanish when $l = 1$ and $l = n$, respectively. The computation is naturally divided into two parts, denoted by S_1 and S_2 . In detail,

$$\begin{aligned} S_1 &= \sum_{l=1}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l-1}(U^{-1})_{l,j} + \sum_{l=2}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l-1}(U^{-1})_{l-1,j} \\ &= \sum_{l=1}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l-1}((U^{-1})_{l,j} + (U^{-1})_{l-1,j}) \\ &= \sum_{l=1}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l-1}(-1)^{j+l-1} \left(\binom{n-l+1}{n-j} - \binom{n-l}{n-j} \right) \\ &= \sum_{l=1}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l-1}(-1)^{j+l-1} \binom{n-l}{n-j-1} \quad (\text{by (2.1)}). \end{aligned}$$

Similarly,

$$\begin{aligned} S_2 &= \sum_{l=1}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l+1}(U^{-1})_{l,j} + \sum_{l=1}^{n-1} U_{i,l}(\widetilde{C}^{(n)})_{l,l+1}(U^{-1})_{l+1,j} \\ &= \sum_{l=1}^n U_{i,l}(\widetilde{C}^{(n)})_{l,l+1}(-1)^{j+l} \binom{n-l-1}{n-j-1}. \end{aligned}$$

Using (2.2), we can further compute S_2 :

$$\begin{aligned} S_2 &= \sum_{l=1}^n U_{i,l}(-1)^{j+l}(n+1-l)(n-l) \binom{n-l-1}{n-j-1} \\ &= \sum_{l=1}^n U_{i,l}(-1)^{j+l}(n+1-l)(j-l+1) \binom{n-l}{n-j-1}. \end{aligned}$$

Then,

$$\begin{aligned} (U\widetilde{C}^{(n)}U^{-1})_{i,j} &= S_1 + S_2 = j \sum_{l=1}^n (-1)^{j+l} \binom{n-i}{n-l} \binom{n-l}{n-j-1} (n+1-l) \\ &= j \sum_{l=1}^n (-1)^{j+l} \binom{n-i}{n-j-1} \binom{j+1-i}{j+1-l} (n+1-l) \quad (\text{by (2.3)}). \end{aligned}$$

It is clear that the above formula equals 0 when $i > j+1$ and equals $(1-i)(n+1-i)$ when $i = j+1$. When $i \leq j$, we can further write

$$\begin{aligned} (U\widetilde{C}^{(n)}U^{-1})_{i,j} &= j \sum_{l=i}^{j+1} (-1)^{j+l} \binom{n-i}{n-j-1} \binom{j+1-i}{j+1-l} ((n-j) + (j+1-l)) \\ &= -j(n-j) \binom{n-i}{n-j-1} \sum_{l=i}^{j+1} \binom{j+1-i}{j+1-l} (-1)^{j+1-l} \\ &\quad + j(j+1-i) \binom{n-i}{n-j-1} \sum_{l=i}^j \binom{j-i}{j-l} (-1)^{j-l} \quad (\text{by (2.2)}) \\ &= \binom{n-i}{n-j-1} (-j(n-j)(1+(-1))^{j+1-i} + j(j+1-i)(1+(-1))^{j-i}) \\ &= \begin{cases} i(n-i) & \text{if } i = j, \\ 0 & \text{if } i < j. \end{cases} \end{aligned}$$

This completes the proof. $\square$

As a byproduct, we have the following result.

PROPOSITION 2.1. *Let U be the matrix with entries as in (1.3) and M be a matrix satisfying $M_{i,j} = 0$ whenever $i - j \geq 2$. Then, $(UMU^{-1})_{i,j} = M_{i,j}$ whenever $i - j \geq 1$.*

Proposition 2.1 can be visualised as

$$U \begin{pmatrix} * & * & * & \cdots & * \\ M_{2,1} & * & * & \cdots & * \\ 0 & M_{3,2} & * & \cdots & * \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & M_{n,n-1} & * \end{pmatrix} U^{-1} = \begin{pmatrix} \Delta & \Delta & \Delta & \cdots & \Delta \\ M_{2,1} & \Delta & \Delta & \cdots & \Delta \\ 0 & M_{3,2} & \Delta & \cdots & \Delta \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & M_{n,n-1} & \Delta \end{pmatrix},$$

where ‘*’ and ‘ Δ ’ stand for unspecified entries.

PROOF OF PROPOSITION 2.1. Consider the (i, j) -entry of UMU^{-1} :

$$(UMU^{-1})_{i,j} = \sum_{1 \leq l, m \leq n} U_{i,l} M_{l,m} (U^{-1})_{m,j} = \sum_{1 \leq l, m \leq n} (-1)^{m+j} \binom{n-i}{n-l} \binom{n-m}{n-j} M_{l,m}.$$

For fixed i and j , we have $\binom{n-i}{n-l} \binom{n-m}{n-j} = 0$ unless $l \geq i$ and $m \leq j$.

In the case where $i - j \geq 2$, the condition $l \geq i$ and $m \leq j$ implies that $l - m \geq i - j \geq 2$. Hence, $M_{l,m} = 0$ and so $(UMU^{-1})_{i,j} = 0 = M_{i,j}$.

In the case where $i - j = 1$,

$$(-1)^{m+i-1} \binom{n-i}{n-l} \binom{n-m}{n-i+1} M_{l,m} = \begin{cases} M_{i,i-1} & \text{if } l = i \text{ and } m = i - 1, \\ 0 & \text{otherwise.} \end{cases}$$

Hence, $(UMU^{-1})_{i,i-1} = M_{i,i-1}$. This completes the proof. $\square$

3. Concluding remarks

The observation $(\widetilde{C}^{(n)})_{i,i} = (\widetilde{C}^{(n)})_{i,i-1} + (\widetilde{C}^{(n)})_{i,i+1}$ leads us to consider a matrix A satisfying $A_{i,i} = r + A_{i,i-1} + A_{i,i+1}$, where r is a parameter. Further exploration gives a generalisation of Theorem 1.2, which requires a cumbersome condition. We elaborate on this generalisation in the extended version [7, Section 3].

Acknowledgements

The authors would like to thank Professors Guoce Xin (Capital Normal University) and Arthur L. B. Yang (Nankai University) for helpful discussions.

References

- [1] P. Alexandersson, S. Hopkins and G. Zaimi, ‘Restricted Birkhoff polytopes and Ehrhart period collapse’, *Discrete Comput. Geom.* **73** (2025), 62–78.
- [2] W. Baldoni-Silva and M. Vergne, ‘Residues formulae for volumes and Ehrhart polynomials of convex polytopes’, Preprint, 2001, [arXiv:math/0103097v1](https://arxiv.org/abs/math/0103097v1).
- [3] M. Beck and D. Pixton, ‘The Ehrhart polynomial of the Birkhoff polytope’, *Discrete Comput. Geom.* **30** (2003), 623–637.
- [4] G. S. Call and D. J. Velleman, ‘Pascal’s matrices’, *Amer. Math. Monthly* **100** (1993), 372–376.
- [5] C. S. Chan and D. P. Robbins, ‘On the volume of the polytope of doubly stochastic matrices’, *Exp. Math.* **8** (1999), 291–300.

- [6] J. A. De Loera, F. Liu and R. Yoshida, ‘A generating function for all semi-magic squares and the volume of the Birkhoff polytope’, *J. Algebraic. Combin.* **30** (2009), 113–139.
- [7] J. Hu and C. Zhang, ‘A proof of Xin–Zhang’s tridiagonal determinant conjecture (extended version)’, Preprint, 2026, [arXiv:2601.03082](https://arxiv.org/abs/2601.03082).
- [8] P. A. MacMahon, *Combinatory Analysis* (Chelsea Publishing Co., New York, 1960).
- [9] J. Mount, ‘Fast unimodular counting’, *Combin. Probab. Comput.* **9** (2000), 277–285.
- [10] R. P. Stanley, ‘Linear homogeneous Diophantine equations and magic labelings of graphs’, *Duke Math. J.* **40** (1973), 607–632.
- [11] R. P. Stanley, ‘Magic labelings of graphs, symmetric magic squares, systems of parameters, and Cohen-Macaulay rings’, *Duke Math. J.* **43** (1976), 511–531.
- [12] G. Xin, ‘The Ring of Malcev–Neumann series and the residue theorem’, PhD Thesis, Brandeis University, Preprint, 2004, [arXiv:math.CO/0405133](https://arxiv.org/abs/math.CO/0405133).
- [13] G. Xin and C. Zhang, ‘A variation of the Morris constant term’, Preprint, 2024, [arXiv:2409.14356v2](https://arxiv.org/abs/2409.14356v2).
- [14] G. Xin, C. Zhang, Y. Zhou and Y. Zhong, ‘The constant term algebra of type A: The structure’, *Adv. Math.* **465** (2025), Article no. 110154.
- [15] C. Zhang, ‘The constant term of the hook shape arising from the Birkhoff polytope’, in preparation.

JIAQIANG HU, School of Mathematical Sciences,
LPMC, Nankai University, Tianjin 300071, PR China
e-mail: 2210472@mail.nankai.edu.cn

CHEN ZHANG, Center for Combinatorics,
LPMC, Nankai University, Tianjin 300071, PR China
e-mail: ch_enz@163.com